

## 27. Verdugo Mountains SEA

### *Location*

#### *General*

The Verdugo Mountains Significant Ecological Area (SEA) is located within the Verdugo Mountains. This SEA encompasses the Verdugo Mountains south of Interstate-210 and east of the Interstate-5, as well as a portion of these mountains north of Interstate-210.

The Verdugo Mountains are a wilderness island in the middle of the urbanized metropolitan area of the County, surrounded by the cities of Los Angeles, Burbank and Glendale. This area is cherished by the local communities, much of which are designated agricultural with many equestrian properties. The Verdugo Mountains have retained a rural atmosphere despite their proximity to urban Los Angeles. The Verdugo Mountains currently encompass wilderness area, which ranges through various chaparral, coastal sage chaparral scrub, southern willow scrub, coast live oak woodland and forest ecosystems, and many riparian areas with seasonal waterfalls. It is one of the few remaining natural regions in the Los Angeles area that supports abundant native wildlife and habitats, and also contains several rare and sensitive plant and animal species. The geographic location of the Verdugo Mountains makes them important for scientific study, genetic interchange between otherwise isolated populations, and recreation for urban residents.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Burbank, Sunland, and Pasadena.

#### *General Boundary and Resources Description*

The SEA is an island refuge, providing what remains of a link between plant and animal populations found in the Santa Monica and San Gabriel mountains. Genetic interchange, by way of this linkage is important in perpetuating the genetic variability in isolated populations, and the maintenance of healthy ecosystems. Chaparral and coastal sage chaparral scrub cover the hillsides of the mountains, with lush riparian vegetation, including California bay (*Umbellularia californica*), western sycamore (*Platanus racemosa*), coast live oak (*Quercus agrifolia* var. *agrifolia*), ferns, and ocellated Humboldt lily (*Lilium humboldtii* ssp. *ocellatum*), which are found in most of the stream drainages. These plant communities provide habitat that is essential to the diverse and abundant fauna that are found in the area. The mountains are also home to the northernmost population of mission manzanita (*Xylococcus bicolor*).

The proximity of the mountains to urban areas provides an excellent opportunity to study the interaction between wild animal populations and humans. The area has already been used for studies concerned with public health. The Interstate-210 crosses the northern edge and there is some intrusion of development along La Tuna Canyon Road. Residential development is beginning to trim the area and has been excluded from the SEA. However, present human use of the natural, native area has been low and has not significantly affected the natural resources found in the Verdugo Mountains.

The northernmost point of the SEA is in the Shadow Hills district of the City of Los Angeles at the west side of the undercrossing of Sunland/Foothill Boulevard beneath the Interstate-210. The freeway frontage is native vegetation and connective to the north with the Wentworth Street undercrossing of Interstate-210. The connection is through native vegetation along the freeway frontage and among dispersed equestrian ranchettes in the Shadow Hills. Wentworth Street ends a short distance east of Interstate-210 at the Angeles National Golf Club, which has its greens interspersed among braided stream courses of the Big Tujunga Canyon alluvial fan. The golf course

is part of the SEA. The native vegetation along the freeway, at the restored County Public Works area on the north side of Wentworth, and the native vegetation among the residences of Shadow Hills is all considered an important conduit for wildlife traversing among the San Gabriel Mountains, the SEA, and the Verdugo Mountains.

The SEA boundary follows southward along the southwest side of the Interstate-210 for about a half mile and then crosses to the northeast side of Interstate-210, along a ridge that connects to slopes with natural, native habitat that continues east, along the southern border of the communities of Sunland and Tujunga. In the vicinity of Pasko Peak, the SEA border is drawn around development in the southern edge of the community of Tujunga. The SEA border recrosses the Interstate-210 and then La Tuna Canyon Road. The crossing boundary goes along the east side of a northern tributary of La Tuna Canyon.

A paved road goes along the stream course in La Tuna Canyon, and the SEA includes about two miles of native, natural vegetation bordered on the eastern, undeveloped section of the road. The developed western end of La Tuna Canyon Road is excluded from the SEA in a "cherry stem" configuration.

After crossing Interstate-210 and La Tuna Canyon Road, the SEA tracks the southern side of natural habitat along the Interstate-210 for about a mile. The boundary trends irregularly southeast excluding development in the flatter areas that border the Verdugo Wash in the Verdugo City District of Glendale. The SEA includes the incised canyons of Sheep Corral, Cunningham, Henderson, Engleheard, and unnamed others, which are all tributaries of the channelized Verdugo Wash.

In the relatively narrowed area around the State Route-2 (Verdugo Canyon) between the San Rafael Hills on the west and the Verdugo Mountains on the East, the SEA continues irregularly south along the natural, native vegetation of the steep hillsides of the Verdugo Mountains that border Glendale. The SEA includes many unnamed canyons and also Ayars Canyon and Deer Creek at the end of Beaudry Boulevard. The unnamed canyon north of Dead Horse Canyon has a possible wildlands connection with the San Rafael Hills. Its northeast-facing slope along Sunshine Drive has chiefly native vegetation and ends in the Verdugo Park on Verdugo Boulevard. Across Verdugo Boulevard is Glendale Community College. The College's eastern border is a natural ridge that connects with the Mountain Avenue overpass of State Route-2 and native vegetation of the San Rafael Hills. Verdugo Boulevard and its development is a substantial block to terrestrial wildlife movement, but aerial fauna and plant seeds can connect fairly easily along this path. In the vicinity of Dead Horse Canyon, the SEA boundary turns generally westward and proceeds irregularly around development in Glendale, including natural parts of Toll, Hillcrest, Sherer, Idelwood, Pomeroy, Brand, and Childs canyons.

The lobe of the Verdugo Mountains between Hillcrest and Brand canyons is the area with the shortest possible traverse, which is a little less than two miles, to the Santa Monica Mountains in Griffith Park. Aerial fauna and plant seeds can easily make the journey, and the Los Angeles River channel at the base of the Santa Monica Mountains has developed a natural bottom and riparian habitat that must be inviting to migrants. A city park and a cemetery are on the route. The Verdugo Mountains are often viewed as one of the principle connections between the Santa Monica Mountains and the San Gabriel Mountains. However, it must be noted that this corridor is highly fragmented and probably impassable for most terrestrial wildlife.

Curving to the northwest in Burbank, the SEA boundary includes the natural, native vegetation of Elmwood, Story, Deer, and Sunset canyons. Sunset Canyon excludes a cherry-stem shaped area around a development. The Wildwood Canyon Park is included in the SEA with its many stately sycamore and coast live oak trees along the narrow stream course. In Stough Canyon the DeBell Municipal Golf course is excluded, as is the paved section of Stough Canyon Road and the buildings of the Stough Canyon Nature Center; however, the hillsides around Stough Canyon Road are included with a lobe of the SEA including the west ridge of Stough. On the west side of this ridge are the excluded Starlight Bowl and a nearby landfill. Continuing northwest, the SEA includes the natural,

undeveloped uppermost elevations of McClure, Brace, Cabrini canyons in Burbank, and Fisher, Jeffries, and Chandler canyons of Sun Valley.

North of Chandler is the developed south side of La Tuna Canyon Road, where the SEA boundary turns west along natural vegetation, using lot lines in part, which delineate the extent of fuel modification in this area of high fire hazard. The SEA boundary includes the north-facing south side of La Tuna Canyon for a distance of about three miles. The boundary crosses the road at the point where development stops and natural vegetation is on both sides of the road. From this point, the boundary continues the cherry-stem exclusion westward along the border of natural vegetation on the south-facing slope of La Tuna Canyon. Near the northwestern end of the Verdugo Mountains and La Tuna Canyon, the La Tuna stream joins the West Burbank Flood Control Channel. The SEA boundary includes the McDonald Creek drainage (tributary of La Tuna) and loops around the north ridge of McDonald Creek, changing direction to the northeast. The SEA boundary includes the natural area along the northern edge of the Verdugo Mountains by again following the edge of development in the Shadow Hills district of the City of Los Angeles. The boundary joins the northernmost point of the SEA near the undercrossing of Interstate-210 for Sunland/Foothill Boulevard.

The SEA is wholly within incorporated boundaries (cities of Los Angeles, Glendale, and Burbank), but much is preserved in conservation easements under the guidance of the Santa Monica Mountains Conservancy.

### **Vegetation**

Vegetation within the SEA is comprised of a large variety of community types. The diversity of the communities reflects the topography of the Verdugo Mountains. The southern slopes are affected by moist marine weather conditions, while the northern slopes are influenced by drier inland weather conditions. In addition, the steepness of many slopes causes sharp differences in vegetation on either side of a ridge. All plant species observed or recorded in previous documentation within the SEA are indicated in the *Comprehensive Floral & Faunal Compendium of the Los Angeles County SEA Update Study 2000 Background Report* and other analyses conducted for this area. Sensitive plant species and plant communities occurring or potentially occurring within the SEA are discussed in the Sensitive Biological Resources section.

Descriptions and general locations of the each plant community present within the SEA are given below.

**Chaparral:** A shrub community composed of robust species. Within this SEA, a number of chaparral subcommunities are found, and differentiated by their dominant plant species. These include chamise (*Adenostoma fasciculatum*), buck brush (*Ceanothus* spp.), scrub oak (*Quercus berberidifolia*), coast live oak (*Quercus agrifolia* var. *agrifolia*) and mosaics of these, depending on mixes of species and elevation. These and other shrub species form dense vegetation covers growing 5 to 10 feet in height. The development of chaparral is pronounced over large hillside areas throughout the SEA.

Corresponding MCV communities:

- *Adenostoma fasciculatum* (chamise chaparral) Shrubland Alliance
- *Adenostoma fasciculatum-Salvia apiana* (chamise-white sage chaparral) Shrubland Alliance
- *Arctostaphylos glauca* (bigberry manzanita chaparral) Shrubland Alliance
- *Ceanothus greggii* [vestitus] (cup leaf ceanothus chaparral) Shrubland Alliance
- *Ceanothus leucodermis* (chaparral whitethorn) Shrubland Alliance
- *Ceanothus oliganthus* (hairy leaf ceanothus chaparral) Shrubland Alliance
- *Prunus ilicifolia* (olly leaf cherry chaparral) Shrubland Alliance

Coastal Sage Chaparral Scrub: A shrubland community exhibiting less robust structure found in this SEA is coastal sage chaparral scrub. This plant community is dominated by California sagebrush (*Artemisia tridentata*), California brittle bush (*Encelia californica*), black sage (*Salvia mellifera*), and California buckwheat (*Eriogonum fasciculatum*). It also forms dense stands, which grow three to four feet in height. Within this SEA, it is generally found in scattered patches, which are highly integrated with mixed chaparral. These are primarily located in the lower elevation hillsides of the SEA.

Corresponding MCV communities:

- *Artemisia californica* (California sagebrush scrub) Shrubland Alliance
- *Artemisia californica-Eriogonum fasciculatum* (California sagebrush-California buckwheat scrub) Shrubland Alliance
- *Encelia californica* (California brittle bush scrub) Shrubland Alliance
- *Salvia leucophylla* (purple sage scrub) Shrubland Alliance
- *Salvia mellifera* (black sage scrub) Shrubland Alliance
- *Eriogonum fasciculatum* (California buckwheat scrub) Shrubland Alliance
- *Lotus scoparius* [*Acmispon glaber*] (deer weed scrub) Shrubland Alliance
- *Malacothamnus fasciculatus* (bush mallow scrub) Shrubland Alliance

Coast Live Oak Woodland: A plant community dominated by *Quercus agrifolia*. Within the SEA, this community includes coast live oak, which typically grows to heights of 20 to 40 feet, and forms either closed or open tree canopies. Oak woodland is most commonly found on north-facing slopes and in drainage bottoms and often intergrades with shrub communities. Understory vegetation varies from grassland in level areas to shrubs where topography is steeper.

Corresponding MCV community:

- *Quercus agrifolia* (coast live oak woodland) Woodland Alliance

Southern Willow Scrub: Found along widely scattered reaches of several drainages throughout this SEA. This community is dominated by species of willow (*Salix* spp.), which form nearly monotypic stands due to their dense growth, with an occasional cottonwood. These stands generally reach 10 to 20 feet in height with little understory vegetation.

Corresponding MCV communities:

- *Salix lasiolepis* (arroyo willow thickets) Shrubland Alliance

Riparian Forest: Along the major drainages, riparian forest is found. Typically, riparian forest grows along streams in bedrock-constrained, steep-sided canyons, resulting in a fairly narrow riparian corridor. The specific dominant plants are not known, but riparian trees such as California bay (*Umbellularia californica*), white alder (*Alnus rhombifolia*), coast live oak, western sycamore (*Platanus racemosa*) and willow occur. There are also a greater number of hydrophytic (moister favoring) plant species in the understory.

Corresponding MCV communities:

- *Alnus rhombifolia* (white alder groves) Forest Alliance
- *Umbellularia californica* (California bay forest) Forest Alliance
- *Quercus agrifolia* (coast live oak woodland) Woodland Alliance
- *Platanus racemosa* (California sycamore woodlands) Woodland Alliance

## **Wildlife**

Wildlife within the SEA is generally diverse and abundant due to large acreages of natural open space and diversity of habitat types. While a few wildlife species are entirely dependent on a single vegetative community, the entire mosaic of all the vegetation communities within the SEA and adjoining areas constitutes a functional ecosystem for a variety of wildlife species, both within the SEA and as part of the regional ecosystem.

The analysis of invertebrates is severely limited due to the lack of data; the SEA, however, undoubtedly supports healthy populations of a diverse assortment of invertebrate species. Amphibian populations are plentiful in the SEA due to the high moisture content provided by coastal conditions, as well as the large number of drainages and year-round water supplies. The SEA is also likely to support a variety of amphibians within the moister woodland areas and canyon bottoms. Many essential reptilian habitat characteristics are present within the SEA. These include rock outcroppings that allow for high visibility and small mammal burrows for cover and escape from predators and extreme weather. These characteristics, as well as the variety of habitat types present, are likely to support a wide variety of reptilian species.

The scrubland, woodland, riparian, and grassland habitats in the SEA provide foraging and cover habitat for year-round residents, seasonal residents, and migrating song birds. In addition, the SEA encompasses many year-round water sources that are located throughout the SEA and abundant raptor foraging, perching, and nesting habitat along the northern slopes of the Verdugo Mountains. The combination of these resources, as well as the confluence of many community types provides an unusually high diversity of bird species. Mammal populations within the SEA are diverse and reflective of the large size and variation of topography and community types.

All wildlife species previously recorded, as well as those expected to occur, within the SEA are indicated in the *Comprehensive Floral & Faunal Compendium of the Los Angeles County SEA Update Study 2000 Background Report*. Sensitive wildlife species occurring or potentially occurring within the SEA are discussed in the Sensitive Biological Resources section.

### **Wildlife Movement**

Although wildlife movement is hampered by adjacent rural development in proximity to the SEA, animals are still able to move through the Verdugo Mountains in many areas. Due to its large size and topographic complexity, many linkages occur within the SEA at various bottlenecks. These linkages allow movement between large open space areas within the SEA, as well as between areas outside the SEA toward the Angeles National Forest. The genetic flow through these areas is crucial in maintaining the diversity and viability of the species within the Verdugo Mountains. Some areas of probable, possible, and perhaps future connection have been indicated in the General Boundary and Resources section. Although there are significantly large open spaces within the SEA, contiguous habitat linkages between them is critical in reducing bottlenecks and providing for long-term sustainability. A wide variety of wildlife use linkages throughout the SEA, including mountain lion (*Puma concolor*), coyote (*Canis latrans*), mule deer (*Odocoileus hemionus*), bobcat (*Lynx rufus*), as well as a number of medium-sized mammals.

### **Sensitive Biological Resources**

Sensitive biological resources are habitats or individual species that have special recognition by federal, state, or local conservation agencies and organizations as endangered, threatened, and/or rare. This is due to the species' declining or limited population sizes, which usually results from habitat loss. Watch lists of such resources are maintained by the California Department of Fish and Game (CDFG), the United States Fish and Wildlife Service (USFWS), and special groups, such as the California Native Plant Society (CNPS). The following sections indicate the habitats as well as

plant and animal species present, or potentially present within the SEA, which have been accorded special recognition.

### *Sensitive Plant Communities and Habitats*

The SEA supports several habitat types considered sensitive by resource agencies. These are inventoried by California Department of Fish and Game (CDFG) in the California Natural Diversity Database (CNDDDB) [2011]. The CNDDDB includes state and federally-listed endangered, threatened, and rare vascular plants, as well as several sensitive vertebrate species. These communities include chamise-white sage chaparral, holly leaf cherry chaparral, California brittle bush scrub, California bay forest, and California sycamore woodlands, which occur throughout the SEA. These communities, or closely related designations, are considered high priority communities by the CDFG, which indicates that they are experiencing a decline throughout their range. The array and composition of these communities has been discussed in the Vegetation section.

### *Sensitive Plant Species*

The statuses of rare plants are hierarchically categorized by the CNPS using a rank and decimal system. The initial category level of Rare Plant Rank is indicated by the ranks 1A (presumed extinct in California), 1B (rare or endangered in California and elsewhere), 2 (rare or endangered in California but more common elsewhere), 3 (more information needed, a review list), and 4 (limited distribution). In cases where the CNPS has further identified the specific threat to the species, a decimal or Threat Code is added: .1 (seriously endangered in California), .2 (fairly endangered in California), or .3 (not very endangered in California).

The following special-status plant taxa have been reported or have the potential to occur within the SEA, based on known habitat requirements and geographic range information:

- Sonoran maiden fern (*Thelypteris puberula* var. *sonorensis*) RPR 2.2
- Nevin's barberry (*Berberis nevinii*) FE, SE, RPR 1B.1
- Lewis' evening-primrose (*Camissonia lewisii*) RPR 3
- San Fernando Valley spineflower (*Chorizanthe parryi* var. *fernandina*) FC, SE, RPR 1B.1
- Parry's spineflower (*Chorizanthe parryi* var. *parryi*) RPR 1B.1
- Peruvian dodder (*Cuscuta obtusiflora* var. *glandulosa*) RPR 2.2
- Palmer's grapplinghook (*Harpagonella palmeri*) RPR 4.2
- Davidson's bushmallow (*Malacothamnus davidsonii*) RPR 1B.2
- White rabbit-tobacco (*Pseudognaphalium leucocephalum*) RPR 2.2
- Greata's aster (*Symphotrichum greatae*) RPR 1B.3
- Slender mariposa lily (*Calochortus clavatus* var. *gracilis*) RPR 1B.2
- Plummer's mariposa lily (*Calochortus plummerae*) RPR 1B.2
- California sawgrass (*Cladium californicum*) RPR 2.2
- California satintail (*Imperata brevifolia*) RPR 2.1
- Ocellated Humboldt lily (*Lilium humboldtii* ssp. *ocellatum*) RPR 4.2

### *Sensitive Animal Species*

The following special-status animal species are reported or have the potential to occur within the SEA, based on known habitat requirements and known range attributes:

- Silvery legless lizard (*Anniella pulchra pulchra*) FSS, SSC
- Coastal whiptail (*Aspidoscelis tigris stejnegeri*) CDFG Special Animals List
- Rosy boa (*Charina trivirgata*) BLMS, FSS

- Coast horned lizard (*Phrynosoma blainvillii*) BLMS, FSS, SSC
- Southern California rufous-crowned sparrow (*Aimophila ruficeps canescens*) CDFG Watch List
- Bell's sage sparrow (*Amphispiza belli belli*) BCC, CDFG Watch List
- Southwestern willow flycatcher (*Empidonax traillii extimus*) FE, FSS, SE, USBC, AWL, ABC
- Least Bell's vireo (*Vireo bellii pusillus*) FE, BCC, SE, USBC, AWL, ABC
- Pallid bat (*Antrozous pallidus*) FSS, BLMS, SSC, WBWG High
- Western mastiff bat (*Eumops perotis californicus*) BLMS, SSC, WBWG High
- Silver-haired bat (*Lasionycteris noctivagans*) WBWG Medium
- Hoary bat (*Lasiurus cinereus*) WBWG Medium
- Western yellow bat (*Lasiurus xanthinus*) WBWG High
- Big free-tailed bat (*Nyctinomops macrotis*) SSC, WBWG Medium-High
- American badger (*Taxidea taxus*) SSC

**Ecological Transition Areas (ETAs)**

There are no ETAs designated within this SEA.

**Regional Biological Value**

The SEA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

**CRITERIA ANALYSIS OF THE VERDUGO MOUNTAINS SEA**

	<b>Criterion</b>	<b>Status</b>	<b>Justification</b>
A)	The habitat of core populations of endangered or threatened plant or animal species.	Not Met	No critical habitats are designated within this SEA.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The Verdugo Mountains combined is an extensive, relatively undisturbed island of natural vegetation in an urbanized area, which is very rare in Southern California.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	The Verdugo Mountains combined is an extensive, relatively undisturbed island of natural chaparral and canyon vegetation. It is an important island refuge for migration among the mountain ranges of the northern portion of the County. Aerial animals and plant seeds can easily traverse the distances between the Verdugo Mountains and Santa Monica, Santa Susana, and the San Gabriel mountains. The Verdugo Mountains serve as the centerpiece of these connections. The hillsides are covered by chaparral and coastal sage chaparral scrub. The canyons' riparian

			vegetation includes California bay, sycamores, ferns and tiger lilies. These plant communities provide habitat that is essential to the diverse and abundant fauna found in the area.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The Verdugo Mountains serve as an arm of the San Gabriel Mountains, extending towards the eastern end of the Santa Monica Mountains in Griffith Park—only two miles distant. Aerial animals and plant seeds easily cross the gap. The Verdugo Mountains are exceedingly important for connections among the Santa Monica, Santa Susana, and the San Gabriel mountains. Genetic interchange, by way of this linkage, is important in perpetuating the genetic variability in isolated populations, which maintains healthy ecosystems and resilience to climate change.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The geographic location of the Verdugo Mountains makes them important for scientific study, genetic interchange between otherwise isolated populations, and recreation for urban residents. The area has already been used for studies concerned with public health.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The Verdugo Mountains have been impacted by dirt roads, one paved road in Tuna Canyon, fuel breaks, transmission lines and isolated buildings for houses, radio towers, and water tanks. Because of their extent, however, the Verdugo Mountains are still considered largely natural and little impacted—a prime example of the chaparral and coastal sage chaparral scrub once prevalent in the County coastal areas.

In conclusion, the area is an SEA because it contains: B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

## 27. Verdugo Mountains SEA Sources

Atlantis Scientific. 1990. *Biological Impact Report, Project No. 90088/PM - 21998/CUP*, Beverly Hills, California.

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

Biological Assessment Services. 1992. *Biological Resources Assessment Vesting Tentative Tract No. 47927, SEA No. 12, Palo Comado Canyon*, Los Angeles County, California.

California Department of Fish and Game, Natural Diversity Database. January 2011. *Special Animals* (898 taxa). 60 pp.

California Department of Fish and Game, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Game, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

C.W. Cook Co. Inc. 1990. *SEATAC Information Packet for SEA No. 4, Project No. 86089* (Biology Reports), Los Angeles, California.

Christopher A. Joseph and Associates. 2004. *FINAL ENVIRONMENTAL IMPACT REPORT FOR THE CANYON HILLS PROJECT*, Prepared for the City of Los Angeles

County of Los Angeles, Department of Regional Planning. 2006. *The Verdugo Mountains SEA Description*, Los Angeles County, California.

Dale, N. 1986. *Flowering Plants: The Verdugo Mountains, Coastal & Chaparral Regions of Southern California*.

David Carroll & Associates. 1992. *Biological Constraints Analysis Report, Project No. 91-359, SEA No. 3 Buffer, Latigo Canyon Road*, Los Angeles County, California.

David Carroll & Associates. 1991. *Biota Report for Subdivision Parcel Map No. 89494 Latigo Canyon*, Los Angeles County, California.

David J. Tanner & Associates. 1988. *Biological Resources Assessment and Tree Inventory and Evaluation for Malibu Terrace, Tract No. 45342 and No. 45343, SEA No. 12*, Los Angeles County, California.

Edelman, Paul. 1990 (revised 2/91). *Critical Wildlife Corridor/Habitat Linkage Areas Between the Santa Susana Mountains, The Simi Hills and The Verdugo Mountains*, Los Angeles County, California.

Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.