

DRAFT

**LOS ANGELES COUNTY GENERAL PLAN
APPENDIX E:
CONSERVATION AND NATURAL RESOURCES
ELEMENT**

Appendix E: Conservation and Natural Resources Element Resources

I. Open Space and Natural Areas in Los Angeles County

Los Angeles County offers a wide variety of open space and natural areas. The following open space and natural areas are managed by the County or are located primarily within the unincorporated areas:

Angeles National Forest

The Angeles National Forest was established by Executive Order in 1892 and is managed by the U.S. Forest Service. The Forest covers over 650,000 acres. The Angeles National Forest manages the watersheds within its boundaries to provide water to Southern California and to protect surrounding communities from catastrophic floods. The land within the Angeles National Forest is diverse in appearance and terrain, and provides many opportunities for recreational and scenic enjoyment. Much of the Angeles National Forest is ~~covered with~~ characterized by dense chaparral, pine and fir covered slopes, ~~as with~~ elevations in the Angeles National Forest range ranging from 1,200 to 10,064 feet.

Deane Dana Friendship Park

Deane Dana Friendship Park, which is a 123-acre park located on the Palos Verdes Peninsula, affords dramatic panoramic views of Santa Catalina Island, Los Angeles and Long Beach harbors, the City of Los Angeles to the north, and the San Gabriel and San Bernardino mountain ranges. There are hiking trails throughout the park.

Devil's Punchbowl Natural Area

Devil's Punchbowl is a 1,310-acre natural area that consists of rugged wilderness rock formations along the San Andreas Fault on the northern slope of the San Gabriel Mountains. The terrain climbs from 4,200 feet to 6,500 feet in elevation, with natural plant and animal communities ranging from desert scrub to pine forests. A seasonal stream runs through the natural area.

Eaton Canyon Natural Area

Situated at the base of Mt. Wilson, this 190-acre natural area contains several plant and native habitat communities. Eaton Creek flows through the Canyon during all but the summer months. The 7,600-square-foot nature center contains displays of local flora and fauna, ecosystem concepts and live animals. The natural area offers five miles of nature trails and an equestrian trail, and serves as a trailhead to the Mt. Wilson Toll Road and Henninger Flats.

High Desert Wildlife and Wild Flower Sanctuaries

The County currently operates ~~eight~~ twelve wildlife sanctuaries and one wildflower sanctuary in the high desert of Antelope Valley. Ranging from 2,500 to over 3,600 feet in elevation and encompassing more than 2,000 acres, the sanctuaries offer opportunities for spring wildflower viewing, bird watching, hiking and horseback riding. Wildlife seen on the preserves vary from horned lizards, chuckwallas and rattlesnakes, to prairie falcons and golden eagles. Insect life is most abundant during the warmer months, and in spring, the Joshua tree and other large shrubs provide nesting sites for a variety of songbirds. Other protected animals are the kit fox, desert tortoise and Mojave ground squirrel.

Kenneth Hahn State Recreation Area

The 338-acre Kenneth Hahn State Recreation Area, managed by the County, includes large areas of native coastal sage scrub habitat, lawns and landscaped areas, picnic sites, ~~tot lots~~ playgrounds, a fishing lake, a lotus pond, a community center, and five miles of trails. One of the most actively used features is the park's more than seven miles of footpaths and trails.

Michael D. Antonovich Open Space Preserve

The Michael D. Antonovich Open Space Preserve offers 500 acres of dedicated open space in the Santa Susana Mountains and is managed by the Mountains Recreation and Conservation Authority (MRCA). Located on the northern border of Los Angeles, this open space preserve contains ~~a diversity~~ of diverse flora and fauna, from big cone Douglas fir, California walnut and oak trees to black bears, deer and mountain lions. The Preserve also provides important habitat connections through its numerous wilderness trails in the Rim of the Valley corridor of the Santa Clarita Woodlands Park.

Placerita Canyon Natural Area

This 350-acre natural area is located in an east-west running canyon featuring oak groves, chaparral-covered slopes and a sycamore-lined stream. Placerita Canyon is home to the Oak of the Golden Dream, where gold was "first" discovered in California in 1842, and Walker's Cabin, which is a relic of early frontier living. Placerita Canyon contains eight miles of hiking trails.

Santa Monica Mountains National Recreation Area

The 150,000-acre Santa Monica Mountains National Recreation Area, ~~is~~ is a part of the National Park System, ~~which~~ encompasses the mountain range from the Oxnard Plain in Ventura County east through, ~~past~~ Topanga State Park to Franklin Canyon and the Hollywood Bowl in Los Angeles. The Recreation Area preserves natural habitats, includes historical and cultural sites, offers recreational opportunities, and acts to improve the air quality for the Los Angeles Basin. Covered by chaparral, oak woodlands, and coastal sage scrub, it is home to many species listed as rare, threatened, or endangered.

Santa Catalina Island Regional Park

Santa Catalina Island Regional Park consists of 41,000 acres of natural open space, with miles of hiking trails, camp sites and a wildlife sanctuary. The Santa Catalina Island Interpretive Center is located within the park.

Schabarum Regional Park

The 640-acres Schabarum Regional Park ~~is comprised~~ consists of open space and natural areas, including picturesque canyons and rolling hills for hiking, biking and horseback riding. ~~Over 90 percent of this park has been left in its natural state for the public to enjoy.~~ The park features equestrian areas, a dog park, tennis and volleyball courts, multiple picnic areas, and playgrounds, in a pastoral setting.

Vasquez Rocks Natural Area Park

This 945-acre natural area is a hiking, picnicking, and equestrian area. The park is located in the high desert near Agua Dulce Springs and features rock formations, ~~Tatavium Indian~~ Tatavium sacred sites,

and a seasonal stream. The principal plant communities are desert, chaparral, and riparian.

Whittier Narrows Natural Area

This natural area occupies approximately 300 acres in the southern portion of the Whittier Narrows Recreation Area. Bordering the San Gabriel River, the Natural Area is home to several habitats, with the dominant one being a riparian woodland. The southeastern portion of the site features four seasonal lakes that provide a winter sanctuary for migrating waterfowl and are opened by special permit for birding and photography. ~~This area is near lakes and contains many plants and animals that are typically found within a wetland community.~~ The area provides habitats for a wide variety of wildlife, including least Bell's vireos and California gnatcatchers.

II. Conservancies

The County works with various conservancies to maintain and protect open space land in Los Angeles County. Land conservancies are private, nonprofit organizations and public agencies that share a common goal: to conserve land for the benefit of people and nature. Land conservancies are generally started by community residents who wish to preserve a certain area or piece of open space land on a local or regional scale. Land conservancies have the flexibility to acquire, hold and manage land in the public interest, and also to preserve open space through voluntary conservation agreements with landowners, ~~which permanently protecting~~ the land from development, while the title remains with the landowner. ~~Most conservancies work~~ Often working in partnership with local governments, ~~land conservancies often~~ and provide various levels of educational programs and land restoration and/or land enhancement projects. undertake projects to restore and enhance the natural habitats they protect.

~~The County works with a number of conservancies to preserve and protect the County's open spaces:~~

Antelope Valley Conservancy

~~The Antelope Valley Conservancy is a local land trust conservancy that obtains and stewards lands that are important to the community for quality of life, scenic beauty, and plant and animal habitat. AVC focuses on Joshua tree woodlands, the keystone species of the Mojave Desert, which supports a wide variety of native species. Most of the Conservancy's targeted preservation lands are in the County's designated Significant Ecological Areas. (<http://www.avconservancy.org/>)~~

Baldwin Hills Conservancy

~~The Baldwin Hills Conservancy (BHC) was created by the State in 2000. Specifically, BHC is responsible for: implementing the Baldwin Hills Park Master Plan; prioritizing and implementing acquisition of additional recreational and open space land for the expansion of Kenneth Hahn State Recreation Area; conducting planning activities for the area; and developing and coordinating a program of resource stewardship for optimum recreational and natural resource value based on the needs of the surrounding community. (<http://www.bhc.ca.gov/>)~~

California Coastal Conservancy

~~Established in 1976, the California Coastal Conservancy is a state agency that purchases, protects, restores, and enhances coastal resources, and provides access to the shore. The Conservancy works in partnership with local governments, other public agencies, nonprofit organizations, and private landowners. To date, the Conservancy has undertaken more than 1,800 projects along the 1,100 mile~~

California coastline. (<http://www.scc.ca.gov/>)

Catalina Island Conservancy

In 1974, the Santa Catalina Island Company entered into a 50-year Open Space Easement agreement with the County, guaranteeing public recreational and educational use of 41,000 acres of Santa Catalina Island, consistent with good land conservation practices. The Santa Catalina Island Company subsequently deeded this land to the non-profit Catalina Island Conservancy (formed in 1972), along with an additional 1,135 acres. The Catalina Island Conservancy continues to manage 42,135 acres on Santa Catalina Island, providing the highest level of conservation protection to 88 percent of Santa Catalina Island. The Conservancy's mission is "to be a responsible steward of its lands through a balance of conservation, education and recreation." The conservation activities include wildlife management, a plant ecology program, invasive plant management, restoration of degraded areas and scientific research by many different researchers from throughout the country. Other activities include various educational camps for youth, several campgrounds and hiking and biking trails. (<http://www.catalinaconservancy.org/>)

Mountains Recreation and Conservation Authority

Established in 1985, the Mountains Recreation and Conservation Authority (MRCA) is a partnership between the Santa Monica Mountains Conservancy (SMMC), the Conejo Recreation and Park District, and the Rancho Simi Recreation and Park District. The MRCA is dedicated to the preservation and management of open space and parkland, watershed lands, trails, and wildlife habitat. The MRCA manages almost 60,000 acres of public lands and parks, and provides comprehensive education and interpretation programs for the public. The MRCA works in cooperation with the SMMC and other local partners to acquire parkland, participate in vital planning processes, and complete major park improvement projects. The MRCA also provides natural resources and scientific expertise, critical regional planning services, park construction services, park operations, fire prevention, ranger services, and educational and leadership programs for youth. (<http://www.mrca.ca.gov/>)

Newhall Ranch High Country Recreation and Conservation Authority

The Newhall Ranch High Country Recreation and Conservation Authority is a joint powers authority formed by the SMMC, the County, and the City of Santa Clarita. Its purpose is the conservation and management of public open space lands set aside for habitat and recreation pursuant to the Los Angeles County Board of Supervisors approval of the Newhall Ranch project, the Westridge project, and any other such open space lands dedicated by Newhall Land Company or its successors. (<http://smmc.ca.gov/NRHCRCA.asp>)

Puente Hills Habitat Preservation Authority

The Puente Hills Habitat Preservation Authority (PHHPA) is a joint powers authority with a board of directors consisting of the City of Whittier, the County, Los Angeles County Sanitation Districts, and the Hacienda Heights Improvement Association. The PHHPA was created in 1994 as mitigation for the Puente Hills Landfill. To date, the PHHPA manages 3,860 acres of preserved public open space. The PHHPA's main focus is to acquire the remaining open space within its jurisdiction, with special consideration given to the Hacienda Heights area. (<http://www.habitatauthority.org/>)

San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy

The San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy (RMC) was created by the State in 1999. RMC is one of nine conservancies within the California Resources Agency. Its mission is to preserve open space and habitat in order to provide for low-impact recreation and educational uses, wildlife habitat restoration and protection, and watershed improvements. RMC's territory covers eastern portions of Los Angeles County and western Orange County, which is a vast and varied area that includes mountains, valleys, rivers, coastal plain, and coastline. (<http://www.rmc.ca.gov/>)

San Gabriel Mountains Regional Conservancy

The San Gabriel Mountains Regional Conservancy is devoted to watershed management and a great variety of other projects in the San Gabriel River watershed of the eastern portion of Los Angeles County. Included in the region are the San Gabriel Mountains, the San Gabriel River valley and related areas. (<http://www.sgmrc.org/>)

Santa Clarita Watershed Recreation and Conservation Authority

The Santa Clarita Watershed Recreation and Conservation Authority (SCWRCA) was established in 1993 by the City of Santa Clarita and the Santa Monica Mountains Conservancy in order to cooperatively plan for the preservation of open space, trails, parkland, and watershed protection in the Santa Clarita area and the upper Santa Clara River watershed. In 2002, the SCWRCA finalized the City's and Conservancy's long-standing efforts to acquire the 442-acre Whitney Canyon Ranch. (<http://smmc.ca.gov/SCWRCA.asp>)

Santa Monica Mountains Conservancy

The Santa Monica Mountains Conservancy (SMMC) was established by the State in 1980. Since that time, it has helped to preserve over 60,000 acres of parkland in both wilderness and urban settings, and has improved more than 114 public recreational facilities throughout Southern California. Through direct action, alliances, partnerships, and joint powers authorities, the SMMC's mission is to strategically buy back, preserve, protect, restore, and enhance treasured pieces of Southern California to form an interlinking system of urban, rural and river parks, open space, trails, and wildlife habitats that are easily accessible to the general public. (<http://smmc.ca.gov/>)

Watershed Conservation Authority

The Watershed Conservation Authority (WCA) was created in 2003 as a joint powers entity of the RMC and the Los Angeles County Flood Control District. The focus of the WCA is on projects that will provide open space, habitat restoration, and watershed improvement projects in the watersheds of both the San Gabriel River and the lower Los Angeles River. (<http://watershedconservationauthority.org/>)

Wildlife Corridor Conservation Authority

The Wildlife Corridor Conservation Authority (WCCA) was established to provide for the proper planning, conservation, environmental protection, and maintenance of lands within the Puente Chico Hills corridor area. Its goal is to assure that sufficient continuity of habitat can be preserved to maintain a functioning wildlife corridor made up of about 40,000 acres of land located between the Santa Ana

DRAFT

~~Mountains and Whittier Hills. WCCA's governing board consists of representatives from the cities of Brea, Whittier, Diamond Bar, La Habra Heights, the Santa Monica Mountains Conservancy, California Department of Parks and Recreation, California Department of Fish and Wildlife, the County, and two public members. (<http://smmc.ca.gov/WCCA.html>)~~

Antelope Valley Conservancy

Founded in 2005, the Antelope Valley Conservancy (AVC) is a local, volunteer-managed land trust conservancy that undertakes habitat preservation and stewardship projects in partnership with a variety of State, County, and local agencies. Sensitive resources protected by AVC initiatives include wetlands, Joshua tree woodlands, populations of alkali mariposa lily, desert tortoises, and burrowing owls. The AVC offers presentations at schools and community groups to educate the public about desert habitat, wildlands stewardship, climate change and carbon footprint, and sustainability. Most of the Conservancy's targeted preservation lands are in the County's designated Significant Ecological Areas. (<http://www.avconservancy.org/>)

Arroyos and Foothills Conservancy

Founded by a small group of volunteers in 2000, the nonprofit Arroyos and Foothills Conservancy (AFC) identifies and conserves undeveloped private properties situated along the San Gabriel Mountains, from Altadena to Tujunga, and in the San Rafael Hills and Verdugo Mountains. Utilizing a network of expert advisors and collaborative partnerships with conservation organizations and state agencies, the AFC gathers extensive information about the water, habitat, and wildlife at each property, along with its potential for community programs. They then acquire high-priority parcels from willing owners by purchase or donation, or place use restrictions on properties in order to preserve them as open space in perpetuity. One of AFC's focus areas is on working to improve wildlife movement opportunities among the Verdugo Mountains, the San Rafael Hills, and Griffith Park, ultimately linking these areas back to the San Gabriel Mountains. The AFC also works to promote backyard biodiversity through educational communications, plant giveaways, helping create native plant gardens in schools and other private green spaces, and backyard camera trap projects. The organization also provides educational programs to schools across Los Angeles County. (<https://arroyosfoothills.org>)

Baldwin Hills and Urban Watersheds Conservancy

The California Resources Agency established the Baldwin Hills Conservancy in 2000, and expanded its geographic scope in 2023, renaming it the Baldwin Hills and Urban Watersheds Conservancy (BHUWC). The BHUWC acquires open space and manages public lands within the Baldwin Hills, Ballona Creek and Upper Dominguez Channel watersheds, an area of approximately 70 square miles, to provide recreation, restoration and protection of wildlife habitat for the public's enjoyment and educational experience. The County and the BHUWC are members of the Baldwin Hills Regional Conservation Authority (BHRCA), which has a mission to acquire, expand and improve open space within the Baldwin Hills and Ballona Creek, and other natural and recreational areas in the Second Supervisorial District of the County. The BHUWC and BHRCA focus on acquiring and preserving open space lands in the most densely populated part of the County. (<http://www.bhc.ca.gov>)

California Coastal Conservancy

Established in 1976 under the California Resources Agency, the California Coastal Conservancy (CCC) has a mission to protect and improve natural lands and waterways, help people access and

DRAFT

enjoy the outdoors, enhance climate resilience, and sustain local economies along the 1,100-mile length of California's coast, and within the watersheds of rivers and streams that extend inland from the coast. Over the years, the CCC has partnered with local governments, other public agencies, nonprofit organizations, and private landowners on more than 2,300 projects statewide. (<http://www.scc.ca.gov/>)

Catalina Island Conservancy

Formed in 1972, the Catalina Island Conservancy (CIC) manages 88 percent of Santa Catalina Island (42,135 acres). Its mission is "to be an exemplary steward of Island resources through a balance of conservation, education and recreation." The CIC's conservation programs focus on resiliency of the Island and its plant and animal populations, partnerships with researchers and implementation of long-term studies, preservation of historic cultural resources, and synergy of conservation efforts with education and recreation programming. The Conservancy's education programming offers the public opportunities to explore Catalina's wildlands through excursions and training, as well as learning pathways accessible from home. (<https://catalinaconservancy.org>)

Los Angeles Neighborhood Land Trust

The nonprofit Los Angeles Neighborhood Land Trust (LANLT) was founded in 2002 to address park and health inequities in Los Angeles County. The organization's mission is to contribute to the equity and well-being of neighborhoods through the development of parks, gardens, and community-driven organizing across greater Los Angeles. Focusing its efforts in communities that have little access to green space, the LANLT has established 30 parks and gardens providing 22 acres of open space in underserved neighborhoods. (<https://www.lanlt.org>)

Puente Hills Habitat Preservation Authority

Established in 1994 as a condition of approval for the Puente Hills Landfill, the Puente Hills Habitat Preservation Authority (PHHPA) is a joint powers authority with a board of directors consisting of the City of Whittier, the County, Los Angeles County Sanitation Districts, and the Hacienda Heights Improvement Association. The PHHPA currently manages 3,880 acres of preserved public open space, conducts public outreach, and provides trail access, but its main focus is to acquire the remaining open space within its jurisdiction, with special consideration given to the Hacienda Heights area. (<https://www.habitatauthority.org>)

San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy

Established in 1999 under the California Resources Agency, the San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy (RMC) has a mission to preserve open space and habitat in order to provide for low-impact recreation and educational uses, wildlife habitat restoration and protection, and watershed improvements. The RMC's territory includes a wide swath of Los Angeles County, from near Palmdale southwest to Santa Clarita, east across the San Gabriel Mountains, and then south across the lowlands and Puente Hills to the Palos Verdes Peninsula and Long Beach. The RMC issues grants to eligible applicants including non-profits organizations, public agencies, and tribes for projects that preserve and increase public access to open space and habitat, and help create healthy and climate resilient communities. (<https://rmc.ca.gov>)

San Gabriel Mountains Regional Conservancy

Founded in 1997, the San Gabriel Mountains Regional Conservancy (SGMRC; <https://sgmrc.org>) is a nonprofit public benefit corporation dedicated to the preservation of land and structures in the San Gabriel Mountains region for historic, educational, ecological, recreational, and open-space purposes. The SGMRC emphasizes land conservation, management practices, public education, and watershed stewardship to foster sustainable regional ecosystems. As a partner to state entities like the San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy, the SGMRC supports collaborative efforts in environmental protection and education, including operation of the nature center at the County's Santa Fe Dam Recreational Area.

Santa Monica Mountains Conservancy

Established in 1980 under the California Resources Agency, the Santa Monica Mountains Conservancy (SMMC) operates through direct action, alliances, partnerships, and joint powers authorities to strategically buy back, preserve, protect, restore, and enhance natural open space in southern California, establishing an interlinking system of urban, rural and river parks, open space, trails, and wildlife habitats that are easily accessible to the general public. To date, the SMMC has helped to preserve more than 75,000 acres of parkland in both wilderness and urban settings, has improved more than 114 public recreational facilities in the region, and has given grants to nonprofit organizations for educational and interpretation programs that have served hundreds of thousands of children and other park visitors. (<https://smmc.ca.gov>)

The SMMC is a member of six active joint powers authorities in Los Angeles County, which function as legally separate government entities with their own governing boards. Central to their success is the ability to complement the work of other organizations protecting land in the southern California mountains to acquire and improve open space and parklands, affording educational opportunities, and providing stewardship for a wide variety of public park and open space resources. The joint powers authorities are:

- Desert and Mountain Conservation Authority
- Mountains Recreation and Conservation Authority
- Newhall Ranch High Country Recreation and Conservation Authority
- Santa Clarita Watershed Recreation and Conservation Authority
- Whittier-Puente Hills Conservation Authority
- Wildlife Corridor Conservation Authority

Transition Habitat Conservancy

Founded in 2005, the nonprofit Transition Habitat Conservancy (THC) focuses on land acquisitions and habitat stewardship in the West Mojave Desert. Through their mission to “protect transition zones and wildlife corridor ecosystems and their scenic, agricultural, and cultural resource values in the West Mojave Desert,” THC has conserved over 11,000 acres of habitat in the Antelope Valley including ownership of over 4,000 acres of critical desert tortoise habitat. THC also works in partnership with the Bureau of Land Management to manage nearly 250,000 acres of public land in the Freemont-Kramer and Super Cronese Areas of Critical Environmental Concern.

(<https://www.transitionhabitat.org>)

TreePeople Land Trust

Founded in 1981, the nonprofit TreePeople Land Trust (TPLT) works to permanently conserve natural land, mainly in the Santa Monica Mountains, and the organization currently owns and/or manages more than 3,000 acres of undeveloped open space. TPLT operates through property acquisition, conservation easements, and management agreements with its partner agencies. The TPLT manages an extensive network of public hiking trails in the central Santa Monica Mountains, is engaged in ongoing native plant restoration projects, and provides resources for scientific research throughout its preserves and properties. The organization has partnered with the County to implement a residential stormwater retrofit program, to encourage replacement of lawns, and to implement school greening projects. (<https://treepeoplelandtrust.org>)

Watershed Conservation Authority

The Watershed Conservation Authority (WCA) was created in 2003 as a joint powers entity of the San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy and the Los Angeles County Flood Control District. The WCA's mission is to expand and improve open space and recreational opportunities to conserve, restore, and enhance natural habitats within the watersheds of the San Gabriel River and Lower Los Angeles River consistent with the goals of flood protection, water supply, groundwater recharge and water conservation. Embracing a "work with nature" approach, the WCA partners with local and regional entities to plan and implement projects to improve watersheds through investment in open space, parks, trails, bikeways, and greenways. The WCA maintains more than 700 acres of open space, ranging from foothills lands to urban river parks, that serve some of the County's most diverse and densely populated neighborhoods. (<https://www.wca.ca.gov>)

III. Regional Habitat Linkages

Habitat linkages are defined as area within the overall range of a species or suite of species that possess sufficient cover, food, forage, water and other essential elements to serve as a movement pathway, or between two or more larger areas of habitat. Depending on the species, linkages vary in size. For example, a belt of coastal sage scrub traversing a golf course, connecting sage scrub habitat areas on either side, providing a safe passage zone for smaller, slower-moving species (such as lizards and rodents) to maintain population¹ connectivity between the two sides of the golf course is

¹ In population biological terms, a "population" is the group of individuals of any species that may interact or interbreed with each other. Thus, many groups of organisms within Los Angeles County may belong to populations that extend far beyond county borders. Nevertheless, populations are typically structured (often because of intervening unsuitable habitat areas) so that individuals are much more likely to interact with other nearby individuals in what may be termed sub-populations or "demes." However, rather than detracting from the scope of the SEA descriptions by detailing a long discussion of the technicalities of population biology, and to benefit readability for the lay reader, the County has chosen to use the term "population" in a broad colloquial sense throughout this document. Thus, in this document the term "population" may often refer to demes, but the intention is to refer to groups of

one form of habitat linkage.

Wildlife corridors, which are areas of open space of sufficient width to permit larger, mobile species (such as foxes, bobcats and coyote) to pass between larger areas of open space, or to disperse from one major open space region to another, are another type of habitat linkage. Such areas are generally several hundred feet wide, unobstructed, and usually possess cover, food and water. The upland margins of a creek channel, open ridgelines, open valleys or the bottoms of drainages often serve as major corridors locally, as do riparian alignments.

Biological resources are important in a regional context, serving to connect resources in adjacent local jurisdictions. Critical biological resources are maintained through habitat connectivity, which sustains population genetic diversity, and provides refuge for migrant species. Regional habitat linkages are identified in the Conservation and Natural Resources Element. The Antelope Valley, Puente Hills, San Andreas, Santa Clara River, Santa Felicia, Santa Monica Mountains, and Santa Susana Mountains and Simi Hills SEAs serve as important regional habitat linkages. More details about linkages between and within each of these SEAs are provided below:

Antelope Valley SEA

The SEA extends from the Angeles National Forest to the playa lakes within Edwards Air Force Base, encompassing most of the ~~two~~three largest drainages systems exiting the northern slope of the San Gabriel Mountain range. The geographical features of the SEA serve as a major habitat linkage and movement corridor for all wildlife species within its vicinity and, in an intergenerational sense, many of the plant species. Ecologically generalist species (such as mountain lion, bobcat, coyote, gray fox,) have the ability to move across such vast areas and through changing habitat types. For such species, the SEA may serve as an important system for long-term and genetic exchange among populations. For smaller or less-mobile species or taxa, which are narrowly restricted in their habitat needs, the SEA can serve as a broad linkage zone, in which individual movement can take place during seasonal population dispersal or over generations. This provides essential genetic exchange within and between metapopulations. The two drainages, combined with the upland terrestrial Desert-Montane transect portion of the SEA, ensure linkage and direct movement areas for all of the wildlife species present within the County portion of the Antelope Valley.

Puente Hills SEA

Evidence of significant wildlife movement throughout the Puente Hills SEA was documented in a two-year carnivore study commissioned by the Santa Monica Mountains Conservancy as part of a multi-jurisdictional effort to establish a regionwide wildlife movement linkage. Additional studies on wildlife movement through the area include several focusing on the Harbor Boulevard and Colima Road Underpasses. These studies can be accessed at the following link: <http://www.habitatauthority.org/studies>. This SEA represents the County portion of a continuous series of natural open space within the Puente Hills and Chino Hills. Overall, this open space extends north

individuals that interact in an ecologically meaningful ways on a regular basis.

and west from State Route-91 in Orange and Riverside counties to the Whittier Narrows reach of the San Gabriel River. The Puente and Chino Hills are a natural, physical link between the Santa Ana Mountains and the San Gabriel River. The San Gabriel River flows from and links to the San Gabriel Mountains. By virtue of these linkages and a complex of interconnected habitat units, the Puente and Chino Hills function as both an important wildlife linkage and resident habitat area for regional wildlife populations.

San Andreas SEA

The SEA includes several important linkages for wildlife movement. The foothills in the ~~western-most~~ westernmost part of the SEA are an important linkage between the San Gabriel Mountains, the Tehachapi Mountains, and the Coastal Ranges. The linkage to the Tehachapi Mountains is important because the Tehachapis connect to the southern-most extent of the Sierra Nevada Mountains. The Tehachapi Mountains represent the only mountain linkage from the Transverse Ranges and the Coast Ranges to the Sierra Nevada Range. This feature may be an important topographic reference for migrating birds, and provides high elevation foraging grounds along the migratory route. The several ranges that meet at the western end of the SEA provide a valuable link for gene flow between divergent subspecies, varieties, and populations of many species. The SEA includes numerous drainages that extend onto the Antelope Valley floor towards resources such as the Fairmont and Antelope buttes. These washes provide an important linkage for animals traveling between the Valley floor, the buttes and the western part of the San Gabriel Mountains. In addition, Anaverde Creek, Amargosa Creek, and Pine Canyon facilitate east-west wildlife movement through the mountains, Portal Ridge, and Ritter Ridge. Tributary drainages from the Santa Clara River, such as Elizabeth Lake Canyon and San Francisquito Canyon, connect coastal drainages and the coastal ecoregion to the San Andreas Fault and interior watersheds. The frequency of valuable riparian communities along this travel route, which is located within an otherwise arid climate, further contributes to the SEA's importance for wildlife and habitat linkages in the region.

Santa Clara River SEA

Historically (and prehistorically) the riparian corridor along the Santa Clara River has served as the primary east-west linkage between the Pacific coastline, Coast Ranges, interior ranges, high desert and southern Sierra (via the Tehachapi Range). Animals moving through the Santa Clara River at one time had unobstructed passage along the river and within its tributaries. The present configuration of the tributary drainages has reduced connectivity from the Santa Clarita Valley to the north, but the Santa Clara River remains relatively intact and open. The SEA embraces the river corridor and the linkage zones that are considered essential to ensuring connectivity and resource values within the historic movement zones for all of the wildlife species present within the County portion of the Santa Clara River.

Santa Felicia SEA

Historically, riparian corridors have served as linkages between the Pacific coastline, Coast Ranges, interior ranges, the high desert and southern Sierras (via the Tehachapi Range). The Santa Felicia stream corridor likely serves the functions today. The elevation in this area is lower than that of the Los Padres National Forest, to the north, which facilitates animal movement within the riparian systems between Piru Lake in Ventura County and the San Gabriel Mountain Range in the County. The tributary drainages for Santa Felicia Creek within this SEA remain intact and unobstructed.

Santa Monica Mountains SEA

Although wildlife movement is hampered by rural development in the SEA, animals are still able to move through the Santa Monica Mountains in many areas. Due to its large size and topographic complexity, many linkages are certain to occur within the SEA at various bottlenecks. These linkages allow movement between large open space areas within the SEA, as well as between areas outside the SEA, such as the Simi Hills and the western extent of the Santa Monica Mountains in Ventura County. The genetic flow through these areas is crucial in maintaining the diversity and viability of certain species within the Santa Monica Mountains. Open space linkages between Kanan Road and Calabasas Parkway along ~~State~~US Route-101, as indicated by the National Park Service, are of particular importance for continued wildlife movement due to the lack of alternative routes and encroachment of development. Although there are significantly large open spaces within the SEA, contiguous habitat linkages between them are critical in reducing bottlenecks and providing for long-term sustainability.

Santa Susana Mountains and Simi Hills SEA

The Santa Susana Mountains and Simi Hills SEA includes several important linkages for wildlife movement. The Santa Susana Mountains and Simi Hills provide a vast open space corridor to foster wildlife movement between the Santa Monica Mountains to the south, San Gabriel Mountains to the east, and the Los Padres National Forest to the north. Dense, natural habitat associated with the majority of the study area provides excellent opportunities for concealment and water sources, while the grasslands provide an abundance of prey.

IV. Significant Ecological Areas

History of the SEA Program

Los Angeles County's Significant Ecological Areas (SEAs) Program has schematic roots in an initial General Plan guiding document, the 1970 Environmental Development Guide, which was adopted as a preliminary General Plan for the County. The Open Space Concept Plan and 1990 Open Space Policy Map depict greenbelt areas and rural lands that reasonably correspond to the current SEA map.

The original Significant Ecological Area Report was prepared in 1972 by scientists from the University of California, Los Angeles, the Los Angeles County Museum of Natural History and other local academic institutions, at the request of the Los Angeles County Department of Regional Planning (DRP). The DRP asked the report authors to identify "significant ecological areas," which due to their high biological resource value, should receive special consideration during the formulation of the 1973 General Plan. In the final report, 81 such areas were mapped and brief descriptions of the value of each were given. The 81 areas were then included on the Vegetation and Wildlife Map in the Conservation Element of the 1973 General Plan.

In 1976, following the 1975 court decision requiring the preparation of a revised General Plan, the DRP and the Environmental Systems Research Institute commissioned the Los Angeles County Significant Ecological Area Study (1976 SEA Study), from the environmental consulting firm, England and Nelson. After excluding the Channel Islands and national forest lands from the study area, the 1976 SEA Study reviewed the data and criteria used to establish the original significant ecological area list, analyzed new information, developed a set of eight criteria to be used to select and prioritize significant ecological areas and concluded with individual maps and descriptions for each. From an initial list of 115 sites, 62 areas met the criteria and were recommended for adoption by the

DRAFT

study. In 1980, 61 of these biologically significant areas were adopted as part of the Conservation and Open Space Element of the Los Angeles County General Plan on the Special Management Areas Policy Map and through individual descriptions of the SEAs in Technical Supplement E of the 1980 General Plan.

In 1991, supplemental studies further assessing the biological resources within seven SEAs were conducted. The Phase I Studies, conducted by Michael Brandman Associates, assessed the following SEA areas: Cold Creek Significant Ecological Area No.9, San ~~Francisquito~~-Francisquito Canyon Significant Ecological Area No.19, Dudleya Densiflora Population Significant Ecological Area No.45, Kentucky Springs Significant Ecological Area No.61, Las Virgenes Significant Ecological Area No.6, Tonner Canyon and Chino Hills SEA No. 15, and Tuna Canyon SEA No. 10. The studies looked at current ownership patterns, existing resources, development pressures and made recommendations into the future management of the SEAs. All of the Phase I studies found either that the SEA boundaries were adequate in size or recommended the expansion of the boundaries to better encompass and protect biotic resources.

In 2000, the DRP commissioned the Los Angeles County Significant Ecological Area Update Study (2000 Update SEA Study) prepared by PCR Services Corporation, Frank Hovore & Associates and Forma Systems. The 2000 Update SEA Study included an Executive Summary, Background Report and twelve biological resources assessments for the Proposed Antelope Valley SEA, Proposed Cruzan Mesa Vernal Pools SEA, Proposed East San Gabriel Valley SEA, Proposed Joshua Tree Woodlands SEA, Proposed Puente Hills SEA, Proposed San Andreas SEA, Proposed San Dimas Canyon and San Antonio Wash SEA, Proposed San Gabriel Canyon SEA, Proposed Santa Catalina Island SEA, Proposed Santa Clara River SEA, Proposed Santa Monica Mountains SEA, and the Proposed Santa Susana Mountains and Simi Hills SEA. These twelve biological resource assessment areas consolidated the 1980 unincorporated area SEAs into larger areas for study and proposed inclusion as SEAs.

The 2000 Update SEA Study also examined the assumptions of the original eight SEA designation criteria from the 1976 SEA Study, modifying one criterion and deleting two. The modification of Class 1 changed the language from “the habitat of rare, endangered, and threatened plant and animal species,” to specify “the habitat of *core populations* of rare, endangered and threatened plant and animal species.” Class 6: “areas important as game species habitat or as fisheries” was removed due to the questionable contribution of these areas towards maintaining biotic diversity. Class 8: “special areas” was deleted due to the vague nature of that designation. The six SEA criteria are contained within this Appendix E, and each SEA description lists which criteria it meets.

From 2001 to 2011, the DRP conducted public outreach, solicited additional recommendations on the SEA boundaries and checked the SEA boundaries with an expert panel of biologists convened in 2010. In 2015, the expanded boundaries that incorporated the collective habitat approach, away from the isolated nodal single resource approach, was adopted.

In 2025, the DRP commissioned the Los Angeles County Significant Ecological Area Update Study (2025 SEA Study) led by the Resource Conservation District of the Santa Monica Mountains. This study analyzed the SEA boundaries with focus on access to open space, climate resilience, rewilding opportunities, and habitat connectivity in response to Senate Bill 1425 (Stern, 2022) and Assembly Bill 1889 (Friedman, 2024). The 2025 SEA Study built upon the 2000 SEA Study by integrating updated biological data, recent scientific research, and field observations to evaluate the continued effectiveness of the SEAs. A central theme of the study was the importance of ecological connectivity for maintenance of biodiversity and ecosystem resilience. This resulted in expansion of existing SEAs

DRAFT

along with consolidations and additions of others. Consolidations included incorporation of the Santa Felicia SEA into the new Liebre-Sierra Pelona SEA, the Rio Hondo College Wildlife Sanctuary SEA into the expanded Puente Hills SEA, and the Valley Oaks Savannah SEA into the expanded Santa Clara River SEA. Additions included the designation of nine new SEAs: Baldwin Hills, Colorado Lagoon, El Dorado Regional Park Lakes, Liebre-Sierra Pelona, Lower Los Angeles River, Pacoima Wash, San Gabriel Mountains, San Rafael Hills, and Sepulveda Basin.

The 2025 SEA Study maintained the six SEA designation criteria and added one new criterion to address habitat connectivity: "Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another." Although wildlife movement has long been recognized as an important component of SEA function, such connectivity has not previously been articulated as a distinct SEA criterion. Consistent with the framework for existing criteria, areas whose primary ecological value is maintaining such connectivity may qualify under this criterion even if they do not independently meet other SEA designation criteria. These areas facilitate species movement, genetic exchange, and other ecological processes where connectivity remains viable or restorable with the surrounding landscape.

SEA Designation Principles

Previously, areas were assigned SEA designations in an attempt to slow or modify the type of development within their boundaries. However, as the County underwent a period of unanticipated growth, many of the SEAs experienced a reduction and/or degradation of their biotic diversity. Appendix E uses the definition of biotic or biological diversity provided by the 1990 U.S. Congressional Biodiversity Act, HR1268, which is defined as a full range of variety and variability within and among living organisms and the ecological complexes in which they occur.

Currently, the design of the SEAs is based on scientifically-grounded concepts regarding size and connectivity. Where feasible, SEAs form linkages between core habitats, which are large blocks of habitat generally conforming to a significant topographical feature, such as a watershed, major river, butte, etc., in order to ensure regional species movement.

Most SEA designations do not focus on a single resource or habitat type and, over time, conservation plans have come to employ a fluid approach to conserving an everincreasing list of sensitive resources (e.g., endangered species, habitats of limited distribution, and "patchy" habitats such as coastal sage scrub). The SEA designations rely on two primary conservation principles: namely that species extinction rates are lower on larger "islands," or blocks of land, than smaller islands; and that isolated habitat areas have less opportunity to regain species by re-colonization from other areas.

Many wildlife species, particularly carnivores and other wide ranging species, require large areas of suitable habitat for genetically and demographically viable populations. In addition, large islands are more likely to encompass diverse habitat types and are more easily buffered against potential impacts from surrounding developed lands. The SEAs are designed to provide habitat linkages between related habitat types (such as the Antelope Valley buttes, or the San Andreas Rift Zone wetlands), by encompassing areas of sufficient width to function as wildlife movement routes between these open space areas.

The current SEA designations provide local resources (such as sensitive species) and their habitats, as well as the seasonal support habitats for those resources, with connections to essential sustaining resource areas (such as corridor areas and hydrological systems). For example, zones of lower

DRAFT

intensity human impacts that exist between essential habitat resources have been included in the current SEA designations, thereby helping to maintain the biotic diversity in the County. The designation of Coastal Resource Area (CRA) is given to those SEAs located with the California Coastal Zone.

SEA Selection Criteria

All of the County's SEAs and CRAs must satisfy at least one of the six SEA selection criteria:

A. The habitat of core populations of endangered or threatened plant or animal species.

Intent of Criterion A: These areas are important in maintaining viable plant and/or animal populations for those species recognized by state and or federal resource agencies as being extremely low in numbers or having a very limited amount of suitable habitat available. The terms "endangered" and "threatened" have precise meanings defined in both state and federal law. The identification of "core population" will be determined by the U.S. Fish and Wildlife Service (USFWS) and the California Department of Fish and Wildlife (CDFW). The term "core population" as used here is a general biological term referring to a known and/or a viable population. Other locations of endangered or threatened plant or animal species may also occur in the County, which are not within an SEA. It should also be noted that the concept of core populations is consistent with current thinking of the USFWS and the CDFW.

This criterion is not meant to constitute a recovery program for listed species, but one element of a more comprehensive conservation effort for the long term sustainment of listed species within the County. At the local level, recovery programs of both the CDFW and the USFWS have measures in place that can impose severe penalties for the "take" of listed species or their habitat.

- Federally Endangered: "any species which is in danger of extinction throughout all or a significant portion of its range...."
- Federally Threatened: "any species which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range."
- State Endangered: "...a native species or subspecies of a bird, mammal, fish, amphibian, reptile, or plant which is in serious danger of becoming extinct throughout all, or a significant portion, of its range due to one or more causes, including loss of habitat, change in habitat, overexploitation, predation, competition, or disease."
- State Threatened: "...a native species or subspecies of a bird, mammal, fish, amphibian, reptile, or plant that, although not presently threatened with extinction, is likely to become an endangered species in the foreseeable future in the absence of the special protection and management efforts required by this chapter." [California Code of Regulations, Title 1, Sec 670.5]

B. On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.

Intent of Criterion B: The purpose of this criterion is to identify biotic resources that are uncommon on a regional basis. The geographical region considered could be as small as the Southern California coastal plains, the Transverse Mountain Ranges, the Mojave Desert, the Southern California coastline, etc. The geographical region could also be as large as Southern California, the Pacific coast, all of California, the western United States, or even larger. The community, association, or habitat is either

DRAFT

unique or restricted in distribution in an area larger than the political boundaries of the County (i.e., coastal sage scrub, native grasslands, or vernal pools). Resources that are limited in distribution in the region being considered, but common elsewhere, are also included under this category.

C. Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.

Intent of Criterion C: The purpose of this criterion is to identify biotic resources that are uncommon within the political boundaries of the County, regardless of their availability elsewhere. The County has a high diversity of biological components. The County and San Diego County are the only counties in the U.S. that possess coastal, montane, and desert subregions within their boundaries. It is a rich heritage that few local governments have an opportunity to preserve.

Many biotic communities that were once common in the County have been severely reduced due to urban and agricultural development. This is especially true south of the San Gabriel Mountains, and among the agricultural fields of the North County. Other biotic features have never been common.

D. Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, migrating grounds and is limited in availability either regionally or in the County.

Intent of Criterion D: Species or groups of species, at various points in their life cycles, tend to congregate in certain areas. These areas possess resources that are essential to the maintenance of specific wildlife species. This criterion is intended to identify those areas that are limited in distribution either regionally or in the County, and not the primary habitat of common species or groups of species.

E. Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.

Intent of Criterion E: Oftentimes scientists learn the most about a biological phenomenon by studying it at an extreme in its distribution. This frequently reveals the biological and ecological parameters under which it can survive. In addition, isolated populations and communities often are relicts of what was present in an area at some previous time, and may show genetic traits not found elsewhere in the species. These biological and ecological parameters may be useful in determining taxonomic relationships.

F. Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.

Intent of Criterion F: The intent of this criterion was to identify examples of the primary biotic resources in the County. At least one example (e.g., native grassland, valley oak savannah) of each vegetation type will be selected from the various geographical regions in the County in order to preserve basic biogeographic diversity.

SEA Descriptions

The following descriptions of the 21 SEAs include descriptions of the boundaries, resources, wildlife movement, and designation criteria for each. More detailed information about the specific plant and animal species of interest for each SEA is contained within the SEA Program Guide, which is

maintained by the Department of Regional Planning. The SEA descriptions, followed by the CRA descriptions, are listed in alphabetical order.

Altadena Foothills and Arroyos SEA

Boundary and Resources Description

The Altadena Foothills and Arroyos SEA is located in the westernmost portion of the San Gabriel Valley. This SEA includes incorporated and unincorporated areas. The SEA represents the lower elevation/urban interface portions of Millard, Alzada, Chiquita, Las Flores, Rubio, and Eaton canyons from the urban edge, to undeveloped wildland areas of the lower elevations of the Angeles National Forest.

The SEA is located within the Mount Wilson and Pasadena United States Geological Survey (USGS) 7.5' California Quadrangles.

On the west side of the Altadena Foothills and Arroyos, the western and southwestern boundaries track along the urban-wildland interface in the undeveloped areas of the Arroyo Seco, Fern, and El Prieto canyons, and the boundary pulls back around a small area of development at the northern-eastern edge of La Cañada Flintridge. A developed area northeast of the junction of Millard Canyon and El Prieto is excluded. The SEA designation includes the undeveloped portions of sub-watersheds of the Arroyo Seco, and also encompasses undeveloped parts of drainages, including Alzada and Chiquita, which flow into the Devils Gate Reservoir of the Arroyo Seco. The Arroyo Seco is within the Los Angeles River watershed. This SEA includes portions of the cities of Pasadena and La Cañada-Flintridge, the unincorporated community of Altadena, and the Angeles National Forest. The eastern side of the southern boundary encompasses undeveloped areas of the sub-watersheds of Las Flores, Rubio and Eaton canyons, which are tributary to the Rio Hondo and historically to the San Gabriel River. Much, but not all, of the Rio Hondo catchment is diverted via flood control channels to the Los Angeles River. The southern boundary of the SEA is bordered by developed properties. The southern boundary moves east along the urban-wildland interface to include undeveloped parts of watersheds, which closely follow the perimeter of Devil's Gate Reservoir, in the Hahamongna Park in Pasadena. From Hahamongna Park, the SEA boundary continues east along the edge of development into the San Gabriel River watershed. The eastern border of the SEA is the eastern ridge of Eaton Canyon near the canyon mouth. A finger of the SEA extends downstream along Eaton Wash to include the Eaton Debris Basin and Reservoir. The northern boundary is formed along ridgelines within the Angeles National Forest that define the catchment of the local canyons. Within the Angeles National Forest, development is much less dense, in the form of in-holdings and Angeles National Forest leases, and is often naturally landscaped, albeit disturbed.

The chief attribute of this SEA is a high diversity of species, which is due to the SEA's position between the mountain biome and the valley biome, caused by an abrupt change of slope formed by the thrust fault complex that borders the San Gabriel Mountains. Furthermore, the SEA has as its center the dividing ridge between the two principal rivers of the Los Angeles Basin, the Los Angeles River and the San Gabriel River.

The wide range of elevation, topography, aspect, and geology represent a diverse array of physical habitats within this SEA. In general, the topography of the SEA is moderately steep to very steep, which results in a number of very narrow corridors with elevations ranging from a high of approximately 2,400 feet above mean sea level (MSL) to a low of approximately 1,200 feet above MSL. Consequently, a variety of plant communities exist, including riparian and upland shrublands and

DRAFT

woodlands. Within these major community types, there are many vegetation series that vary according to plant species dominance.

Of particular note for this SEA is its potential to accommodate lower elevation east-west linkages. This is significant because of the constraints of development at lower elevations, the very steep terrain, and seasonal snow storms above the SEA, beginning at about 3000 feet—all of which limit potential movement for many species. There is also potential for north-south wildlife movement between the Angeles National Forest and the Verdugo Mountains via the Arroyo Seco and the San Rafael Hills. The Arroyo Seco is the eastern limit of this link and creates a potential movement corridor from the Angeles National Forest, over and under the Interstate 210. Across the Interstate 210, the linkage enters the San Rafael Hills, where blocks of habitat remain in the cities. Some are conserved in natural open space, such as the Cherry Canyon Park and Open Space Preserve of the City of La Cañada-Flintridge, just south of the County Descanso Gardens. These open spaces are interspersed with residential development and are not part of the SEA. From the San Rafael Hills, linkage potential may be traced to the west across State Route 2 and Verdugo Wash, past enclaves of residential development to access the Verdugo Mountains.

Wildlife Movement

Wildlife movement within the SEA takes on two major forms. First, due to the extremely steep intervening topography, considerable movement of wildlife up and down the drainages, which course through this SEA to connect the forest interior with foothill areas, is expected. Consequently, this type of movement occurs on a seasonal basis, particularly for large mobile mammals that typically meet their full range of habitat needs over broad areas.

The second major type of movement occurs across the flanks of the foothills in an east-west direction. Particularly for riparian-obligate and riparian-favoring migratory birds, the corridor linking lower elevation riparian habitats in the SEA are of high importance and heavily utilized.

Regional Biological Value

The SEA meets important SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE ALTADENA FOOTHILLS AND ARROYOS SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Not Met	None within this SEA.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The SEA is designating one of the principle ecotones of the Southern California coastal areas: the area where the sediment of the coastal alluvial fans from the mountain streams and drainages is exiting the abrupt upthrust rock of the mountains. Here one finds the biotic communities of the mountains meeting the communities of the coastal plain areas, combining with the organisms that are only found at the

DRAFT

	Criterion	Status	Justification
			junction. The natural habitats of this kind of biological area are fast dwindling as urban communities expand to the limits of easily buildable space.
G)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The SEA is designating one of the principle ecotone areas of the County coastal exposure: the area where the sediment of the alluvial fans from the mountain streams and drainages is adding to the mile deep sediments of the Los Angeles Basin, as the watercourses exit the abrupt upthrust rock of the San Gabriel Mountains. It is an area where one can often encounter flora that is characteristic of the Peninsular Ranges to the south and flora of the coastal ranges and Sierra Nevada to the north, among typical flora of the Transverse Ranges. The SEA contains prime examples of coastal sage scrub and other kinds of chaparral, riparian oaks, woodlands of the canyon oak of the mountains, woodlands of the coast live oak, which occurs both in the lower mountains and the valleys, good stands of the San Gabriel endemic oak (<i>Quercus dumosa</i> var. <i>gabrielensis</i>), diverse and beautiful flora characteristic of the continually changing beds of the mountain streams, both perennial and intermittent, and the wildlife that reside in these various habitats.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The SEA provides a low elevation constrained corridor. The SEA serves as the only corridor to provide interacting component habitat areas for species to feed, rest, and migrate from low basin and foothill elevations to the sub-alpine elevations of the high San Gabriel Mountains.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Not met	None within this SEA.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Areas encompassed within the SEA represent the only remaining stands of low elevation foothill scrub, chaparral, and canyon woodland communities within the north San Gabriel Valley. These communities once extended throughout what are now the communities of the north San Gabriel Valley, bridging the transition between high chaparral on the southern slope of the San Gabriel Mountains to the alluvial fans extending beneath the mountains to the coastal basin.

In conclusion, the area is an SEA because it contains (B - C) a good example of the biotic communities typical of the area where the abrupt upthrust of the mountains meets the alluvial fans of the valleys, a natural habitat that is limited in availability in the County and the coastal Southern California region; (D) it has a constrained connective corridor area near the Devil's Gate Dam where the freeway

~~underpasses provide access between the San Rafael Hills and the San Gabriel Mountains; and (F) it supports intact remnant stands of low-elevation chaparral and scrub communities that were once more widespread within the region.~~

Antelope Valley SEA

Boundary and Resources Description

~~The Antelope Valley SEA is located in the central portion of the Antelope Valley, primarily east of the cities of Palmdale and Lancaster, within a predominantly unincorporated area of the County. The SEA is focused on the principal watercourses of the area: Little Rock Wash and Big Rock Wash and tributaries, such as Mescal Creek. Audubon California recognizes the area of Edwards Air Force Base as a Globally Important Bird Area (IBA), which is visited by tens of thousands of migrant birds during the spring and fall migratory seasons, and supports the breeding of rare and endangered birds during the spring and summer months.~~

~~The SEA is located, at least partially, in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Rosamond, Rosamond Lake, Redman, Rogers Lake South, Jackrabbit Hill, Lancaster East, Alpine Butte, Hi Vista, Adobe Mountain, Palmdale, Littlerock, Lovejoy Buttes, El Mirage, Pacifico Mountain, Juniper Hills, Valyermo, and Mescal Creek.~~

~~Watercourses and water features, such as dry lakes and springs, are the focus for desert wildlife and central to connectivity and biodiversity in this region. The SEA was delineated to emphasize the importance of the Little Rock Wash and Big Rock Wash watersheds to the surface and subsurface hydrology of the Antelope Valley and to the dry lakes. The western portion of the SEA extends along the margin of the Little Rock Wash and floodplain zone, while the eastern margin follows a tributary of Big Rock Wash, which is Mescal Creek Wash and its tributaries. The origins of the watercourses in the Angeles National Forest are an important aspect of their diversity and connectivity, and the importance of the diverse forest vegetation of this SEA is discussed below. The SEA includes several major buttes and numerous minor ones, which have highly diverse biota along with diverse desert habitats, which range from sand dunes formed from the wind-blown dust that the buttes collect, to rocky crags, which are home to various raptors. The SEA includes the County's portion of the watershed basin for dry lakes, which are the destination for the watercourses. There are three dry lakes and their adjacent plains (protected as part of Edwards Air Force Base) included in the SEA: Rosamond Dry Lake with the adjacent Piute Ponds, Buckhorn Lake, and Rogers Lake. These lakes and ponds are often flooded during the rainy winter-spring seasons, and are the principal resting areas in the region on the Pacific Flyway. The northeastern portion of the SEA encompasses some agricultural cropland (portions of which are fallow) and dispersed rural residential uses; however, the underlying hydrology of the washes remains intact throughout the entire SEA.~~

~~Three main watercourse segments originate in the San Gabriel Mountains and flow through the Antelope Valley to dry lakes near the northern County boundary: 1) Little Rock Wash; 2) Big Rock Wash; and 3) Desert Montane. Desert Montane centers on Mescal Creek and includes adjacent drainages. The flows of all three drainages are subsurface for much of the year and may be on the surface during rain and snowmelt.~~

~~The Little Rock Wash segment (the westernmost segment), goes north from Little Rock Palmdale Dam as its southern barrier. Upstream from the reservoir is critical habitat for the endangered arroyo toad (*Anaxyrus californicus* FE, SSC). The toad could occur from time to time in the downstream area of the SEA. Heading north to Mount Emma Road, the boundaries follow the flood zone of the Little Rock~~

DRAFT

Wash and also incorporate some of the vegetated slopes that drain to the wash. North from Mount Emma Road, the boundaries generally follow Federal Emergency Management Agency (FEMA) boundaries. On the west side, south of Edwards Air Force Base and north of Avenue F, the SEA boundary follows the Economic Opportunity Area boundary.

All of Edwards Air Force Base that is in the County is included in the SEA because the restricted entry and use protect the dry lakes and their neighboring areas. Many desert plants and wildlife species once found broadly across the Antelope Valley are now found only or primarily within Edwards Air Force Base. The ponds and dry lakes have distributed habitat of marshy alkali grassland, alkali flats, and cattail and bulrush marsh augmented by wastewater treatment facilities that have additional ponds. Some of the nesting rare and uncommon birds include white-faced ibis (*Plegadis chihi*), tricolored blackbird (*Agelaius tricolor*), redhead (*Aythya americana*), gadwall (*Anas strepera*), yellow-headed blackbird (*Xanthocephalus xanthocephalus*), least bittern (*Ixobrychus exilis*), and federally threatened western snowy plover (*Charadrius alexandrinus nivosus*).

The Big Rock Wash area has western and eastern segments in the SEA. The western arm of the Big Rock Wash segment begins near the northern boundary of the Angeles National Forest, heads north out of the Forest along Pallett Creek. The SEA includes parts of Cruthers and Holmes creeks near their junctions with Pallett Creek. SEA boundaries follow the braided stream channel toward the confluence with Big Rock Wash. From the aqueduct at Big Rock Wash to Edwards Air Force Base, the western boundary line follows recently active braids of Big Rock Wash, encompassing Alpine Butte, and joining to the Little Rock Wash segment within Edwards Air Force Base. On the eastern arm of the Big Rock Wash segment, the SEA boundaries head north from the Angeles National Forest headwaters of Dorr Canyon (a Big Rock Wash tributary) and the headwater area of Big Rock Wash near State Route 2. The boundaries travel through the Angeles National Forest and follow the wash area of the streams toward their confluence with Pallett Creek. The Angeles National Forest floodplain of the widened area of South Fork of Big Rock Wash is included in the SEA.

South Fork of Big Rock Wash is part of the federally-designated critical habitat of the mountain yellow-legged frog (*Rana muscosa*, FE, SE). This frog is known in the County from only a few high mountain streams in the San Gabriel Mountains. A fungal pathogen is principally responsible for its decline; however, climate change, air pollution and non-native predators are also likely contributing factors.

Another broad area of the San Andreas Fault Zone near the Valyermo Ranch follows the FEMA boundaries and includes a nesting area for gray vireos near Bobs Gap. Between the Angeles National Forest and the aqueduct, the SEA boundaries follow FEMA boundaries. The eastern boundary generally follows the FEMA boundary and recently active braids along the main course of Big Rock Wash to the vicinity of Avenue Q East, at which point it projects east to encompass Lovejoy Buttes. At Avenue O, the eastern boundary rejoins the main active portion of Big Rock Wash, continuing northeastward to skirt development in Lake Los Angeles. In the vicinity of Avenue M, the boundary projects eastward from about 156th Street East to 180th Street East) to encompass Rocky, Piute, and Saddleback Buttes, and connect with the Desert Montane transect segment.

The Desert Montane Transect segment begins in the Angeles National Forest along the ridge of Table Mountain at the San Bernardino-Los Angeles County line. Table Mountain is known for its diverse flora, which includes desert and mountain elements, and some unusual limestone-obligate species. The SEA southern boundary along the ridgeline meets the western boundary as it skirts the camp developments along the southern base of Table Mountain. The boundary turns north along the western ridge of the Mescal Creek drainage, crossing the California Aqueduct with the State Route 138. From the aqueduct to Avenue R, the western boundary buffers the westernmost portion of the drainage by

DRAFT

~~200 feet, protecting the braided area of the watercourse. This part of the SEA includes Black Butte and the Three Sisters Buttes, and many smaller unnamed buttes, as well as Mescal and Theodore Payne County wildlife sanctuaries. The east side of the transect is the San Bernardino-Los Angeles County line. At about Avenue U East, the eastern boundary veers off the San Bernardino-Los Angeles County line to the north-northwest, buffering the Puzzle Creek watercourse by about 200 feet, protecting the braiding of the easternmost drainages. Near Avenue R, the boundary trends north, and goes north-northwest near Avenue P to include Moody Butte, lesser unnamed rises, and Blue Rock Butte.~~

~~The Desert-Montane segment largely avoids drainages that flow into and out of the Lake Los Angeles community, but the transect includes diffuse watercourses on the south side of Saddleback Butte, Saddleback Butte and the surrounding Saddleback Butte State Park, the Antelope Valley Indian Museum State Park at the base of Piute Butte, and Piute Butte. At about Avenue H and 170th Street East, the boundary turns to the northeast following natural vegetation to the County boundary near Avenue C. Here the boundary turns north along the line to where San Bernardino, Kern and Los Angeles counties meet. This northeastern part of the SEA has WEMO conservation areas for the threatened desert tortoise and state-threatened Mojave ground squirrel. The northeastern area has some BLM land and the County Phacelia Wildlife Sanctuary, which is also County Wildflower Preserve A. The SEA includes large parts of County Wildflower Preserve F.~~

~~On Edwards Air Force Base, north to south between Avenues B and E East, and west to east between 140th Street East and the San Bernardino-Los Angeles County line, there is federally designated critical habitat for the state and federally-threatened desert tortoise (*Gopherus agassizii*). At 190th Street, the critical habitat widens to extend north beyond the County and the SEA into Kern County. At 200th Street, the critical habitat widens to the south to extend to Avenue H and then goes east across the San Bernardino-Los Angeles County line. The desert tortoise critical habitat area on Edwards Air Force Base is included in the SEA, and much of the SEA area north of Avenue H in the eastern drainages of the SEA is designated critical habitat for the tortoise.~~

~~The SEA traverses the Antelope Valley from the foothills of the San Gabriel Mountains, to the low elevations of the dry lake basins, and its expanse and considerable topographical relief is reflected in its relatively high floral and faunal diversity. The SEA includes playa lake, alkali marsh, alluvial fan scrub, a mosaic of xeric desert scrubs, Joshua tree woodland, desert riparian woodlands, juniper scrub, pinyon pine, chaparral and mixed conifer, oak, and riparian communities of higher elevations. Transitional zones (ecotones) between these communities often contain unusual species compositions, such as pinyon pine, juniper and Joshua trees together, or Joshua trees adjacent to cottonwood forest.~~

~~Edwards Air Force Base has the only good stands of mesquite (*Prosopis glandulosa*) remaining in the County. It has areas of Mojave spineflower (*Chorizanthe spinosa*), creosote bush scrub, alkali sink, and the transition vegetation between the two. Rosamond Lake has the best example of the shadscale scrub and alkali sink biotic communities in the County. Shadscale scrub needs heavy soil with underlying hardpan between 3000-6000 feet elevation, which is unusual in the County, and more common in the north Mojave Desert and Owens Valley. In addition, the playa has the southernmost extension of the Great Basin kangaroo rat (*Dipodomys microps*), which is an isolated geographic population of scientific interest.~~

~~The southernmost portions of the three “legs” of the SEA lie within the Angeles National Forest, and include the upper tributary watersheds and streams for Little Rock Wash, Big Rock Wash, and Mescal Creek. These areas support multi-species oak and conifer woodlands that are common to the middle-~~

DRAFT

elevation zones on the north face of the San Gabriel Mountains. The creeks are higher energy systems at those elevations, as they collect water from the surrounding terrain, and are typically lined with woodlands of alder, willow, sycamore and cottonwood, with varying densities and with various compositions of species.

As the creeks drop north of the pressure ridges of the San Andreas Fault Zone, they lose gradient and widen, and most of the flow becomes sub-surface, except during high energy storms or in the spring (depending upon rainfall totals in the watersheds). The vegetation becomes sparser and less evenly distributed along the channel margins. Crossing the lowlands of the Antelope Valley, the channels support a variety of desert scrub vegetation within the alluvial plains. Where the alluvial plains are wide and shallow, cottonwood-willow woodland and sycamore woodland vegetation communities often occur within the overall floodplain on stable terraces; around oxbow flow zones in the Antelope Valley; or where the groundwater table is replaced or augmented by agricultural runoff. The surrounding upland habitats are primarily desert scrubs, including creosote and chenopod scrubs, sand sheets (chiefly around the buttes), and Joshua tree woodland. Intact Joshua tree woodland, with native understories present, supports a relatively high diversity of annual wildflowers, reptiles and mammals. The Joshua trees also provide nest sites for many resident and migratory bird species.

Lovejoy, Alpine, Piute, Black and Saddleback buttes, along with other, smaller unnamed buttes, form most of the topographical relief within the SEA. These areas offer different ecological conditions that are associated with rock shelter, perching sites, nesting sites, denning areas, wind protection and sand sheet accumulation areas. Local and migratory bat species roost and reproduce in the caves and crevices of the butte formations. The higher buttes provide local nesting sites for owls and other birds of prey.

Alpine Butte is the least disturbed butte in the County, with excellent stands of Joshua tree woodland and creosote bush scrub, and impressive wildflower displays when rainfall creates appropriate conditions. Lovejoy Butte has Joshua tree woodland and creosote bush scrub, with a central wind-blown sand community for a good mixture of rock and sand habitats. In addition, the close proximity of Lovejoy Butte to Big Rock Wash increases the diversity of habitats in the area. Nevertheless, it also suffers from impact from the Lake Los Angeles community, which borders the butte on three sides. The clustering of buttes in the SEA may be important to the abundant, diverse wildlife that inhabits the various vegetation communities around and in the buttes. Saddleback Butte and Piute Butte together are protected as a state park, but Saddleback Butte is also subject to development for campsites and hiking trails. Piute Butte has a prehistoric site that may protect it from much future recreational development. All of the buttes harbor diverse wildlife and flora. Most of them are critical habitat for the state and federally threatened desert tortoise. Some buttes within the desert tortoise's critical habitat are not included in the SEA.

The active and fallow open agricultural lands support a diversity of wildlife species, which essentially regard the fields and ditches as irrigated desert. Birds of prey frequently hunt over the open agricultural areas, including fallow fields; wide-ranging predators also find excellent hunting conditions in and around agricultural areas. A spectrum of local and migratory bat species feed over the irrigated fields in the spring and summer, when insect numbers are the highest, and at least one sensitive bat species, the pallid bat, forages in open scrub or ruderal desert habitats.

The northern portion of the SEA contains several unique habitat types, including mesquite bosque (threatened locally by lowering water tables and harvest for firewood), clay pan pools, vernal pools, alkali grasslands, alkali and freshwater marshes, and permanent ponds. Hundreds of bird species have been recorded from the pond and marsh habitats around the dry lakes and ponds, and numerous

DRAFT

species nest on the playa margins or in the associated riparian habitats. The open creosote scrub and other xeric habitats on the slopes surrounding the lake playas serve as important wintering areas for many raptor species, as well as large numbers of songbirds.

Wildlife Movement

The SEA extends from the Angeles National Forest to the playa lakes within Edwards Air Force Base, encompassing most of the two largest drainages exiting the northern slope of the San Gabriel Mountain range. The geographical features of the SEA serve as a major habitat linkage and movement corridor for all wildlife species within its vicinity and in an intergenerational sense, many of the plant species. Ecologically generalist species (mountain lion, bobcat, coyote, gray fox, etc.) have the ability to move across such vast areas and through changing habitat types. For such species, the SEA may serve as an important system for long term and genetic exchange among populations. For smaller or less mobile species or taxa, which are narrowly restricted in their habitat needs, the SEA can serve as a broad linkage zone, in which individual movement can take place during seasonal population dispersal or over generations. This provides essential genetic exchange within and between metapopulations. The two drainages, combined with the upland terrestrial Desert Montane transect portion of the SEA, ensure linkage and direct movement areas for all of the wildlife species present within the County portion of the Antelope Valley.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE ANTELOPE VALLEY SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	Critical habitat for the only known Antelope Valley population of the federally endangered arroyo toad is adjacent to Little Rock Reservoir, upstream in Little Rock Creek, and some may still be found downstream of the dam in the SEA. The SEA encompasses much of the County ranges of the federally threatened California desert tortoise, including much of the County critical habitat for the tortoise. The state threatened Mohave ground squirrel occurs throughout much of the SEA. The SEA includes some of the critical habitat of mountain yellow legged frog in the South Fork of Big Rock Creek. It includes habitat designated in the Western Mojave Plan (WEMO) for the alkali mariposa lily, which is a rare lily of the desert floor.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The mesquite bosque, sand sheet, rocky butte, desert riparian woodland, and alluvial fan sage scrub habitats are unique and regionally restricted biotic communities encompassed by the SEA. Desert species not, or rarely, found elsewhere in the County, such as verdin, black throated sparrow, Mojave rattlesnake, desert banded gecko, Leech's prionid borer, and mesquite borer, occur within these habitats. Additionally, the ponds and other riparian and wetland systems in the northern portion of the SEA support numerous water birds and raptors not found elsewhere in the County.

DRAFT

	Criterion	Status	Justification
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The desert alluvial fan sage scrub, Joshua tree woodland, desert riparian woodland, mesquite bosque, alkali meadow/marsh, desert freshwater marsh, playa-lake and seasonal pool habitats are located within, are unique to, or best represented within, the SEA.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, migrating grounds and is limited in availability either regionally or in the County.	Met	The freshwater habitats within and around Rosamond, Buckhorn and Rogers dry lake basins have large concentrations of migratory and resident waterfowl and birds of prey, providing them with essential seasonal and permanent resources. The rocky desert buttes are unique roosting, sheltering, perching and nesting sites for birds of prey and bats. This SEA is centered on migratory routes for both plants and animals along principal desert washes and buttes that connect the mountains to freshwater playas.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The mesquite bosque that is located within the SEA is clearly at an extreme of its geographical range, along with its associated biota, such as the mesquite borer. Edge populations usually represent an unusual genetic variation in a population or community, and therefore meet the criterion of scientific interest as well as the criterion of a population at the extreme physical/geographical limit of its range.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The SEA encompasses some of the most biotically intact acreages of Joshua tree woodland, desert riparian woodland, and desert alluvial fan sage scrub remaining in the County. Mesquite was formerly widely distributed in the Antelope Valley, but due to harvesting and drawdown of groundwater, is now limited to a few protected areas, such as the Edwards Air Force Base.

In conclusion, the area described is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) populations of scientific interest at the edge of their range including the desert tortoise, the mesquite bosque, and the Mojave ground squirrel; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Cruzan Mesa Vernal Pools SEA

Boundary and Resources Description

The Cruzan Mesa Vernal Pools SEA lies north of the Santa Clara River southeast of Bouquet Canyon. The SEA boundaries encompass the watershed and drainages of the Cruzan Mesa and Plum Canyon vernal pools, which are considered as a single ecosystem within the SEA. Vernal pools are a rare habitat. They are floored by clay or other substrates that retain the water of spring rains for a period

DRAFT

~~of months. The pool dries primarily by evaporation. The fauna and flora are adapted to this regimen, which often forms propagules that can last through years of desiccation. The pools are ephemeral, and may not fill every year. The fauna may be specific to an area of pools. Most biota is spread from pool to pool in mud, which adheres to the feet of birds and other animals that visit the pools.~~

~~The SEA is located within the unincorporated area of the County and lies entirely within the United States Geological Survey (USGS) 7.5' California Mint Canyon Quadrangle.~~

~~The SEA boundaries generally follow sub-watershed boundaries and occasionally extend beyond the sub-watershed boundary into areas to encompass vegetation, such as coastal sage scrub. The vernal pools in the northwest segment of the SEA are bounded in part by a sub-watershed boundary, but are primarily based on land formations that would adversely affect the vernal pools if altered to a great degree. This segment encompasses the vernal pool upland areas that contribute to the biological function of the vernal pool ecosystem. This SEA area includes most of the local federally designated critical habitat for the threatened vernal pool plant, spreading navarretia (*Navarretia fossalis*, FT).~~

~~The SEA includes mesas, canyons and interior slopes, with Plum Canyon Creek running east west through the southern portion of the SEA. The extent of the SEA encompasses the watershed, which supports both regionally unique vernal pools, including their immediate watersheds and the corridor between them. Plum Canyon forms the major drainage running east west through the southern portion of the SEA, which drains west toward Bouquet Canyon. Uplands within the SEA are comprised of slopes and canyons, which support coastal sage scrub or scrub chaparral vegetation. The Cruzan Mesa vernal pool complex lies within an elevated, topographically enclosed basin atop an eroded foothill between Mint and Bouquet canyons. Plum Canyon vernal pool, which is situated in a landslide depression on a hillside terrace, is smaller than the Cruzan Mesa pools, but possesses the same essential vernal pool characteristics as the larger system, and the two areas together form an ecologically functional unit.~~

~~The seasonally wet vernal pools and surrounding open coastal sage scrub and chaparral slopes support migrant and resident birds, other native sage scrub vertebrate species, and a number of sensitive taxa, including Riverside fairy shrimp, western spadefoot toad, ashy rufous-crowned sparrow, Bell's sparrow, and possibly coastal California gnatcatcher. The steep cliffs that surround the Cruzan Mesa vernal pools, especially along the southeast and north margins, provide protected sites for perching, roosting and nesting by a variety of birds of prey, including great horned owl, barn owl, red-tailed hawk, prairie falcon, and golden eagle. The Plum Canyon vernal pool is hidden from view from the Plum Canyon roadway, and receives little human attention. Trash dumping, shooting and off-road vehicle activities have occurred within a few meters of the margin of the pool basin, but the pool shows only limited evidence of human intrusion.~~

Wildlife Movement

~~The vernal pools situated within this SEA serve as isolated, high quality, habitat linkage sites for migratory waterfowl. The vernal pools teem with arthropod and amphibian activity, and provide essential feeding grounds for long-distance migrants, as well as for resident species of reptiles, birds and mammals. The ponds do not lie within any identified terrestrial movement routes for wildlife, but may serve as important seasonal watering sites for species moving through and across the Plum Canyon divide between Mint and Bouquet canyons. The Plum Canyon stream channel undoubtedly serves as a movement pathway for more mobile species of terrestrial mammals, but it no longer links any larger habitat areas directly due to land conversion in Mint and Bouquet canyons.~~

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE CRUZAN MESA VERNAL POOLS SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	The only known County populations of the federally endangered Riverside fairy shrimp, and the state and federally endangered California Orcutt grass, are found in the vernal pools within the SEA. The federally threatened plant, spreading navarretia, also occurs in these pools, and the pools and much of the SEA are designated critical habitat for this plant.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The Cruzan Mesa and Plum Canyon vernal pools are regionally unique biotic communities with several plants found only in such habitat types. The pools support the Riverside fairy shrimp, western spadefoot toad, and at least one vernal pool endemic ground beetle.
C)	Within County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The Cruzan Mesa and Plum Canyon vernal pools are unique biotic communities within the County, with several plants only found in such habitat types. The pools support the Riverside fairy shrimp, western spadefoot toad, and at least one vernal pool endemic ground beetle species.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The vernal pools serve as concentrated breeding areas for several species of amphibians, including the sensitive western spadefoot toad. They also attract a diversity of waterfowl seasonally—chiefly species migrating through the area—which use the pools for resting and feeding. While other open water systems attract and support waterfowl, the vernal pools are located in remote, upland sites, away from other freshwater features.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Not met	Although the SEA does not contain biotic resources that are clearly an extreme in physical/geographical limitations, or represent unusual variation in a population or community (and therefore does not meet this criterion), it is of scientific interest due the extreme rarity of vernal pool communities, especially in the County, and due to their susceptibility to drought and climate change.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic	Met	Protection and sensitive management of the Cruzan Mesa and Plum Canyon vernal pools would preserve one of the few examples of such habitat type in the County.

Criterion	Status	Justification
communities in the County.		

The SEA meets several designation criteria and supports several regional biological values. These values include A) sensitive plant species unique to seasonal pools on heavy clay soils—several of which are at the northernmost point in their overall ranges, and seasonal surface water, which provide breeding sites for sensitive amphibians, including western spadefoot and Riverside fairy shrimp; B-C) a concentration of vernal pools, found in few places in the County and rare in the region, and their coastal sage scrub watershed, which serves as a hydrological filter, and seasonal ponds and surrounding mesic vegetation; D) vegetation that provides essential foraging and wintering sites for migrating birds that are otherwise uncommon on the coastal side of the San Gabriel Mountains, and steep cliffs surrounding the mesa tops with crevices and cavities providing roosting and nesting sites in the otherwise brush covered hillsides; E-F) Rare habitat in Southern California, and the support of sensitive resources that are unique locally and regionally, and that biologists consider to be among the most sensitive habitat types in Southern California.

East San Gabriel Valley SEA

Boundary and Resources Description

The East San Gabriel Valley SEA is located in the eastern end of the San Gabriel Valley. The SEA includes some of the low hills that rim the San Gabriel Valley on the east. The SEA has several natural, interacting components that constitute an area wide ecological unit. The SEA consists of five units. The location and configuration of this SEA and its parts are primarily defined by the urbanization of the eastern San Gabriel Valley, which has occurred over the more developable valley floor and lower slopes of the San Jose Hills. As a consequence, the SEA resembles an "archipelago" that encompasses portions, or islands, of undeveloped ridgelines, hilltops and drainages between the San Gabriel Mountains to the north and the Puente Hills to the south.

Generally, the topography within this SEA consists of moderate to steep hillsides with north, south, east and west slope aspects. Ridgelines vary in width, from narrow to broad, with well defined drainages in between. One major drainage, Walnut Creek, and a man-made reservoir, Puddingstone Reservoir, are found within this SEA. Elevations range from a low of approximately 560 feet above Mean Sea Level (MSL) in the Walnut Creek drainage, to a high of approximately 1,375 feet above MSL at Buzzard Peak.

The SEA has several ridgelines and hilltops and the major drainage of Walnut Creek at the northern side of the San Jose Hills. In the last half of the 20th century, these areas have been surrounded by urban development. Units 1-5 include 1) South Hills Park and the surrounding undeveloped land in the City of Glendora north of Interstate 210; 2) The natural riparian section of Walnut Creek County Park. The SEA includes sections in the City of Covina and City of San Dimas; 3) Frank G. Bonelli Regional County Park (Bonelli Park); 4) Buzzard Peak, which is an extended ridge of undeveloped slopes to the west of Bonelli Park within the City of San Dimas, City of Walnut, and City of West Covina; and 5) Elephant Hill and an adjoining ridgeline in the City of Pomona. Along most of its boundaries, the SEA is bordered by developed properties. Large parts of this SEA are designated critical habitat for the federally threatened coastal California gnatcatcher (*Poliioptila californica californica*).

DRAFT

In its entirety, the SEA is located within the San Dimas, Glendora and Baldwin Park U.S. Geological Survey (USGS) 7.5' California Quadrangles.

Unit 1: The northernmost section of the SEA is the South Hills unit. Much of the SEA is "protected" in the South Hills Park of the City of Glendora. The SEA lies north of the Interstate 210, has its northwestern boundary along Big Dalton Wash (a concrete-lined flood control channel), and northeastern boundary along existing residential development in the floodplain of the wash and adjoining hills. The eastern boundary is also bordered by existing residential development. The southern boundary consists of the Interstate 210 right of way, and the western boundary lies along Glendora Avenue. Residential neighborhoods on the periphery (and not inside) of the South Hills Park are excluded, while external natural habitat is included in the SEA. An area of nursery fields in a central position of the park is included, as it is less than 40 acres in extent. Toward the eastern side of the SEA, there are some estate residences on the hills that are included, as they incorporate much natural habitat; however, denser neighborhoods are excluded from the SEA.

The South Hills are an area of chaparral and grasslands, and a northern stepping stone for wildlife (chiefly aerial) that move along the spine of the San Jose Hills between the San Gabriel Mountains of the Transverse Ranges, and the Puente and Chino Hills of the Peninsular Ranges on the southern side of the San Gabriel Valley. In South Hills Park, there are fine stands of walnut woodland in the upper elevations; northern ravines of oak woodland with walnut woodland in the upper elevations; what remains of the coast live oak woodland that once bordered the Big Dalton Wash; and excellent stands of prickly pear cactus covering some of the slopes. There are also grasslands with a dominant component of introduced mustards and some rocky exposures. Coastal sage scrub with alternating dominant components of California sagebrush (*Artemisia californica*), sumacs (*Rhus* spp.) and elderberries (*Sambucus* spp.) cover many of the slopes. Big Dalton Wash is channelized, and much development has been inserted into what was formerly Big Dalton's riparian oak woodland. This residential area forms a border for the northeast area of the SEA.

Unit 2: The westernmost portion of Walnut Creek begins where South Reeder Avenue crosses the Walnut Creek drainage on the border of the City of Covina and City of San Dimas. Walnut Creek runs along the northern base of the San Jose Hills and is a riparian interface between the rolling hill habitat and the coastal plain of the San Gabriel Valley. This segment encompasses the undeveloped creek bottom and flood plain, which is a relatively natural area along and adjacent to Walnut Creek, most of it preserved as Walnut Creek Park. The Creek and lower slopes of the San Jose Hills contain one of the best riparian oak woodlands in the County, with a mixture of ash, walnut, willows, and sycamores. The upper areas of the oak woodland transition into excellent examples of walnut woodlands. Traveling east from this point, the SEA is bordered to the north by residential areas on the plains and to the south by residential areas on the ridgelines above the Creek. Much of the habitat of interest is on the slope of the ridgeline above the Creek.

Unit 3: The Walnut Creek Park unit is connected to the Bonelli Park unit. The SEA continues east to the Interstate 210, where the drainage and the SEA underpass along with San Dimas Avenue. There is a fenced footpath on the south side of the underpass that wildlife and humans use extensively. On the east side of the Interstate 210, the boundaries diverge and follow the freeway right of way to the north and to the south. At the edge of natural habitat about a 0.2 mile north, the boundary turns northeast with Walnut Creek. There are two fingers of creek tributaries: one encircles its stream course as far as the Arrow Highway, and the other includes the undeveloped north facing slope of a low ridge that has residential development along the top. The SEA ends at a Union Pacific rail line. The edges of Walnut Creek have business parks and industrial properties, as well as residential areas.

DRAFT

The SEA boundary returns back west from the two fingers to circle around the west side, and excludes the Raging Waters theme park. From Raging Waters, the SEA boundary follows along the northern boundary of Bonelli Park, which is the largest unit of the SEA. This area of Bonelli Park includes the 250-acre Puddingstone Reservoir, which is a flood control basin created by damming the main course of Walnut Creek. Puddingstone Reservoir is home to many resident birds, including a large population of American coots (*Fulica americana*). Notably, Puddingstone Reservoir also hosts a great variety and number of migrating waterfowl and other birds during the spring and fall, and provides riparian habitat for birds and other animals that are riparian obligates. Following along Puddingstone Drive to the Puddingstone Channel, the SEA eastern boundary turns south to conform to the eastern boundary of Bonelli Park. Within Bonelli Park, developed areas include the Mountain Meadows Golf Course on the west side of the Park, the campground area for recreational vehicles, the northern parking area of Puddingstone Reservoir, and the developed south side of Puddingstone Reservoir. On the south side of the park, the boundary borders the Interstate 10 to the junction with State Route 57, and then turns north following the east side of State Route 57, back to the undercrossing of San Dimas Avenue.

Bonelli Park (with the exception of the built areas above), Puddingstone Reservoir, and Raging Waters are critical habitat for the coastal California gnatcatcher. There are a number of breeding pairs that occur in natural areas of coastal sage scrub in Bonelli Park. Another species of note is the many-stemmed dudleya or live forever (*Dudleya multicaulis*), which is an uncommon plant found on the picturesque rocky outcrops of the Glendora volcanics formation. This plant is at its northernmost extent in this area of the San Gabriel Valley, and is more common in Orange County.

Unit 4: Across the State Route 57 from Bonelli Park, the San Jose Hills continue southwest, with residential areas dispersed among intervening grassland and chaparral natural areas. This area is considered an extension of the Buzzard Peak ridgeline. The natural area that continues the habitat in Bonelli Park is roughly a central area of a north-facing wooded area and a southern area of a ridgeline with steep slopes along the Interstate 10. The SEA is chiefly on the undeveloped upper ridge areas, and critical habitat for the coastal California gnatcatcher closely follows the SEA in the southern part along the ridgeline and the Interstate 10. The SEA and critical habitat for the gnatcatcher cross the highway with the ridgeline, and extend to Buzzard Peak

Buzzard Peak and its associated lands begin in the east along a ridgeline that is immediately north of the California State Polytechnic University (Cal Poly), Pomona campus. The western section of the SEA is undeveloped, but may in the path of future growth. This component follows the ridgeline west to where it crosses Grand Avenue and continues in a westerly direction, which encompasses a series of minor ridgelines and drainages with highly dispersed residential areas. Moving east to west, its northern boundary is marked by Interstate 10, developed portions of the Forest Lawn Memorial Park Covina Hills and residential areas. Moving from east to west, its southern boundary is marked by developed and cultivated areas of the Cal Poly campus, Amar Road and residential areas. Along their entire lengths, the northern and southern boundaries follow highly circuitous alignments at the edge of developed landscape. A lobe of the SEA extends into the natural area that separates Cal Poly Pomona from Mt. San Antonio College. Gnatcatcher critical habitat covers most of this segment of the SEA, and extends beyond Grand Avenue as far as State Route 39 (Azusa Avenue), with some islands of critical habitat in nearby hilltops.

Unit 5: A final component of this SEA is located south of the intersection of State Route 71 and State Route 57 in the City of Pomona, at a ridgeline that is bordered chiefly by development that has an undisturbed remnant of the original habitat: north-facing slopes of oak and walnut woodland. South-facing slopes consist of a mixture of dispersed chaparral, coastal sage scrub, and grasslands. There is a lobe that is north of Mission Boulevard in the City of Pomona on Elephant Hill, and a lobe south

DRAFT

of Mission Boulevard. On Elephant Hill, the northern boundary is the Metrolink track. The eastern boundary is bordered by industrial development; the southern boundary is Mission Boulevard in the southeast; and the periphery of apartments along Brea Canyon Road, Appian Way, and Ferrara Court in the southwest. The western boundary is State Route 57. The Spadra cemetery with many introduced eucalyptus trees is not included. The ravines of Elephant Hill have fine examples of walnut woodland and the upper slopes have grassland, often dominated by mustards, with some areas of prickly pear scrub.

Southeast of Mission Boulevard, the SEA is bordered by Mission Boulevard. The north-facing slopes with excellent oak and walnut woodland are bordered by residential areas at the base of the steep slopes. The eastern end is bordered by North Ranch Road, and the southwest is bordered by the dense residential community of Phillips Ranch in the City of Pomona. The crest of the ridge is rolling grasslands that are dominated by introduced mustards, and the slopes have a mix of dispersed grasslands, chaparral, coastal sage scrub, and prickly pear scrub.

The SEA encompasses several different local jurisdictions, including the unincorporated area of the County, City of Covina, City of Glendora, City of La Verne, City of Pomona, City of San Dimas, City of Walnut, and City of West Covina.

Wildlife Movement

The SEA represents the only regional wildlife linkage between the San Gabriel Mountains and the Puente Hills and Chino Hills complex. Unlike the commonly held concept of a corridor, however, this SEA contains a series of discontinuous habitat blocks and patches, rather than an unbroken corridor for movement. This SEA facilitates movement and exchange between larger habitat areas by permitting terrestrial "island-hopping" between the SEA components.

Using birds as an example, movement may be initiated by an individual or group of birds in either the San Gabriel Mountains or the Puente Hills. Larger species, with the capacity to cover long distances, may make the passage as one segment of its journey. Smaller species, however, that lack the physical or behavioral capacity may not be able to attain this movement under normal circumstances. By utilizing various components of the SEA, the same species can cover this journey in several smaller trips. The same example may also apply to winged insects and wind-borne plant propagules and pollen. Interaction between, not just through the components, can occur as well.

This same function probably does not apply to other taxonomic groups. It is highly doubtful that amphibian, reptile and most mammal populations use this corridor as effectively as birds, if at all. Mule deer (*Odocoileus hemionus*), for example, do not occur within Bonelli Park, but are common in the San Gabriel Mountains and the Puente Hills. However, some mammals that tolerate urban environments, such as Virginia opossum (*Didelphis virginiana*), raccoon (*Procyon lotor*), and striped skunk (*Mephitis mephitis*), use the corridor in the manner previously described. Even mountain lion (*Puma concolor*) periodically enter Bonelli Park and Walnut Creek Park from the outside, by way of routes related to SEA components.

The manner in which the SEA allows wildlife populations in different areas to interact is less than ideal. However, exchange in the manner described above is dictated by the widespread urbanization of the region. "Island hopping" is the only remaining connection for regional interaction that can contribute to the maintenance of genetic variability and health of regional wildlife populations.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE EAST SAN GABRIEL VALLEY SEA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Met	This SEA contains various alliances of coastal sage scrub that are found in scattered patches over hillside habitat, which support a core population of the federally-threatened coastal California gnatcatcher. The gnatcatcher population has been slowly increasing, as recently observed in coastal sage scrub at two locations in the area, Bonelli Park and Buzzard Peak. There are an estimated 10 and 15 pairs of gnatcatchers in this population. The SEA also has a population of breeding coastal cactus wren. The rare multi-stemmed dudleya has its northernmost population in Bonelli Park.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Several plant communities within this SEA are CDFW highest priority communities due to their restricted distribution in the Southern California region. These communities include walnut woodlands, which are scattered throughout the SEA; oak riparian woodland, which is excellent within the Walnut Creek drainage; isolated stands of willow woodland along many of the drainages; freshwater marsh and open water in association with Puddingstone Reservoir; and coastal sage scrub in scattered patches over hillsides. Coastal sage scrub serves as the habitat for the coastal California gnatcatcher, which has been slowly increasing in Bonelli Park and Buzzard Peak. These areas also have a population of breeding cactus wren. The rare multi-stemmed dudleya has its northernmost population in Bonelli Park.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	All of the plant communities and habitats indicated as restricted in distribution on a regional basis, are also restricted in distribution within the County.
D) Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	Any relatively large body of water with pockets of natural lakeside vegetation along its shoreline potentially meets this criterion, particularly within the context of an arid to semi-arid environment that is characteristic of the County. Although subjected to boating activities and shoreline recreational use, Puddingstone Reservoir serves as an important habitat for migrating water fowl and water birds, which is evident in the high diversity of birds recorded at the Bonelli Park over the past several years. The natural areas of the east San Gabriel Valley serve as a migration and habitat connection between the

DRAFT

			San Gabriel Mountains and the Puente Hills.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	Bonelli Park contains a population of the rare many-stemmed dudleya, which is at the northern limit of its range. Most of the species is found in the Peninsular Ranges in Orange County. The coastal California gnatcatchers are in the northern extent of their range here, are clearly at an extreme in physical/geographical limitations, and may represent unusual variation in a population or community. The SEA, therefore, meets this criterion.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Not Met	The SEA does not contain areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.

In conclusion, the area is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; and E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.

Griffith Park SEA

Boundary and Resources Description

The Griffith Park SEA is located within Griffith Park, the central park of the City of Los Angeles, situated on the extreme eastern end of the Santa Monica Mountains. The SEA is an extensive, relatively undisturbed island of natural vegetation in an urbanized, metropolitan area. It supports the coastal sage scrub, chaparral, riparian, and southern oak woodland plant communities that are typical in the interior mountain ranges of Southern California. What makes the SEA important is its geographical location. It has become an island of natural vegetation surrounded by urban and suburban development. The geographic location makes the area important for scientific study, for genetic interchange between otherwise isolated populations, and for recreation of urban residents.

The SEA is located partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Burbank and Hollywood.

The SEA encompasses most of Griffith Park, south of the State Route 134, and west of Interstate 5. The SEA boundary generally follows the natural area near the Griffith Park boundaries in most cases. Isolated areas are important for preserving and documenting the geographical variability of vegetation and wildlife that formerly occurred throughout the region. They serve as reservoirs of native species that could be of scientific and economic value in the future. In addition, birds rely on these islands for areas to rest and feed along their north-south and east-west migration routes. In the case of Griffith Park, this function is made even greater than might be expected because it serves as a corridor for any gene flow and species movement that may take place between the Santa Monica and San Gabriel mountains via the Verdugo Mountains.

Beginning in the northwest corner, and proceeding eastward, the SEA follows the natural vegetation

DRAFT

~~on the mountain slopes at the junction with the flood plain of the former Dark Canyon and the Los Angeles River. This area of the SEA includes the recently acquired (2010) Cahuenga Peak, at 1820 feet, which is now the highest point of Griffith Park. Cahuenga Peak slopes have rocky outcrops, chaparral, and regenerating oak woodland and chaparral on the north-facing slopes. (This area was part of the 800 acres burned in the Griffith Park Fire of 2007.)~~

~~The Los Angeles River is channelized, but there is remnant oak riparian woodland in this area. Bordering the apartment complex on the east side of Barham Boulevard, there is a somewhat abrupt change in slope where the previous Dark Canyon Creek flowed (Barham Boulevard was evidently constructed in this Canyon.). The SEA includes the remnant riparian coast live oak woodland (*Quercus agrifolia*), which has many jurisdictional oak trees and in many places, the natural understory. Residents and staff at the apartments report frequent sightings of wildlife, particularly mule deer (*Odocoileus hemionus*) and coyotes (*Canis latrans*), in their parking lots, which line the Griffith Park side of the complex. On the slopes above, the chaparral of this west-facing slope grades upward into an extensive area of coastal sage scrub. The SEA includes these natural areas. From the natural areas on slopes above the junction of Barham Boulevard and Forest Lawn Boulevard, the SEA boundary continues eastward along the border of natural vegetation on the slopes above Forest Lawn Boulevard, including oak woodland in the ravines and mixed chaparral and grassland on the upper slopes. Occasionally, an ash (*Fraxinus velutina*) or Southern California black walnut (*Juglans californica*) are in these ravines, along with oak trees and other chaparral plants.~~

~~The boundary follows natural vegetation southward, away from the Los Angeles River, at the boundary of Forest Lawn Memorial Park (Forest Lawn). A slope and ridge top that have been cleared by Forest Lawn have been excluded from the SEA, but the chaparral on the east-facing side of the slope is included. From this ridge, the SEA roughly follows at the edge of the natural areas around the south side of the Forest Lawn and returns northward on the parcel line between the Forest Lawn and Griffith Park.~~

~~From the east side of Forest Lawn, the SEA boundary includes a chaparral-covered slope that is south of Travel Town and Zoo Drive. Cooper and Mathewson (2008) describe how coastal sage scrub occurs through a broad section of the northern part of Griffith Park, from end to end with patches of the sensitive valley needlegrass grassland. From the natural area near the Interstate 5 and State Route 134 interchange, the SEA boundary swings around westward, north of the Los Angeles Zoo, and forming a lobe on the chaparral-covered slopes. This area has ravines and a gradually sloping area near Travel Town, with riparian forest that includes sycamores (*Platanus racemosa*), oaks, willows (*Salix* spp.), and mulefat (*Baccharis salicifolia*), which are easily seen along Griffith Park Drive. Travel Town is not in the SEA, but its periphery of native riparian and chaparral is included. The north-facing upper slopes have chaparral, and the south-facing upper slopes have coastal sage scrub or grassland with chaparral plants here and there, especially elderberry (*Sambucus* spp.). Along Zoo Drive, ravines have typical chaparral of north-facing slopes. The SEA boundary continues past the Los Angeles Zoo along a road to a landfill area within Griffith Park, and goes around the landfill, forming a cherry-stem shaped area at the landfill road, and then continuing southeastward on the west side of Griffith Park Drive, excluding the Harding Municipal Golf Course. The Spring Canyon picnic area is excluded, as the understory of the sycamores and oaks is unnatural lawn, and the SEA boundary continues south along natural vegetation along Griffith Park Drive to the southern boundary of Griffith Park, near the Los Feliz offramp from the Interstate 5. A golf course practice area at the corner is excluded from the SEA.~~

~~From the southeast corner, the SEA boundary goes west along with the Griffith Park boundary at the edge of development to another golf course, which is excluded due to extensive modification of~~

DRAFT

theslopes. The Greek Theater in Vermont Canyon and Griffith Park Observatory on the slope beyond are included, as the modified vegetation for each covers less than 40 acres. The SEA boundary continues west and then north with the Griffith Park boundary at the edge of development. A small quarry is excluded. The undeveloped upper Brush Canyon in Griffith Park is included. Griffith Park and SEA have oak woodland along the drainages, transitioning uphill into chaparral and then grassland on the upper slopes. Within Griffith Park, north-facing sides of rocky outcrops often have a cliffside vegetation that is characterized by multiple kinds of lichens, mosses, liverworts and other non-vascular plants along with live forever (*Dudleya* spp.), and other flowering plants. The SEA boundary follows Griffith Park boundaries around the development in the Blackwood Canyon area. A ridge area in Griffith Park on the south side of Mulholland Drive overlook is excluded. The SEA boundary follows Griffith Park boundaries on the southern edge and then turns north after including the grassland and coastal sage scrub covered slopes that cover the open area between the two northern arms of the Hollywood Reservoir. On the west side of the SEA, the boundaries lap west outside of Griffith Park boundaries to include the oak woodland and chaparral of the lower elevations of Cahuenga Peak in the neighborhood of Dark Canyon (Barham Boulevard) and Cahuenga Pass.

Wildlife Movement

Griffith Park has become increasingly isolated from the rest of the Santa Monica Mountain Range, the Los Angeles River, the Los Angeles Basin, the San Fernando Valley, and the Verdugo Mountains (a little less than two miles to the east) because of the freeways, concrete river projects and urbanization that surround Griffith Park. Although some species have disappeared, including the ringtail (*Bassariscus astutus*), the gray fox is still seen.

River bed vegetation is quickly returning in the Los Angeles River as sand deposits on the hard channel bottom, and re-vegetation should be encouraged. Major bird and mammal populations exist on the re-vegetated portions of the Los Angeles River. Although some stretches of the Los Angeles River may not provide suitable primary corridors, it is important to reinstate Griffith Park's connection to the Los Angeles River for the future of wildlife and plant connectivity. In the management draft for Griffith Park wildlife (Cooper and Mathewson 2008), the authors outline some of the important connections to maintain or enhance: bridges and underpasses over and under State Route 101 and culverts that feed into the Los Angeles River Channel.

Griffith Park is viewed as an important connective island for the Santa Monica Mountains to the west of State Route 101 and the Verdugo Mountains and San Gabriel Mountains to the east. Wildlife may also use the natural areas and even concrete channels of the Los Angeles River to connect to the Tujunga Wash and Hansen Dam SEA and to the San Gabriel Mountains.

Regional Biological Value

The SEA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

Criteria Analysis of the Griffith Park SEA

Criterion		Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Not Met	No known core populations occur within this SEA.

DRAFT

B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Not Met	No known unique or rare plant or animal species occur within this SEA that would be regionally uncommon. No rare plant habitats occur in Griffith Park. Griffith Park has extensive wild areas that are little studied according to Cooper and Mathewson 2008. Such areas could be discovered.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Not Met	No known unique or rare plant or animal species occur within this SEA that would be particularly uncommon in the County. No rare plant habitats are known in Griffith Park. Griffith Park has extensive wild areas that are little studied according to Cooper and Mathewson 2008. Such areas could be discovered.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	Griffith Park is the easternmost extent of the Santa Monica Mountains, and a stepping stone to the Verdugo and San Gabriel mountains, which are only two miles distant. It is a very important natural area for animals and plants species that go between the Santa Monica and San Gabriel mountains. Because of its large acreage, Griffith Park maintains sizable populations of biological communities, even including top predators, such as bobcats and mountain lions. Griffith Park is teetering between becoming an island of natural habitat in a metropolis and maintaining connections to the rest of the Santa Monica Mountains to the west.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	Griffith Park is the easternmost extent of the Santa Monica Mountains, and a stepping stone to the Verdugo and San Gabriel Mountains, which are only two miles distant. It is a very important natural area for animals and plants species that go between the Santa Monica and San Gabriel mountains. Because of its large acreage, Griffith Park maintains sizable populations of biological communities, even including top predators, such as bobcats and mountain lions. Griffith Park is teetering between becoming an island of natural habitat in a metropolis and maintaining connections to the rest of the Santa Monica Mountains to the west.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Griffith Park has extensive areas of coastal chaparral and is an island of refuge for native animals in the Santa Monica Mountains. Its mosaic of habitats includes coastal sage scrub, riparian areas, and southern oak woodland. The mosaic of habitats is especially valuable to preserve. Griffith Park is in the City of Los Angeles and protected in this respect, but no management plan preserves its natural habitat in perpetuity.

In conclusion, the area is an SEA because it contains D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent

DRAFT

unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Harbor Lake Regional Park SEA

Boundary and Resources Description

The Harbor Lake Regional Park SEA is located [within Kenneth Malloy Harbor Regional Park] in the Harbor City community of the City of Los Angeles, which is approximately 15 miles south of downtown Los Angeles and just west of the Interstate-110. The Lake is named "Machado Lake." The SEA supports one of three remaining wetlands that once covered the southern and western areas of the Los Angeles Basin. The SEA supports significant areas of aquatic and terrestrial plant communities, which provide habitat to a variety of birds and wildlife.

The site is located in the Torrance Quadrangle of the United States Geological Survey (USGS) 7.5 Minute Map Series (USGS, 1964).

The SEA boundaries encompass the lake areas that contribute to the biological function of the Harbor Lake ecosystem. It is bordered to the north by the Pacific Coast Highway, but includes a small segment of drainage, the Wilmington Drain ("Bixby Slough") north of Pacific Coast Highway. The slough is a secluded marsh that supports wildlife and waterfowl common to marsh areas. Thirty-five species of native birds have been observed breeding in the Wilmington Drain and a large number of interesting vagrants are recorded from this area. Following the Wilmington Drain segment, the boundary travels east along the south side of Pacific Coast Highway and south along the east boundary of Harbor Regional Park to the Harbor Park Municipal Golf Course. The SEA goes east to a parking area for the golf course. The golf course is included in the SEA to emphasize the need to keep this area green. Wildlife from the adjacent woodlands use the golf course to transit between natural areas of the park in a less populated area. The eastern boundary of the SEA includes the greens and excludes parking areas around the periphery of the golf course. The SEA goes west along the boundary of the natural riparian woodland of the park and golf course and then southwest along the boundary of marsh habitats of the park and the golf course. At the south end of the golf course, the SEA boundary goes generally southeast along the periphery of park and developed environment, much of it along the outer southern boundary of Los Angeles Harbor College. The SEA stops at the college southern frontage paved area, then turns west along Anaheim Street with the park boundary, excludes a graded and cleared area of the park in the southwest corner, and then goes with the park boundary north along Vermont Avenue to the Pacific Coast Highway. At the Pacific Coast Highway, the SEA boundary goes with the park boundary to the Wilmington Drain. Machado Lake is chiefly bordered by native marsh vegetation of cattails, rushes, reeds, and mulefat with some invasive giant cane (*Arundo donax*) and upland native riparian forest, which primarily consists of willows. The exception to the natural vegetation is along Vermont Avenue where lawns are maintained between parking areas and the lake for public recreation and picnicking. The bird species list has over 330 species, and the area is used by migrant birds on the Pacific Flyway during the spring and fall migration periods. The SEA lies generally west of the Interstate-110.

The SEA encompasses regionally unique areas, including one of three remaining wetlands that once covered the South Bay area. The freshwater plants and animals found here are completely surrounded by residential and industrial facilities. This type of habitat has been filled, drained, and lost to development throughout most of the County. In some areas, man-made lakes and ponds have created small freshwater marshes along their edges, but this is minimal in comparison to the large expanses of freshwater marsh that were once found in the Los Angeles Basin.

DRAFT

Freshwater marsh habitat supports a great diversity of wildlife. Most of the bird species found here are dependent in some way on the surface moisture and vegetation, and would not be able to survive without it. It is also a habitat that supports several species of amphibians. Frogs and toads can be found here that are becoming extremely difficult to find throughout Southern California. The marsh is also an important area for migratory birds. Because Harbor Lake Regional Park and Madrona Marsh are the only habitat of this type in the southern portion of the County, they serve as small scale wildlife refuges. Waterfowl, shorebirds, marsh birds, and others can be found on the marsh in numbers during the spring and fall migration.

Wildlife Movement

The vernal pools situated within this SEA serve as isolated, high-quality habitat and linkage resource for migratory waterfowl. The vernal pools teem with arthropod and amphibian activity, and provide essential feeding grounds for long distance migrants, as well as for resident species of reptiles, birds and mammals. The SEA does not lie within any identified terrestrial movement routes for wildlife.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE HARBOR LAKE REGIONAL PARK SEA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Not Met	This SEA contains habitat that supports a population of vernal pool fairy shrimp.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	SEA supports a sizeable lake with a freshwater marsh along its northern, eastern and southern shores and the Bixby Slough, habitats that once covered the South Bay area. This type of habitat has been filled, drained, and lost to development throughout most of Southern California.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Harbor Lake Regional Park is one of three remaining wetlands with freshwater marsh in the County (The others are Madrona Marsh and recreated Ballona Freshwater Marsh, which are also in SEAs.). This type of habitat once covered much of the southern and western Los Angeles Basin area, and supports several species of amphibians including frogs and toads that are becoming rare throughout Southern California.
D) Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in County.	Met	Harbor Lake Regional Park and its marshes is an important area for migratory birds. The list of number of bird species seen has over 330 species. Freshwater marshes are important breeding areas for a number of birds and amphibian species.

DRAFT

E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The Harbor Lake freshwater marsh has attracted considerable attention from the academic and scientific communities, and the resources of the area are well documented and continue to be studied.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The freshwater marsh is a good example of the freshwater marshes that used to occur along the fault lines of the Los Angeles Basin.

In conclusion, the area is an SEA because it contains B-C) a sizeable lake with a freshwater marsh along its eastern shore and the Bixby Slough, habitats that once covered the South Bay area. This type of habitat has been filled, drained, and lost to development throughout most of Southern California and the County. Three of these habitats remain in the County; D) the Harbor Lake Regional Park is habitat that serves as concentrated breeding, feeding, resting, and migrating grounds and is limited in availability both regionally and in the County; E) the Harbor Lake freshwater marsh has attracted considerable attention from the academic and scientific communities, and the resources of the area are well documented and continue to be studied; and F) the freshwater marsh is a good example of the freshwater marshes that used to occur along the fault lines of the Los Angeles Basin.

Joshua Tree Woodlands SEA

Boundary and Resources Description

The Joshua Tree Woodlands SEA is located in the western portion of the Antelope Valley west and northwest of the Antelope Valley California Poppy Reserve State Natural Reserve in an unincorporated area of the County. This SEA encompasses many of the remaining old-growth stands of Joshua trees (*Yucca brevifolia*) on the west side of the Antelope Valley. Joshua tree woodland is a complex biological community of the gradual slopes of higher elevation desert area as that once covered much of this part of the Antelope Valley around the Antelope Wash. Joshua trees only occur within the Mojave Desert, and Los Angeles County populations are at the western limit of the species' range.

Because Joshua trees live in areas that are easily developed for residences and agriculture, this habitat has become very fragmented in the County. The SEA consists of eight separate units, seven of which are in close proximity to each other between the Kern Los Angeles County line to the north, and the California Aqueduct and Fairmont Butte to the south. The eighth unit is in an arroyo on the north side of the principal western ridgeline of Liebre Mountain, which is near the furthest western extent of Joshua tree woodland in Southern California. This woodland is located partially within the Angeles National Forest, and east and adjacent to the Interstate-5. The eighth unit is bordered on three sides by the San Andreas SEA.

All of the SEA except Unit 8 is within an area designated as the Antelope Valley Globally Important Bird Area (IBA) by Audubon California. This part of the Antelope Valley is very important as a resource area that supports spring and fall migration of birds, from the small passerines to the larger raptors, such as the state-threatened Swainson's hawk (*Buteo swainsoni*) and turkey vultures (*Cathartes aura*). The Joshua tree woodland is a very important resource to these migrations by supplying perches and food for these animals on their journeys. The SEA is near the San Andreas SEA, the Antelope Valley California Poppy Reserve State Natural Reserve, the Arthur B. Ripley Desert Woodland State Park, and the County George F. Bones Desert Pines Wildlife Reserve; however, many

DRAFT

of these areas are not contiguous with one another nor with the SEA. Unit 2 of the SEA includes much of the Arthur B. Ripley Desert Woodland State Park. Unit 8 of the SEA is contiguous with the San Andreas SEA.

Fragmentation is a concern because the Joshua trees depend on a small moth for reproduction. Only two species of moth can successfully pollinate Joshua trees, and in the SEA, there is only the yucca moth (*Togeticula synthetica*). The moth may have limited dispersal abilities, and the Joshua trees cannot reproduce from seeds without pollination from this particular moth. Cross pollination is regarded as essential to a species' genetic diversity, which is essential to adaptation to environmental change.

The Joshua trees in the seven units have the growth form of the lower elevation woodlands of the flatter areas, and somewhat spaced from one another and less clumped. The Joshua trees in the eighth unit have a growth form that is more common in the hilly areas, where the individuals sprout from connected rhizomes and are clumped. Many times, these clumps are clones, with individuals all sharing the same genetic identity.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Neenach School, Fairmont Butte, Black Mountain, and Lebec.

The SEA is composed of eight units. The overall boundaries are as follows: The western boundary for units 1-7 terminates at 220th Street West (the border between Ranges 15W and 16W). The eastern boundary is 145th Street West. The northern boundary is on Avenue A at the Kern-Los Angeles Countyline. The southern boundary straddles the California Aqueduct, touches the Los Angeles Aqueduct, and is approximately on Avenue F. The southernmost area is located close to the foothills of the western San Gabriel Mountains.

Unit 1: The northernmost unit is bounded by Avenue A on the Kern-Los Angeles County line on the north between 200th Street West and approximately on 218th Street West. It extends irregularly to the south along a desert wash contour, about a 0.7 mile at its greatest extent. The current southern boundary is determined by agricultural clearing. This unit has a Joshua tree woodland with many shrub components of the biological community intact, including a floor covered by the wildflower slender goldfields (*Lasthenia gracilis*) in the spring.

Unit 2: Another unit is located between Avenue C to the north and Avenue F to the south (straddling part of State Route 138 on Avenue D and part of Lancaster Road on Avenue E), and east to west from about 200th Street to about 220th Street West. Vegetation clearance in various parcel units accounts for this unit's irregular shape. Agricultural clearing on both sides of the Antelope Wash has separated this unit from Unit 1 to the north. The intervening area is a broad wash plain with rich alluvial soils. The former agricultural fields may now become fields of photovoltaic panels to generate renewable energy. This unit has a southern square mile that straddles the California Aqueduct and touches the Los Angeles Aqueduct at the base of the San Gabriel Mountains. In the northern area, this unit has old-growth Joshua tree woodlands on a rocky ridge that grades into stands of Joshua trees and woodland that includes California junipers (*Juniperus californica*) in flatter areas toward the south. The southern and eastern parts of this unit overlap with much of the Arthur B. Ripley Desert Woodland State Park. The California Aqueduct is open in this area and is an important resource for bird migration along the desert slopes of the western San Gabriel Mountains, particularly waterfowl. The Los Angeles Aqueduct is generally in concrete pipe for most of its extent, and in this area, is covered by a berm and road. A colony of burrowing owls (*Athene cunicularia*), which is a state species of special concern, was discovered during surveys for an adjacent photovoltaic panel development, and probably other colonies or individuals of the owl live within this unit.

DRAFT

Unit 3: Another unit is located between Avenue D to the north and Avenue E to the south, and between 190th Street and 195th Street West. It is on the broad outwash alluvial area of Kings Canyon and adjacent drainages. This outwash area is somewhat blocked by the aqueducts, but both aqueducts are provided with underpass channels for outflow of the canyons onto the desert floor. The SEA includes a central cleared area that is regenerating the Joshua tree woodland and a residence with less than 40 acres cleared. The area next to Avenue D that has been cleared of Joshua trees is not included.

Unit 4: The square mile between Avenue C and Avenue D, and between 180th Street and 190th Street West has a good stand of Joshua tree and juniper woodland. This is also in the Kings Canyon alluvial wash area. There is a known area of Joshua tree regeneration to the east that is not included in the SEA.

Unit 5: The quarter square mile between Avenue C-5 and Avenue E, and between 180th Street and 185th Street West, is also on the Kings Canyon alluvial wash area and has a good stand of Joshua tree and juniper woodland.

Units 6: An area of a little over one eighth square mile is located at the corners of both units 4 and 5. It is between Avenues D and E and between 180th Street and what would be 174th Street West. This is also in the Kings Canyon alluvial wash area and has a good stand of Joshua tree and juniper woodland.

Unit 7: A large irregular unit is located roughly between Avenue B, Avenue C5, 145th Street and 180th Street West. It has an extensive area of Joshua tree juniper woodland that grades into stands of Joshua trees towards the east. There is a known area of Joshua tree regeneration in former agricultural fields between 160th Street West and 170th Street West that is not included in the SEA. The alluvial wash in the SEA is a combined area of outflow from Kings Canyon, unnamed canyons, and Broad Canyon.

Unit 8: The eighth unit is in an arroyo on the north side of the principal western ridgeline of Liebre Mountain, which is near the furthest western extent of Joshua tree woodland in Southern California. This woodland is located partially within the Angeles National Forest. It is east and adjacent to the Interstate 5. The eighth unit is bordered on three sides by the San Andreas SEA. This woodland has the clonal growth that is typical of Joshua trees in hilly areas.

The SEA is located primarily on the western Antelope Valley floor between the Tehachapi Mountains and the western San Gabriel Mountains. The topography of the SEA is extremely flat with the land sloping less than 200 feet in approximately five miles. The location and orientation of the SEA represents a matrix of remnant stands of Joshua tree woodland among a patchwork of disturbed areas. Nearly all of the land within the SEA is undisturbed and vegetated. Most of the land surrounding the SEA is disturbed by agricultural use, and also has some scattered rural residences. The SEA is entirely within the unincorporated area of the County.

Wildlife Movement

Wildlife movement within the SEA is possibly limited to local movement, but large scale movement across the Antelope Valley floor is probably much facilitated by the Joshua tree habitat as island like stepping stones. Typically in burned over areas, animal paths tend to orient toward the Joshua tree habitat. Birds, and possibly bats, and other aerial organisms that use the habitat linkage along the desert side of the San Gabriel Mountains probably use the woodland in the SEA for resting and feeding. Animals foraging within the SEA are unlikely to occur in concentrated numbers due to the

DRAFT

heterogeneity of the topography and habitat of the SEA. However, local movement to and from the different SEA areas, as well as to and from the San Gabriel Mountains and the Tehachapi Mountains may be restricted due to the disturbed nature of the Antelope Valley floor. Wildlife movement is likely to converge in areas where movement is still possible, which produces concentrated movement areas or “bottlenecks.”

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE JOSHUA TREE WOODLANDS SEA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Not Met	Although there are several listed species that occur within the SEA, this criterion is not met due to the lack of known core population areas.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The SEA contains large patches of undisturbed Joshua tree woodland habitat, which has become increasingly rare in the region.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	As stated above, Joshua tree woodlands have become rare in the region, and are even more rare in the County.
D) Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The habitat within the SEA has been studied for how it may serve as a concentrated breeding, feeding, resting, or migrating ground for any species. Some cross-desert migratory routes depend, in part, on the cover and habitat of the Joshua tree woodland. The units 1-7 of the SEA on the Antelope Valley floor are in a globally IBA, known as a bird migration route. The Joshua tree woodland is an important component of resources that supports this migration.
E) Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	Due to the scarcity of Joshua tree woodland, specimens of the quality found in the SEA are important to science and have become living laboratories. The SEA contains the most westerly extent of this habitat type.
F) Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The Joshua tree woodland contained within the SEA is an excellent example of this community type.

In conclusion, the area is an SEA because it contains B-C) Joshua tree woodland, a rare community both regionally and within the County; D) habitat important to breeding, feeding, and migration; E) the

geographic limit of Joshua tree woodland; and F) an excellent undisturbed example of Joshua tree woodland.

Madrona Marsh Preserve SEA

Boundary and Resources Description

The Madrona Marsh Preserve SEA lies on the northeast corner of Madrona Avenue and Sepulveda Boulevard, within the City of Torrance. The SEA boundaries encompass the natural part of the Madrona Marsh Preserve, which is a remnant of one of the last natural vernal wetlands in the County. Vernal marshes fill in the rainy season (winter and spring in our area) and dry completely by the end of summer. They are heavily used by migrant birds in the spring, and in the fall if rains are early. They usually have other resident flora and fauna that are specially adapted to the seasonal cycle of wetting and gradual complete desiccation.

The SEA is located within the Torrance United States Geological Survey (USGS) 7.5' California Quadrangle.

The SEA boundaries generally follow the Madrona Marsh Preserve boundary, which is surrounded by urban development. The Madrona Marsh Preserve is a park of the City of Torrance that is situated to the west of Maple Avenue, north of West Sepulveda Boulevard, east of Madrona Avenue and directly south of Plaza Del Amo. The Madrona Marsh Nature Center has offices and visitor activities to the north of Plaza Del Amo. The Nature Center supports an active volunteer and educational program for natural areas in this part of the Los Angeles Basin.

The SEA supports one of three remaining freshwater wetlands that once covered much of the southern and western Los Angeles Basin area. In the lowland area are vernal marshes and an alkaline margin, and the upland supports a back dune system and vernal pools.

The freshwater plants and animals found here are completely surrounded by residential development and industrial facilities. This type of habitat has been filled, drained, and lost to development throughout most of the County. In some areas, man-made lakes and ponds have created small freshwater marshes along their edges, but this is minimal in comparison to the large expanses of freshwater marsh that were once found in the Los Angeles Basin.

Freshwater marsh habitat supports a great diversity of wildlife. Most of the bird species are dependent in some way on the surface moisture and vegetation, and would not be able to survive without it. It is also a habitat that supports several species of amphibians. Frogs and toads can be found here that are becoming extremely difficult to find throughout Southern California. The marsh is also an important area for migratory birds. Over 150 bird species have been recorded as using Madrona Marsh. Because the remnant freshwater marshes like Madrona Marsh are the only habitat of this type in southern portion of the County, they serve as miniature wildlife refuges. Kenneth Malloy Harbor Regional Park (Harbor Lake Regional Park SEA) and Ballona Freshwater Marsh (Ballona Wetlands CRA) are the two other freshwater marshes in this area. Waterfowl, shorebirds, marsh birds, and others can be found on the marsh in numbers during the spring and fall migration.

Wildlife Movement

The vernal pools situated within this SEA serve as an isolated, high-quality resource providing habitat linkage for migratory waterfowl. The vernal pools teem with arthropod and amphibian activity, and provide essential feeding grounds for long-distance migrants, as well as for resident species of reptiles,

DRAFT

birds and mammals. The ponds do not lie within any identified terrestrial movement routes for wildlife.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

Criteria Analysis of the Madrona Marsh Preserve SEA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Not Met	Not met, although the Madrona Marsh contains vernal pools, which may harbor threatened and endangered species in a unique location for the County.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Madrona Marsh is one of three remaining wetlands with freshwater marsh in the County (the others are Kenneth Malloy Harbor Regional Park and the Ballona Wetlands freshwater marsh.). This type of habitat once covered the South Bay area of the Los Angeles Basin, and supports several species of amphibians and vernal pools that are becoming rare throughout Southern California.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	Madrona Marsh is one of three remaining wetlands with freshwater marsh in the County. This type of habitat once covered the South Bay area, and supports several species of amphibians including frogs and toads that are becoming rare throughout Southern California. In addition it has some vernal pools, which are very rare in the County.
D) Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	Any natural wetland like the Madrona Marsh is an important area for amphibians, which need water for their aquatic larvae to complete their life cycle. The Marsh is very important for migratory birds as a waypoint during the spring and fall migration periods.
E) Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	Because the Madrona Marsh is one of only three remnants of the formerly extensive habitat, there is much interest in the local community and at academic and scientific institutions in preserving the area and acquiring adjacent habitat currently used for gas and oil extraction that can be restored to marsh habitat.
F) Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The Madrona Marsh is a remnant of the freshwater marsh once prevalent in this area of the County, and it is important to recognize protection of the habitat.

In conclusion, the area is an SEA because it contains B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in

DRAFT

distribution in the County and regionally; D) concentrated breeding, feeding, resting, and migrating grounds, which are limited in availability in the County; E) unique habitat that is of scientific and educational value; and F) a very rare habitat of vernal marsh that has retained its unique yearly cycle.

Palos Verdes Peninsula and Coastline SEA

Boundary and Resources Description

The Palos Verdes Peninsula and Coastline SEA is chiefly located offshore around the Palos Verdes Peninsula, but also includes terrestrial habitat in Portuguese Bend and several other disjunct drainages and ridgelines. All of the many offshore rocks within 12 nautical miles of the coast are part of the California Coastal National Monument that managed by the Bureau of Land Management in the U.S. Department of the Interior. Many endangered marine mammals use this habitat. The SEA includes the inshore part of the Point Vicente State Marine Conservation Area (SMCA) and the Abalone Cove SMCA. Most of the interior Peninsula canyons and slopes are critical habitat for the federally-threatened coastal California gnatcatcher (*Poliophtila californica californica*). Many of the interior areas are also critical habitat for the federally endangered Palos Verde blue butterfly (*Glaucopsyche lygdamus palosverdesensis*). The gnatcatcher favors the coastal sage scrub or chaparral vegetation; the butterfly favors flatter areas with grassland plants of the upper marine terraces, such as Ventura milk-vetch (*Astragalus pycnostachyus* var. *lanosissimus*), which is a principal food plant for its larvae. The gnatcatcher critical habitat covers virtually all of the terrestrial part of the SEA and extends well beyond the SEA on the Peninsula. The butterfly's critical habitat is chiefly in flatter areas of the marine terraces, but can include broader areas of the canyons.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Torrance, San Pedro, and Redondo Beach

Most of the SEA is in the nearshore marine habitat and extends from the high tide line of the beach to about one mile offshore. The designated SEA traverses along the coastline of the Palos Verdes Peninsula from Flat Rock Point at the northern end of Bluff Cove to the west end of the breakwater of Los Angeles Outer Harbor. The terrestrial part of the SEA includes several non contiguous ravines, canyons and streambeds that are adjacent to rural residential areas. The area known as Portuguese Bend is one of the largest inland areas within the SEA.

The SEA has the inshore part of the Point Vicente SMCA, which extends from mean high tide due east west to three nautical miles offshore. The Point Vicente SMCA is like a reserve in that no take of any kind is permitted. The western line goes due east-west off an unnamed point about a 0.4 mile north of Point Vicente. The eastern line goes due north-south off Long Point. Adjacent is the Abalone Cove SMCA, which extends from mean high tide to 3 nautical miles offshore, also due north-south. The eastern boundary of the Abalone Cove SMCA extends off Portuguese Point.

Disjunct interior parts of the SEA include 1) Agua Amarga Canyon, which drains to Lunada Bay (critical habitat for both the Palos Verdes blue butterfly and the coastal California gnatcatcher; 2) an area of coastal sage scrub and bluffs north of Point Vicente; 3) the upper bluff area of chaparral and grassland that is on the northeast side of Pacific Coast Highway at Point Vicente; 4) the beach and cliff area south of Pacific Coast Highway at Abalone Cove, Portuguese Point, Inspiration Point and Portuguese Bend; 5) the unbuildable areas of the Portuguese Bend Landslide, which include grassland, chaparral, and canyons and surround the community of Portuguese Bend on the north side of Pacific Coast Highway, and border the southern side of the community of Rolling Hills; 6) cliffs and headland areas with coastal bluff, coastal chaparral, and grassland extending from the east side of Portuguese Bend

DRAFT

~~to the City of Los Angeles border at the west end of Royal Palms Beach Park; 7) a lobe bordering the switchbacks of Palos Verdes Drive East that connects to the coast and covers the natural cliff, bluff, and chaparral vegetation up to about one mile inland; and 8) the intertidal area on the west side of Point Fermin.~~

~~The other disjunct interior parts of the SEA are within the community of Rolling Hills, with numerous natural parts of the canyons and ridges have been too steep to develop. These are designated as part of the SEA because the native vegetation is essential for the native birds and the migrants, which include those that use the Palos Verdes Peninsula as a rest stop and those that overwinter. They include the following canyons, which are on the north side of the Palos Verdes Range crest, and drain to the Los Angeles Basin: 9) Agua Negra Canyon on the west side of Crenshaw Avenue; 10) Agua Magna Canyon; 11) Sepulveda Canyon; 12) Blackwater Canyon, including two upland forks; and 13) George F. Canyon with all three upland forks.~~

~~Finally, the remaining disjunct interior parts of the SEA are in the Miraleste district of Rancho Palos Verdes, on the east side of Palos Verdes Drive East, naturally vegetated canyons trend eastward to the flat area of San Pedro: 14) north of Rockinghorse Road; 15) north of Colt Road; 16) north of Coral Ridge Road; 17) Miraleste Canyon (north of Miraleste Road); and 18) two branches of San Pedro Canyon (south of Via Siena and north of Via Colinta).~~

~~The SEA is formed by the Palos Verdes Range, and is characterized by marine terraces atop rough terrain that is created by the erosion of the upland areas into many canyons. The flatter areas on the marine terraces are the areas that have been developed. Unparalleled headlands, rocky shoreline, and the land-sea interface provide for a tremendous variety of biotic resources in this area. It is one of the most biologically diverse and productive regions in the County, and contains several biotic communities, including rocky intertidal, kelp bed, coastal strand, and coastal sage scrub. One small sandy beach is periodically present on an ephemeral basis at Portuguese Bend. This 10 mile stretch of coastline, between Point Fermin and Bluff Cove, is the only sizeable rocky intertidal area in the County.~~

~~The Peninsula was once an island before sediments filled the Los Angeles Basin, and it has many biological associations today that originated during the time it was insular. A number of the uncommon plants, for example, are also only known from offshore islands, and some of the bird subspecies on the Peninsula are also only known elsewhere on the offshore islands. They are not found on the mainland except on the Palos Verdes Peninsula.~~

~~Rocky shores support a great number of species. This is primarily due to the interface of the ocean and the land, the highly diverse natural communities that both contain, and the oxygen and food-rich environment offered by this habitat. The natural features include a variety of substrates, both stable and unstable, the aeration of water and humidification of the air through wave splash, and the upwelling of nutrient rich waters along the Southern California coast.~~

~~Subtidally, one can find representatives of every phylum of animals (major groupings, such as mollusks, echinoderms, jellyfish, etc.), many of which are not represented on land. Rocky substrates are as diverse as the intertidal areas. Sandy subtidal areas also support a great variety of species, but most are cryptic and submerge into the depths of the sand because of the high energy and unstable nature of their environment.~~

~~Kelp beds dominated by giant kelp (*Macrocystis pyrifera*) are an extremely productive habitat that provides food, attachment sites, and shelter for a diverse community of invertebrates, fishes, and sea mammals. They are an important nursery habitat for juvenile fishes. Kelp beds were originally common off the~~

DRAFT

Southern California coast wherever rocks were present at shallow depths. However, due to man-made and natural phenomena, this habitat has been severely diminished in the region, and is now rare in the County. A kelp bed habitat restoration program was begun in the area in the 1960s, and appears to have been successful. Kelp has been reestablished west of Point Fermin, along Abalone Cove, and offshore of Halfway Point. Smaller colonies are now reestablishing at other locations offshore of the southern coast of the Peninsula.

The coastal cliffs found in the area range in elevation from 100 to 300 feet and support coastal sage scrub and coastal strand bluff scrub. These and offshore rocks offer ideal roosting and feeding sites for numerous shorebirds, gulls, and other seabirds, including the fully protected California brown pelican (*Pelecanus occidentalis californicus*). The area is an important stop for migrating birds as they fly along the coast or across the Santa Monica Bay. In addition, the bluff tops that are now abandoned agricultural fields are utilized by many species as wintering feeding grounds. One fully protected species, the American peregrine falcon (*Falco peregrinus anatum*), and one very uncommon species of conservation concern, the prairie falcon (*F. mexicanus*), have been known to winter here in recent years.

The bluff tops and cliffs have been disturbed by pedestrian use, residential development, and agriculture. Only very small, isolated ravines remain in a natural state. The shoreline has suffered major biological impairment, commonly blamed on over-collection by humans and intense pollution. The health of the marine environment has been relatively poor due to human influences, such as outfalls with toxic contents, but appears to be slowly recovering.

Wildlife Movement

The SEA is on the Pacific Flyway migration route, which is used by many birds seasonally and plays an important role as a stopover because of its extensive undisturbed marine resources. It does not fall within any identified terrestrial movement routes for wildlife because it is isolated by the ocean on one side, and enclosed by miles of developed land on the remaining terrestrial sides.

The offshore major part of the SEA is on the annual migration route of the California gray whale (*Eschrichtius robustus*), which is a Convention on International Trade in Endangered Species (CITES) Appendix I animal. This means that they have endangered status, and no trade or harvest is permitted, except by strictly controlled export and import permits that are issued in countries that are party to the convention and have legislation of adoption, which includes the U.S. The gray whale migrates close to shore on its transit during the winter months, from the Bering Sea to lagoons of Baja California for calving and breeding. A whale count station is manned by volunteers at the Point Vicente Park and Interpretive Center that has yearly records, starting in 1979. The whales generally return north offshore. The area may be a migration corridor for other marine animals, as this is part of the great Southern California Bight, which has an eddy circling counterclockwise to the north off the southward flow of the California Current. This flow brings marine organisms (fishes, invertebrates, and plants) that may have major population concentrations in the south to this part of the ocean habitat. Most of these organisms have a planktonic stage that is dependent on the current system to maintain their populations and distribution. Pollution of the ocean waters by development on land will have significant effects that may not be appreciated immediately because of the cryptic nature of subtidal animals and plants.

Other Cites Appendix I mammals have been recorded as using the SEA in the Point Vicente records including sperm whales (*Physeter macrocephalus*), minke whales (*Balaenoptera acutorostrata*), humpback whales (*Megaptera novaeangliae*), blue whales (*Balaenoptera musculus*), fin whales (*Balaenoptera physalus*), and southern sea otters (*Enhydra lutris nereis*).

Regional Biological Value

The SEA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE PALOS VERDES PENINSULA AND COASTLINE SEA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Met	The Palos Verdes Peninsula is the western-most area of breeding for the threatened coastal California gnatcatcher. The headlands and rocky shoreline provide wintering habitat for two important birds: the fully protected California brown pelican and the state-fully protected American peregrine falcon. The canyons of the SEA support three races of birds that are residents only on the Peninsula and the Channel Islands: insular forms of the orange-crowned warbler, western flycatcher, and Allen's hummingbird. Rare plants, such as southern island mallow and Island cressosoma, make this area a very special mainland population for species that otherwise occur only on the offshore islands. Island green dudleya, a rare bluff plant known only from Palos Verdes Peninsula, Santa Catalina Island, and San Nicolas Island, occurs in the Portuguese Bend Landslide.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Unparalleled headlands, rocky shoreline, and the land-sea interface provide for a tremendous variety of biotic resources in the coastline area. Rocky shores support a great number of species, and the upwelling of deep waters provides nutrient rich waters for the area's kelp beds, which are dominated by giant kelp. Kelp beds were formerly common off the Southern California coast wherever rocks were present at shallow depths. However, due to man-made and natural phenomena, this habitat has been severely diminished in the region. The mainland SEA areas contain mainland populations of species that occur chiefly on the offshore Channel Islands. Any population characteristic of an island is of extremely restricted regional distribution.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	Kelp beds are now rare in the County. These algal communities can account for 90% of the biomass where they occur, and provide food and habitat for hundreds of species. Like terrestrial forests, they moderate the microclimate, reduce wave shock to shorelines and provide shade and shelter to their myriad denizens. The coastal cliffs support coastal sage and coastal strand vegetation, which are rare in Los Angeles because of the scarcity of rocky headlands. The Peninsula's former island status and current maintenance of mainland populations of species that occur chiefly on the offshore Channel Islands are very special. Any population characteristic

DRAFT

			of an island is of extremely restricted distribution in the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The coastal cliffs provide ideal roosting and feeding sites for numerous shorebirds, gulls, and other seabirds, including the fully protected California brown pelican. This headland is a principal rest stop for migrating terrestrial and marine birds on the Pacific Flyway and a wintering area for some migrants. Bluff tops and marine terraces of abandoned agricultural fields are utilized by many species as winter feeding grounds. The Palos Verde Peninsula is the western-most area of breeding for the federally threatened coastal California gnatcatcher. It has numerous plant and animal species that only occur here and otherwise on the offshore islands, so this is a special area for all phases of their life cycles. The marine area is an important migration area for all of the sealife and marine mammals.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	Much scientific and educational work has been done on this part of the coastline. State and county agencies have generated a great deal of information about the area. The Peninsula was once an island and has biotic characteristics more like the offshore Channel Islands than the rest of the County. It is an anomaly of an island now attached to the mainland, and thus interesting for biogeographic studies. The Peninsula canyons are the western-most area of breeding coastal California gnatcatchers. Because of the resident birds and plants that occur only on the Palos Verdes Peninsula and the Channel Islands, this area is of scientific interest for study of island biogeography and evolutionary ecology.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	As the only extensive (10 miles long) rocky intertidal shoreline in the County, the coastline is an important area for preserving its intense biodiversity. It has headlands, rocky shoreline cliffs, rocky intertidal areas, boulder field intertidal areas, kelp beds, coastal strand, ephemeral coastal strand, and coastal sage scrub. The bluffs have a special array of plants found nowhere else on the County mainland. The Peninsula has diverse communities typical of the offshore Channel Islands including coastal sage scrub (which is used by the mainland threatened bird, coastal California gnatcatcher), chaparral, and riparian habitats. The SEA canyons are the least disturbed coastal drainages of the County.

In conclusion, the area is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, and migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either

an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Puente Hills SEA

Boundary and Resources Description

The Puente Hills SEA is located in the Puente Hills in the southeastern portion of the County. The Puente Hills are an inland topographical feature that separates the San Gabriel Valley to the north and the coastal plain to the south. The hills are oriented east-west and stretch from the San Gabriel River on the west approximately to the San Bernardino-Los Angeles County line to the east, where they transition into the Chino Hills. The SEA includes portions of the Whittier Narrows Dam Recreation Area and Flood Control Basin, and much of the undeveloped land throughout the Puente Hills. Nearly the entire SEA is designated as the Puente-Chino Hills State Important Bird Area (IBA) by Audubon California. The main area hosts migrating and resident birds that use the extensive mosaic of lowland terrestrial habitats, and notable extensive areas of grassland and oak and walnut woodlands. This IBA extends well beyond the SEA into Orange and San Bernardino counties, and in general, goes beyond the SEA boundaries in most places. The northwestern disjunct area of the SEA is part of the Los Angeles Flood Control Basin IBA, which hosts many resident and migrating birds that use the wetlands. This IBA extends beyond the SEA on both the Rio Hondo and a long distance upstream along the San Gabriel River.

The SEA encompasses portions of the El Monte, Baldwin Park, Whittier, La Habra and Yorba Linda United States Geological Survey (USGS) 7.5' California Quadrangles. The majority of this SEA lies within unincorporated area of the County.

In the westernmost part of the SEA, boundaries encompass the undeveloped portions of the Montebello Hills. The oil field and transmission lines that occupy this area have protectively fenced and left most of the coastal sage scrub, which is inhabited by the largest population of the federally-threatened coastal California gnatcatcher (*Poliptila californica californica*) in the County. This area is part of critical habitat for the gnatcatcher. Boundaries of critical habitat extend to Montebello Boulevard on the north side and to the west side of San Gabriel Boulevard, while the SEA boundary stops at the transmission lines' southern edge. Some of the canyons in the oil field have oak woodland.

The SEA has a finger that extends from the Montebello Hills section over San Gabriel Boulevard to the oak woodland (among oil field structures) that borders the Rio Hondo Channel. The SEA finger continues upstream along the natural riparian course of the Rio Hondo to a point where the Rio Hondo is encased as a concrete flood control channel. This area of the Rio Hondo usually has surface water and is on the migration route for migrating and wintering waterfowl. The riparian area of the Rio Hondo immediately north of San Gabriel Boulevard has long been recognized as important to birds and has been called "Bosque Del Rio Hondo" since Spanish occupation of this area. Abundance of birdlife is frequently mentioned, but the area also has amphibians, reptiles, mammals, and probably fishes.

It is intended that the SEA encompass only natural areas of the basin and portions of the San Gabriel River and Rio Hondo, and the SEA in this area is disjunct from the rest of the Puente Hills. A golf course that is part of the Whittier Narrows Dam County Recreation Area is on the west side of Rio Hondo, and the golf course is not included in the SEA. Likewise, the manicured areas of the County Recreation Area on the east side of Rio Hondo are not included.

Moving east, the southern boundary of the SEA follows the upstream edge of the Whittier Narrows

DRAFT

~~Dam, while the northern boundary goes east along the south side of Durfee Avenue from the east bank of the Rio Hondo. This captures the Flood Control Basin at the confluence of the San Gabriel River and Rio Hondo. The pool at the base of the Whittier Narrows Dam, which is regularly utilized by migrant and resident waterfowl, is included in the SEA. In the Flood Control Basin the connective channel between the Rio Hondo and the San Gabriel River has native riparian vegetation and usually has water. The connective channel is included in both the SEA and the critical habitat for the coastal California gnatcatcher. The northern SEA boundary follows the south side of Durfee Avenue eastward to near the junction with Peck Road. Here the northern SEA boundary follows the northern side of the grounds of the Whittier Narrows Nature Center to the San Gabriel River, which is naturally vegetated with a soft bottom. The southern SEA boundary has outlined the natural vegetation of the flood control basin and connective channel, with a finger extending south downstream along natural riparian vegetation in the San Gabriel River, to the crossing of San Gabriel Boulevard. Across from the Whittier Narrows Nature Center, the southern boundary is along the southeast side of the San Gabriel River. A finger of the SEA extends upstream along the area of the San Gabriel River, which has soft bottom and native riparian vegetation upstream about one mile (three quarters of a mile beyond the crossing of State Route 60, which is a short distance upstream from the Whittier Narrows Nature Center area). This includes the confluence with San Jose Creek that drains the south side of the San Jose Hills, and the north side of the Puente Hills. Currently, the Whittier Narrows area of the SEA outlined above is not physically connected to the remainder of the SEA due to urban development on both sides of Interstate 605.~~

~~Much of San Jose Creek is channelized, and the City of Industry is located in the flood plain. Because of this, most wildlife movement cannot use the main watercourse route, which makes Puente Hills the wildlife corridor through the area. A plan for the Whittier Narrows area, sometimes called "the Emerald Necklace," proposes to eventually have a string of parks along the San Gabriel River and Rio Hondo, stretching from the Whittier Narrows Dam County Recreation Area along the rivers to the Santa Fe Dam area, into the City of Irwindale. This would better connect wildlife from the Puente Hills, Chino Hills, Santa Ana Mountains and the other Peninsular Ranges of Southern California with the San Gabriel Mountains, and the Transverse Ranges across the northern side of the San Gabriel Valley.~~

~~Moving east across Interstate 605, the SEA begins at the end of Sycamore Canyon Road and the mouth of Sycamore Canyon (off Workman Mill Road on the south side of the west section of Rose Hills Memorial Park). The northern boundary essentially follows the edge of developed portions of the Rose Hills Memorial Park, the Puente Hills Landfill, and rural residential and suburban developments of the Hacienda Heights area, eastward to Peter F. Schabarum Regional Park. The southern boundary in this area tracks the edge of urban development along the southern slopes of the Puente Hills bordering the City of Whittier and the City of La Habra Heights. The SEA in this southern slope area includes some areas impacted by oil production that are otherwise connective by virtue of largely natural habitat of chaparral and grasslands. Much of this area of the SEA is critical habitat for the coastal California gnatcatcher. The critical habitat of the gnatcatcher extends into areas with appropriate habitat for the gnatcatcher among some development, for example, Rose Hills Memorial Park and the Puente Hills Landfill. Active operation areas within these properties were not included in the SEA because they did not meet the SEA mapping criteria.~~

~~Much of the summit area in the Puente Hills is conserved and under restoration by the Puente Hills Habitat Preservation Authority (PHHPA), which has sponsored important studies on habitat and wildlife movement, restoration, and the wildlife movement tunnel under Harbor Boulevard at the eastern side of the SEA. The SEA in this western end of the Puente Hills includes important natural drainages on the west side of the Puente Hills, which drain to the San Gabriel River: much of Sycamore Canyon, with walnut woodland and riparian scrub; and Turnbull Canyon with a mixed riparian forest~~

DRAFT

~~containing ash, sycamore, and oak. The ridges around Turnbull Canyon are mixed grassland and chaparral, with coastal sage scrub and a scattered population of the federally-threatened coastal California gnatcatcher. Mixed chaparral and cactus scrub on the Turnbull Canyon and Sycamore Canyon slopes support a local population of the sensitive coastal cactus wren (*Campylorhynchus brunneicapillus sandiegensis*). Other areas of the southwest slope drainages include Worsham Creek, La Cañada Verde Creek, Arroyo Pescadero, and San Miguel Creek. Underpasses of Colima Road, which could otherwise block the east-west wildlife movement along the Puente Hills, are used by wildlife as shown by studies sponsored by the PHHPA. (SWCA Environmental Consultants, 2012)~~

~~A lobe of the SEA goes over the ridge to natural habitat along north-facing slopes. These tributaries of Joe Hill Wash have a mix of oak and walnut woodland habitats along multiple drainages that end in San Jose Creek. Other areas of north slope drainage to San Jose Creek are included in the SEA on the southeast side of Workman Mill and along the transmission line corridor in the Workman Mill area. The north slope boundary skirts the developed areas of the community of Hacienda Heights. An area of planned development in the oak woodland just west of Peter F. Schabarum Regional Park has been excluded from the SEA.~~

~~Continuing east, the northern SEA boundary crosses the Schabarum Park accessory corridor and includes the undeveloped portions, as well as those of neighboring Trailview County Park in the SEA. Here the northern boundary is trending southeast, skirting along the edge of residential development within unincorporated Rowland Heights. The southern SEA boundary in this area skirts development in the community of City of La Habra Heights and includes the southern slope of Powder Canyon, which has a variety of natural habitats including riparian, walnut, and north-slope oak woodlands, California walnut woodland, rocky hillsides, chaparral, and some grassland. Powder Canyon is part of the watershed of San Jose Creek. Here the SEA crosses Fullerton Road in a broad swath about a 0.4-mile wide that includes the Powder Canyon undercrossing and Pathfinder Park on the east side. Pathfinder Park has riparian and chaparral habitat. There is an area around the ridgeline development along Vantage Pointe Drive that has been excluded from the SEA.~~

~~The northern SEA boundary continues along the edge of development at the southern edge of the communities of Rowland Heights to include walnut woodland, oak woodland, grassland, and chaparral vegetation on the north-facing slopes and central ridge of the Puente Hills. This area includes tributaries in the Puente Hills for the Brea Canyon watershed. The boundary crosses Brea Canyon (Santa Ana River watershed) and State Route 57 at the edge of development in the City of Diamond Bar, and continues along the boundary of natural vegetation to the San Bernardino-Los Angeles County line. East of State Route 57, this urban-wildland interface is more or less the boundary between development in the eastern San Gabriel Valley and the natural areas of the Firestone-Boy Scout Reservation, with its focal area on Tonner Canyon (also in the Santa Ana River watershed).~~

~~From the crossing of Fullerton Road, the southern SEA boundary encompasses the naturally-vegetated central ridgeline of the Puente Hills, and extends south along the west side of Harbor Boulevard to include natural area around a wildlife passage tunnel under Harbor Boulevard. This tunnel was constructed by the Puente Hills Habitat Preservation Authority, and its mud floor has many track prints that attest to frequent use by deer and other animals. The tunnel has the ridgeline area on its west side (with scattered residences), and on the east side are a deep canyon with fine riparian oak woodland, hills with scattered oil wells, walnut woodland, and grasslands. The southern boundary includes the canyon, but excludes the hills and grasslands of the oil field that are on the ridgeline and south of the canyon. On the northern slope of the included canyon is the Vantage Pointe development, which is excluded from the SEA. On the east side of the oil field, the southern SEA boundary trends southeast along natural-vegetation boundaries to the Orange-Los Angeles County line. At the Orange-~~

DRAFT

Los Angeles County line, the SEA borders the development in the City of Brea in Orange County for a short distance, and continues through grassland and chaparral to cross the State Route 57 into walnut woodlands, southern oak woodland, chaparral, coastal sage scrub, and riparian woodlands of Tonner Canyon on the Firestone Boy Scout Reservation. Tonner Canyon is in the Santa Ana River watershed, and the ridge separating Brea and Tonner canyons is generally considered the divide between the Puente Hills and the Chino Hills. From here the southern SEA boundary continues eastward on the Orange-Los Angeles County line to the three-way junction with the San Bernardino-Los Angeles County line. For its eastern border in the Chino Hills, the SEA boundary turns north and follows along the San Bernardino-Los Angeles County line to contact with the northern boundary. In Orange County, south of the Orange-Los Angeles County line, is the Chino Hills State Park, with grasslands, chaparral, and riparian oak woodlands that extend across into the Firestone Boy Scout Reservation.

Critical habitat for the coastal California gnatcatcher is generally coincident with the SEA in the western part of the SEA, but has lobes and fingers that extend into development areas where the preferred natural habitat of the gnatcatcher, coastal sage scrub, coexists with less dense residential areas. East of Fullerton Road, the gnatcatcher critical habitat differs from the SEA. Gnatcatcher critical habitat narrowly includes the Harbor Boulevard wildlife passage tunnel, goes on the south side of the Vantage Point exclusion area, and trends into Orange County on the eastern side of the oilfield that borders the Vantage Point development. There is a small area of gnatcatcher critical habitat in the Firestone Boy Scout Reservation in the County that connects to the large area of critical habitat in Orange County and the Chino Hills.

The majority of the SEA lies within unincorporated County jurisdiction. Other local jurisdictions have also been included within the SEA in order to delineate the boundaries of functioning habitat units. These include the City of Diamond Bar, the City of Industry, the City of La Habra Heights, the City of Montebello, the City of Pico Rivera, the City of South El Monte, and the City of Whittier.

Wildlife Movement

Evidence of significant wildlife movement throughout the Puente Hills SEA has recently been documented in a two year carnivore study commissioned by the Santa Monica Mountains Conservancy as part of a multi-jurisdictional effort to establish a region wide wildlife movement linkage. Additional studies on wildlife movement through the area include several focusing on the Harbor Boulevard and Colima Road Underpasses. These studies can be accessed at the following link: <http://www.habitatauthority.org/studies>. This SEA represents the County portion of a continuous series of natural open space within the Puente Hills and Chino Hills. Overall, this open space extends north and west from State Route 91 (SR 91) in Orange and Riverside Counties to the Whittier Narrows reach of the San Gabriel River. The open space physically links the Puente/Chino Hills with the Santa Ana Mountains and the San Gabriel Mountains, respectively. By virtue of these linkages and a complex of interconnected habitat units throughout the hills, the Puente/Chino Hills function as both an important wildlife linkage and resident habitat area for regional wildlife populations.

Within the SEA itself several habitat units, well defined by major canyons, exist. These include Sycamore Canyon, Turnbull Canyon, Powder Canyon, Brea Canyon and Tonner Canyon. Each of these, in and of themselves, is capable of supporting a diversity and abundance of wildlife. More importantly, however, these habitat units are connected by a series of open space corridors, which allows population exchange to occur. Maintenance of biological diversity and population viability is accorded throughout the SEA and the chance of local species extinctions due to isolation is minimized. This function is acutely important for wide-ranging species that meet their breeding and/or habitat

requirements over broad areas.

Although several major arterial roads and highways cross the hills, continued use of undercrossings and surface crossings by wildlife has been documented. This movement is largely east-west trending between large habitat blocks located in the western, central and eastern portions of the SEA. Species documented as moving through the area include bobcat, coyote (*Canis latrans*), gray fox (*Urocyon cinereoargenteus*), and mule deer (*Odocoileus hemionus*).

Regional Biological Value

The SEA meets several SEA designation criteria that consider regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE PUENTE HILLS SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	The population of the California gnatcatcher at the Montebello Hills is the largest in Los Angeles County. Pairs occur throughout the County portion of the Puente Hills, especially in Sycamore Canyon and Arroyo San Miguel. The coastal cactus wren has significant populations in the Puente Hills, occurring in the Montebello Hills, Sycamore Canyon, Rose Hills, Hellman Park in Whittier, and through Hacienda Heights into Rowland Heights. Several CNPS Rare plants occur in the Puente Hills, including Weed's mariposa lily.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Several plant communities within this SEA are CDFW highest priority communities due to their restricted distribution in the Southern California region. These communities include walnut woodland, which is scattered throughout this SEA; oak riparian woodland, which is best developed in the major drainages of Sycamore Canyon, Turnbull Canyon, Powder Canyon, Brea Canyon, and Tonner Canyon, but is found elsewhere; stands of southern willow scrub along many of the drainages; scattered freshwater marsh; and coastal sage scrub, which is found in scattered patches over hillsides throughout.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	All of the plant communities and habitats mentioned above as being restricted in distribution on a regional basis are also restricted in distribution within the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	This SEA represents the only large complex of multiple, relatively undisturbed habitats in southeastern portion of the County. It is regionally important to many resident species, as well as migrating species, which would otherwise not be able to meet their habitat requirements. In particular, large mammal and overwintering birds of prey and songbirds make use of this area. The Puente Hills are a well known migration corridor for migratory songbirds during spring migration (April and May). On foggy May mornings, the hilltops can

	Criterion	Status	Justification
			support hundreds of individual migrant songbirds, which forage actively in all available habitats.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The Puente Hills represent the clear northern edge of the Peninsular Ranges. For this reason, taxa such as red diamond rattlesnake, occur here at the edge of their range, where they co-occur with the more widespread Pacific rattlesnake. Several bird species extend west through the Puente Hills into the Los Angeles Basin and the Whittier Hills (Oak Titmouse, Grasshopper Sparrow). These species are absent from the floor of the Los Angeles Basin. The Whittier Narrows Nature Center provides the public with extensive information and opportunities for field study of the Whittier Narrows natural environment.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Both the oak woodlands and walnut woodlands within this SEA represent excellent relatively undisturbed examples. The walnut woodlands in this area are reported to be the best remaining stands south of Ventura County.

In conclusion, the area is an SEA because it contains A) habitat of core populations of endangered or threatened plant species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) populations of scientific interest because they are at range extremes and intermix with species from other areas with known examples of reptiles and birds; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Rio Hondo College Wildlife Sanctuary SEA

Boundary and Resources Description

The Rio Hondo College Wildlife Sanctuary SEA is located on the western edge of the Puente Hills near the San Gabriel River, within the City of Whittier, and south of the Interstate 605. The SEA is the designated a Wildlife Sanctuary of the Rio Hondo College in the northern and eastern part of the campus, and is currently used as a study area by the students and faculty of Rio Hondo College. The SEA includes natural areas bordering two ridgelines—one impacted by transmission line installation and the maintenance roads, and the other with substantially natural areas. Between the two ridgelines is an intermittent drainage with riparian elements. The area includes good examples of riparian woodland, chaparral, oak woodland, coastal sage scrub, and chaparral communities. The entire unpaved campus area is designated critical habitat for the federally threatened coastal California gnatcatcher (*Poliioptila californica californica*).

The SEA is located within the El Monte United States Geological Survey (USGS) 7.5' California Quadrangle.

The SEA boundaries have a roughly triangular shape. The northern boundary begins in an arroyo with walnut woodland toward the west end of the North Entry Road, and goes southeastward along the border

DRAFT

of natural habitat, passing the justice buildings near the bottom of the transmission line ridge. The boundary continues along the base of the ridge to the border with the Rose Hills Memorial Park. On the east side, the boundary tracks the edge of the Rose Hills Memorial Park peripheral road towards the southeast, including a slope of probable restored white sage scrub. Where the peripheral road meets the crest of the southern ridge, the boundary turns northwest and goes along the southern ridge top, continues down off the ridge northwestward around the Rio Hondo College campus parking lots and buildings to connect with the small walnut woodland.

The northern ridge primarily contains non-native grassland, with a strong component of introduced mustards, but also a strong component of scattered native chaparral shrubs, such as elderberry (*Sambucus* spp.), sumacs (*Rhus* spp.); and in the ravines, dense growths of large chaparral shrubs, including coast live oak (*Quercus agrifolia*). Some of the slopes on the north ridge have prickly-pear shrub. The watercourse between the ridges is riparian with plants, such as mulefat (*Senecio douglasii*) and coyote bush (*Baccharis* spp.). The ravines on the north face of the southern ridge that border the watercourse have fine oak woodland, cherry woodland, and walnut woodland at the upper ends. The SEA provides examples of many of the common and cherished natural habitats of the County for study. Sign of coyote (*Canis latrans*), fox (probably *Urocyon cinereoargenteus*) and bobcat (*Lynx rufus*) can be noted while walking the transmission line northern ridge. The biotic communities within the SEA contain a variety of plant life and an abundant fauna.

Due to location near the extreme northwestern end of the Peninsular Ranges, the SEA is an excellent place to observe the geographical range variability of a number of species that are characteristic of the mountains to the south, and have their northernmost occurrences in the Puente Hills, such as the red diamond rattlesnake (co-occurring with the Pacific rattlesnake).

Wildlife Movement

The SEA is located in an area of potentially low to moderate value with regards to regional and local terrestrial wildlife movement. The Interstate 605 and industrial development that borders the freeway serves as a barrier for wildlife movement. The SEA is a gateway area for connectivity between the Peninsular Ranges of Southern California and the Transverse Ranges to the north. Naturally, they are connected by use of wildlife, particularly birds, insects, and plant propagules that are found along the San Gabriel River and Rio Hondo, which are only a 0.5 mile to the west of the Rio Hondo College. The San Gabriel River is designated by Audubon California as a State Important Bird Area (IBA), and extended arms of Semi-Natural habitat are important to connectivity for wildlife of the area and the region. Wildlife species could potentially use the SEA and possibly the ornamentally landscaped areas to facilitate movement and provide access to natural resources located in the Puente Hills. A wide variety of wildlife use linkages throughout the SEA from the extreme southeast up to the Rio Hondo College Wildlife Sanctuary, including mountain lion (*Puma concolor*) and a number of medium-sized mammals.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE RIO HONDO WILDLIFE SANCTUARY SEA

Criterion	Status	Justification
-----------	--------	---------------

DRAFT

A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	The SEA is critical habitat for the coastal California gnatcatcher.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Not Met	The SEA contains walnut woodland, which is restricted in distribution in the region of Southern California.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	The SEA contains walnut woodland, which is restricted in distribution in the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	This SEA is located on the eastern upland area of the San Gabriel River and is considered critical habitat for connectivity of the coastal California gnatcatcher. The largest population of the gnatcatcher in the County is on the west side of the San Gabriel River and the Interstate 605. Critical habitat in the SEA is on the east side of the San Gabriel River and the Interstate 605. The SEA is an arm extending to the rest of the gnatcatcher critical habitat and connecting to the rest of the Puente Hills SEA. The SEA is an important connecting and migration area for plants and wildlife of the Puente Chino Hills of the Peninsular Ranges.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The SEA is part of the education network of the public community colleges of the Los Angeles area. The area is used by the college for scientific study and research on native wildlife and plants. The college maintains field records on the biotic resources of the area. The SEA is relatively undisturbed. As a "choke point" for the wildlife corridor, it is an important area of research and study of connectivity.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	NotMet	The SEA has little disturbed areas, as well as recovering natural habitat.

In conclusion, the area described is an SEA because it contains A) core habitat for a threatened species; D) is an important choke point in a significant migration and connective corridor of the County and the region of Southern California; and E) is an important resource to the education community of the County because of its connective status and its natural and recovering habitats.

San Andreas SEA

Boundary and Resources Description

The San Andreas SEA is located in the western portion of the Antelope Valley in an unincorporated area of the County. The SEA is the second largest SEA and includes many diverse habitats. This is in large part due to the northwestern area being a meeting place for several diverse biomes and wildlife

DRAFT

~~corridors. There are five ecoregions that meet in this area and have biological species that extend along the SEA and San Andreas Fault in the County. These ecoregions include California Coastal Mountains; California Central Valley; Tehachapi Mountains, which extend to the southern Sierra Nevada; San Gabriel Mountains, which extend to other ranges in the Transverse Ranges; and the Antelope Valley, which is the western limit of the Mojave Desert. Wildlife corridors extend along the courses of the mountain ranges, as well as along the San Andreas Fault and Garlock Fault, which provide a great variety of habitats and frequent emergent water that is important for wildlife, plant movement and connectivity. The location and orientation of the SEA coincides with a segment of the San Andreas Fault Zone. The SEA includes a small portion of the western south-facing Tehachapi foothills, which are known for wildflower field displays in years of good rainfall. The SEA extends east and south across grasslands at the western tip of the Antelope Valley, and includes Quail Lake, a sag pond enhanced to receive water from the West Branch of the California Aqueduct. From Quail Lake, the SEA extends up the northern foothills of Liebre Mountain, Sawmill Mountain, and includes Portal Ridge; large portions of Leona Valley; Ritter Ridge, Fairmont and Antelope buttes; and portions of Anaverde Valley. It also includes a disjunct area that encompasses water bodies along the fault, Lake Palmdale, and Una Lake, with a terminus at Barrel Springs.~~

~~The Antelope Valley and adjacent desert slopes of the SEA are recognized by Audubon California as the Antelope Valley (Lancaster) Globally Important Bird Area (IBA). Near Lake Palmdale in the disjunct eastern section of the SEA is part of the Antelope Valley (Lancaster) IBA and near Barrel Springs is part of the Santa Clara River IBA.~~

~~The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Frazier Mountain, Lebec, La Liebre Ranch, Neenach School, Fairmont Butte, Little Buttes, Black Mountain, Liebre Mountain, Burnt Peak, Lake Hughes, Del Sur, Lancaster West, Sleepy Valley, Ritter Ridge, and Palmdale.~~

~~The northwestern tip of the SEA encompasses south-facing foothills at the western end of the Tehachapi Mountains, in the northwest corner of the County, on the eastern side of Tejon Pass.~~

~~From the Tehachapi Foothills, the southern boundary goes south-southeast along Interstate 5, including much of Peace Valley in the Gorman area, which is the broad faulted area that includes Gorman Creek. The SEA boundary crosses the Western Branch of the California Aqueduct, which is south of the junction of Interstate 5 and State Route 138. The boundary continues south along Interstate 5 until the point where the Liebre Mountain ridgeline dips to the highway, and the SEA boundary turns eastward and follows the ridgeline along the northern side of Liebre Mountain.~~

~~Along this section of Interstate 5 are several large underpasses for stream courses that are extremely important for wildlife connectivity across Interstate 5. The Angeles National Forest boundary is just east of the highway, and south of the aqueduct. Just north of the Liebre Mountain ridgeline, the San Andreas SEA borders the north, east, and south sides of the eighth unit of the Joshua Tree Woodlands SEA. This woodland is in an unnamed arroyo, and contains a population of the clonal growth form that Joshua trees (*Yucca brevifolia*) exhibit in colder and more fire-prone areas, sometimes referred to as *Yucca brevifolia* var. *jaegeriana*. The woodland is located near the westernmost limit of the range of the species, with a small number of stands and individuals known west of the Interstate 5. The SEA includes the northern slope area of the Angeles National Forest with its diversity of chaparral, grasslands, and oak and conifer forests.~~

~~After turning east from Interstate 5 and climbing uphill on the northern slope of Liebre Mountain, the SEA boundary crosses the ridgeline to the south to incorporate natural pristine areas of headwaters~~

DRAFT

~~for all the branches of Liebre Gulch, which are part of the headwaters for Piru Creek, the largest tributary of the Santa Clara River in Ventura County. The SEA boundary returns to the north face of Liebre Gulch in the vicinity of Sandberg. The boundary tracks the Sawmill Mountain Maxwell Road, along the broad ridgeline of the mountains and generally trends in a southeasterly direction. This ridgeline is the headwaters of Castaic Creek, which is the largest tributary of the Santa Clara River in Los Angeles County. Castaic Creek is above the Castaic Reservoir, which extends into Cienega Canyon and Fish Creek, which is federally designated critical habitat for the endangered arroyo toad (*Anaxyrus californicus*). In addition, maintenance of clean water in the source areas is critical for the species.~~

~~The boundary turns northeast where it meets Lake Hughes Road. This is an extremely important area of connectivity as the canyon along the Lake Hughes Road (Elizabeth Lake Canyon) drains to Castaic Creek and the Santa Clara River, whereas the Amargosa Creek that goes east and west from the Lake Hughes Road in the fault valley drains to the Antelope Valley in both directions. The junction is topographically broad and well vegetated though residential, which is excellent for wildlife connectivity in spite of a few houses.~~

~~The SEA boundary goes north at the junction with Lake Hughes Road and then skirts the Lake Hughes community's extension into Pine Canyon along the San Andreas Fault. In Pine Canyon, the boundary turns north and returns to its southeasterly direction, skirting the Lake Hughes development along the southern edge of Portal Ridge. Portal Ridge is entirely included in the SEA. A side extension of the southern boundary includes Lake Hughes, which is important for migrating waterfowl, with its sheltered position in the Fault valley. The boundary extends along the southern edge of Lake Hughes, Munz Lake, and Elizabeth Lake, and then trends southeast to go along the Leona Divide, including a large portion of Leona Valley.~~

~~The entire area along the San Andreas Fault is rich in wetlands and bogs, but Leona Valley has these in abundance, even in many yards. All of the wetlands in the San Andreas Fault valley and Portal Ridge are home to the greatest concentration of the tricolored blackbird in Southern California, many of which are year round residents. This bird species has experienced great population declines in recent years and is proposed for listing at both state and federal levels. In the community of Leona Valley, the southern SEA boundary goes along Lost Valley Creek and then along Leona Road to exclude some of the denser residential area in this section. The included area in Leona Valley has many of the bogs that line the Fault and the less populated farm areas along Portal Ridge north of Leona Road.~~

~~North of the Bouquet Canyon watershed, the southern SEA boundary dips south around an expansive area of drainages and bogs used by the tricolored blackbird on the old Ritter Ranch. From Ritter Canyon to the east, the boundary follows the old Ritter Ranch high road along the Sierra Pelona, crosses from 40th Street to the California Aqueduct along vegetation in the Anaverde Valley (where the boundary transitions from the Amargosa Creek drainage to the Anaverde Creek drainage), and then follows the aqueduct to the area where Anaverde Creek exits from the Fault valley. At the Lancaster Landfill boundary, the SEA boundary goes north and becomes the north SEA boundary at Verde Point.~~

~~The northern boundary of the SEA begins at Tejon Pass next to Interstate 5 and follows the Kern-Los Angeles County line eastward to its intersection with the western branch of the California Aqueduct in the western Tehachapi Foothills. This area along the Kern-Los Angeles County line is coincident with the designated critical habitat for the federally endangered California condor (*Gymnogyps californicus*), which is a bird that nearly went extinct and was saved by prodigious efforts in captive~~

DRAFT

~~breeding. The boundary then generally follows the Tehachapi foothills southward to Quail Lake. Here the northern SEA boundary crosses Highway 138 to include the northern foothills of the Liebre Mountains and fallow agricultural fields, which are important for raptor foraging. These fields are often oriented along the Los Angeles Aqueduct, which is a little south of the California Aqueduct in this area, or along the California Aqueduct itself.~~

~~The boundary eventually tracks along the northeast edge of Fairmont Reservoir (another breeding site for the tricolored blackbird), and turns northeast to include a patchwork of farmed areas between the Fairmont and Antelope buttes, which are known to have tricolored blackbird feeding grounds. The boundary makes an inclusive path to encompass the Broad Canyon Wash, the Fairmont and Antelope buttes, and the Antelope Valley California Poppy Reserve State Natural Reserve. These desert buttes are concentrated wintering grounds for birds of prey, and provide roosting sites that are surrounded by cultivated fields that support a plentiful food supply of rodents, rabbits, and hares. They are the most westerly buttes in the Mojave Desert, and with their proximity to the San Gabriel Mountains, have unique ecological relationships of scientific interest. Near the southern area of the buttes, the boundary follows agricultural fields along 130th Street West and then 135th Street West south to Munz Ranch Road (Willow Springs Road on some maps). Along 135th Street West, the boundary crosses Myrick Canyon where it spreads out onto the plain of the desert floor. The upstream areas of Myrick Canyon are included in the SEA.~~

~~The boundary tracks along the northwest side of Munz Ranch Road and then crosses to include Willow Springs Canyon, where Willow Springs Canyon is in its most undisturbed state. Where Willow Springs Canyon crosses the California Aqueduct, the northern SEA boundary turns east along the California Aqueduct as it passes along the northern base of Portal Ridge. Following the southern edge of the California Aqueduct, the boundary continues in a southeasterly direction to the east side of Ritter Ridge to Leona Siphon. A development along Joshua Tree Ranch Road near the summit of Ritter Ridge is excluded from the SEA. The SEA northern boundary turns east for roughly one quarter mile along the southern edge of a tributary to Amargosa Creek. Where the Amargosa Creek terminates Ritter Ridge, the SEA boundary crosses the creek and ascends along the ridgeline of an unnamed ridge to where it meets the southern boundary at Verde Point.~~

~~East across the State Route 14 is a disjunct part of the SEA that incorporates Lake Palmdale and Una Lake and extends along the Fault to 37th Street East, including the ridgelines north and south of Barrel Springs Road, which includes the sag ponds or Barrel Springs. The Palmdale Ditch is included in this part of the SEA. Many migrant birds using the desert water features can be observed at these artificial lakes and the natural springs of this area during the spring and fall migration.~~

~~The gap between the two portions of the SEA includes the Antelope Valley Landfill, disturbed lots, and State Route 14.~~

~~The majority of land within the SEA lies within unincorporated area of the County. Other jurisdictions include the Angeles National Forest, the City of Palmdale, and the City of Lancaster.~~

Wildlife Movement

~~The SEA includes several important linkages for wildlife movement. The foothills in the western-most part of the SEA are an important linkage between the San Gabriel Mountains, the Tehachapi Mountains, and the Coastal Ranges. This linkage to the Tehachapi Mountains is important because they connect to the southern-most extent of the Sierra Nevada Mountains. The Tehachapi Mountains represent the only mountain linkage from the Transverse Ranges and the Coast Ranges to the Sierra~~

DRAFT

Nevada Range. This feature may be an important topographic reference for migrating birds, as well as providing high elevation foraging grounds along the migratory route. The several ranges that meet at the western end of the SEA, provide a valuable link for gene flow between divergent subspecies, varieties, and populations of many species. The SEA includes numerous drainages that extend onto the Antelope Valley floor towards resources, such as the Fairmont and Antelope buttes. These washes provide an important linkage for animals traveling between the Valley floor, the buttes and the western part of the San Gabriel Mountains. In addition, Anaverde Creek, Amargosa Creek, and Pine Canyon facilitate east-west wildlife movement through the mountains, Portal Ridge, and Ritter Ridge. Tributary drainages from the Santa Clara River, such as Elizabeth Lake Canyon and San Francisquito Canyon, connect coastal drainages and the coastal ecoregion to the Fault and interior watersheds. The frequency of valuable riparian communities along this travel route, which are located within an otherwise arid climate, further contributes to the SEA's importance for wildlife and habitat linkages in the region.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE SAN ANDREAS SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Not met Met in Future?	Although there are several listed species that occur within the SEA, this criterion is not met due to the lack of known core population areas. The far northwestern border with Kern County is the edge of critical habitat for the California condor. The tricolored blackbird may soon be listed and has its largest population in Southern California within the SEA.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The SEA encompasses a series of marshes and sinks concentrated along the San Andreas Fault Zone, which are both unique and restricted in distribution. The Fairmont and Antelope buttes represent a unique habitat due to their location, as the most westerly buttes of the Mojave Desert and their close proximity to several geographic regions. As the confluence of a number of major geographical areas, the Mojave Desert, the San Gabriel Mountains of the Transverse Ranges, the Coastal Ranges, and the Tehachapi Mountains produces a unique and regionally rare flora that represents a transition between desert, foothill, and several montane environments.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The confluence of five major geographical areas—the Mojave Desert, the San Gabriel Mountains, the Coastal Ranges, the Tehachapi Mountains, and the Central Valley—has produced the most unique and diverse flora found in the County, and represents a transition between desert, foothill, and montane environments. The SEA also includes the southern limit of the foothill woodland community, blue oak, gray or foothill pine, and California buckeye, rare relic stands of Great Basin

DRAFT

	Criterion	Status	Justification
			sagebrush scrub, and rare wildflower fields.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The Fairmont and Antelope buttes provide vital habitat to many wide ranging species, which forage in outlying habitat, but use the buttes for nesting, roosting, denning, and refuge. The buttes also serve as concentrated wintering grounds for birds of prey, which are rare in the County, and which forage on grassland and agricultural fields in the vicinity. Lakes and other wetland areas along the Fault and throughout the SEA provide breeding habitat for amphibians and feeding habitat for migrating birds that traverse the slopes adjacent to the Mojave Desert. The Fault is one of the principle wildlife corridors and connective areas for in the County. Major drainages (Santa Clara River, San Francisquito Canyon, and Lake Elizabeth Canyon) run from the coast through the San Gabriel Mountains and end at the Fault, which also has extensive riparian habitat that facilitates migration. The Fault provides the final westernmost linkage to the Mojave Desert (Antelope Valley). The tricolored blackbird is a year round resident of the SEA.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The transition of several habitat types including creosote bush scrub, Joshua tree/California juniper mixed woodland, and desert chaparral, makes the SEA valuable for educational and scientific reasons. The close proximity of the Fairmont and Antelope buttes to the San Gabriel Mountains renders them unique in their species composition and ecological relationships and, therefore, of interest to scientists. The concentrated diversity of vegetation types, particularly in the western half of the SEA, creates an outstanding opportunity for educational use. This area also harbors the southern limit of the foothill woodland community; blue oak, gray or foothill pine, and California buckeye, as well as rare relic stands of great basin sagebrush scrub.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The slopes of Ritter Ridge support one of the most pristine mixed stands of Joshua tree and California juniper in Los Angeles County. The location of the SEA at the confluence of five major geographical areas, the Mojave Desert, the Central Valley, the San Gabriel Mountains of the Transverse Ranges, the Coastal Ranges, and the Tehachapi Mountains has produced a community rich area with desert, foothill, and montane environments. The SEA encompasses large, mostly undisturbed examples of all of these communities.

In conclusion, the area is an SEA because it contains B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, and migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original

natural biotic communities in the County.

San Dimas Canyon and San Antonio Wash SEA

Boundary and Resources Description

The San Dimas Canyon and San Antonio Wash SEA is located along the cismontane foothills of the eastern San Gabriel Mountains. Generally, the SEA is centered on the mouths of four major canyons, which flow from the mountains and interconnecting terrain. From east to west, these canyons include San Antonio Canyon above the City of Claremont as one component; and Live Oak, Marshall, and San Dimas canyons above the cities of La Verne and San Dimas as a second component. The SEA incorporates areas with diverse natural habitat ranging from high elevations to the foothill alluvial areas of two of the major drainages of the San Gabriel Mountains. San Dimas Canyon is a tributary of the San Gabriel River. San Antonio Wash is a tributary of the Santa Ana River.

The SEA is found within the Mount Baldy and Ontario U.S. Geological Survey (USGS) 7.5' California Quadrangles.

Over most of its boundaries, particularly to the north, east, and west of both the San Dimas Canyon and San Antonio Wash components, the SEA is bordered by open space within the Angeles National Forest. Generally to the south, however, the borders are mostly defined by the edge of urban development within the San Gabriel Valley. The San Dimas Canyon component covers approximately 5,500 acres and includes portions of Live Oak, Marshall, and San Dimas canyons. The smaller component, San Antonio Canyon, covers approximately 1,200 acres of the San Antonio Canyon alluvial outwash. In total, this SEA encompasses 6,727 acres.

In general, the topography of the SEA is severe, consisting of steep-walled canyons and narrow ridgelines. Elevations range from a high of approximately 3,000 feet above mean sea level (MSL) along the ridges of San Dimas Canyon, to a low of approximately 451 feet above MSL in San Antonio Wash. Several major drainages and numerous tributaries exit the San Gabriel Mountains through this SEA.

The wide range of elevation, topography, slope aspect, and geology represent a wide array of physical habitats within this SEA. Consequently, a number of plant communities exist, including grasslands, riparian, shrublands, woodlands, and forests. Within these major community types, there are many sub-communities, which vary according to plant species dominance. This area contains the last remaining relatively well-developed lower montane riparian habitat in the eastern County. Dammed drainages have created significant reservoirs or flood control basins in the SEA. The SEA is within several jurisdictions including the Angeles National Forest, the unincorporated area of the County, the City of Claremont, the City of Glendora, the City of La Verne, and the City of San Dimas.

The more westerly component of this SEA generally includes portions of the lower watersheds of San Dimas, Marshall, and Live Oak canyons, which is part of the San Dimas Canyon component. The San Dimas Canyon watershed is part of the Experimental Forest section of the Angeles National Forest. Experiments were conducted and data was collected here during the latter half of the 20th century to determine the relationships among rainfall, topography, vegetation, and runoff. Much of the work and results influenced flood control in the Los Angeles Basin and even other areas of the U.S. The area was carefully protected through very limited and monitored access. The terrain chiefly includes undisturbed natural habitats of rocky canyon walls and canyon forest, riparian areas of many vegetation types, coniferous and oak forest, chaparral, and grassland. A few slopes were altered with vegetation removal in order to experiment on the effect of vegetation, and some of these are still

grassland.

~~This SEA area on the border of the granitic San Gabriel Mountains has unusual rock strata, such as the Glendora Volcanics. Much of the grassland is natural and has unusual vegetation, such as wildflowers that prefer clay substrates. Not too distant from this area are critical habitat areas for the endangered thread-leaved brodiaea (*Brodiaea filifolia*). Some of these brodiaea and other rare wildflowers could occur in appropriate habitat of the SEA in undiscovered populations.~~

~~Beginning at Johnstone Peak in the west, the western boundary follows the ridgeline separating Big Dalton Canyon and San Dimas Canyon. Just before this ridgeline is intersected by Big Dalton Canyon Road, the SEA boundary turns east. From the area of Big Dalton Canyon Road, the northern boundary follows and crosses over a series of ridgelines to include the upper portions of several tributary canyons. It continues in this fashion in a southeasterly direction eventually meeting and following the Sunset Ridge Fire Road (Sunset Peak Motorway), which separates Wolfskill and Marshall canyons. The tributaries San Dimas Canyon include Lodi, West Fork of San Dimas, and San Dimas from near the junction with Wolfskill Canyon. The lower section of Wolfskill Canyon with and below the Wolfskill Falls is included in the SEA. The upper section of Wolfskill is not included in the SEA, but much of Marshall Canyon watershed is included, along with watersheds of Live Oak and Webb canyons in the City of Claremont.~~

~~A large lobe of the SEA extends from the Sunset Ridge Fire Road on the dividing ridgeline, to include lush canyon forests and chaparral of the slopes above the City of La Verne and City of Claremont. Most of this lobe is in municipal or private ownership. The Angeles National Forest boundary is about a 0.1 mile south of the Sunset Ridge Fire Road. The eastern boundary leaves the fire road and travels south along a ridgeline, including Live Oak Canyon in the SEA, but separating out the more developed watersheds of Palmer, Cobal, Burbank, and Gail canyons in the City of Claremont. A finger of the SEA includes the lush riparian oak forest of Webb Canyon to the edge of development. The lobe of the SEA excludes an area around the residences and equestrian areas that surround Live Oak Reservoir. Live Oak Canyon Reservoir and its riparian oak woodland is included as far south as Base Line Road. The ridges and dissected canyons that border Live Oak Reservoir are included as far south as Base Line Road. However, the flat area of the ridge around Live Oak Reservoir and development in the periphery are excluded. The northwestern edge of the lobe includes the riparian area and slopes of Marshall Creek, but excludes developed areas, such as the Marshall Canyon Regional Park and Golf Course. The lobe boundary returns north into the Angeles National Forest at the Sunset Ridge Fire Road along the edge of Marshall Creek and the western ridge of Marshall Canyon.~~

~~From Sunset Ridge Fire Road, the southern boundary of the SEA is within the Angeles National Forest and follows the ridgeline that includes the watershed of San Dimas Canyon. The San Dimas Reservoir, with good habitat for waterfowl, is included in the SEA. The SEA extends a finger out of the Angeles National Forest along San Dimas Canyon road to include the riparian habitat along the watercourse, which is a rare example of the lowland riparian community. From the Angeles National Forest boundary and rocky cliffs above the west side of San Dimas Canyon, the SEA boundary follows the ridge of Lodi Canyon (tributary of San Dimas Canyon) to Johnstone Peak.~~

~~The eastern, disjunct segment of the SEA (San Antonio Wash) follows the San Bernardino-Los Angeles County line as its eastern boundary from about a 0.5 mile upstream of the San Antonio Dam through the San Antonio debris basin, past the San Antonio Dam, to the natural extent of alluvial fan vegetation south of the Interstate 210. This is at an area about a 0.1 mile north of Base Line Road. Downstream of the San Antonio Dam has the best example of arroyo or wash vegetation that remains in the County, and it extends onto the adjacent alluvial fan. The vegetation is a dry form of~~

DRAFT

coastal sage scrub, with included desert plants that are adapted to coarse substrate. The vegetation is much more dense and stable than the alluvial fan in the arroyos behind Santa Fe Dam (San Gabriel Canyon SEA) and Hansen Dam (Tujunga Valley Hansen Dam SEA). From its southern point, the SEA turns north to include the natural alluvial fan vegetation and border on the existing residential development on the alluvial fan. At the intersection of the San Antonio Wash with Mount Baldy Road, the SEA boundary follows the southeast side of Mount Baldy Road to the watershed of Chicken Canyon, which is a tributary of San Antonio Wash. The boundary crosses the road and includes the undeveloped part of Chicken Canyon. The boundary follows the minor ridgeline up to Potato Mountain, and goes along the south ridge of Evey Canyon back to cross Mount Baldy Road and return to the San Bernardino-Los Angeles County line in the San Antonio Debris Basin. Evey Canyon is outside the SEA, but is a preserve of the Claremont Colleges, and has excellent riparian canyon habitat. The SEA designation acknowledges the need to protect the Evey Canyon watershed. Small tributary watersheds of San Antonio Canyon with chaparral vegetation are included with the Chicken Canyon area.

Wildlife Movement

Wildlife movement within the SEA takes on two major forms. First, due to the extreme intervening topography, it is logical to expect considerable movement of wildlife up and down the many sizeable drainages, which course through this SEA and connect the forest interior with foothill areas. The larger the watershed of the drainages, the greater the volume of movement. Consequently, this type of movement occurs on a seasonal and more frequent basis, particularly for large mobile mammals, such as American black bear, mountain lion, coyote (*Canis latrans*), bobcat (*Lynx rufus*) and mule deer (*Odocoileus hemionus*), whose full range of habitat needs are typically met over broad areas.

The second major type of movement occurs across the flanks of the foothills and lower mountains, in an east-west direction. Particularly for riparian-favoring migratory birds, a corridor linking lower elevational riparian habitats in the SEA is expected to be of high use and importance. In addition to providing essential habitat for resident riparian birds, this SEA contains some of the best developed riparian habitat for birds, which are seasonal visitors to the cismontane region of the County.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

Criteria Analysis of the San Dimas Canyon and San Antonio Wash SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Not Met	Although the SEA contains rare plant populations, it does not contain a core population of a listed species and therefore does not meet this criterion. The lower slopes in and around San Dimas Canyon support one of the largest populations of the coastal cactus wren in the County, which is a subspecies that is very threatened throughout its range, although not officially recognized by listing.
B)	On a regional basis, biotic communities, vegetative associations,	Met	The SEA contains habitat of the rare rock monardella. In addition, several plant communities within this SEA are

DRAFT

	and habitat of plant or animal species that are either unique or are restricted in distribution.		CDFW highest priority communities due to their restricted distribution in the Southern California region, including walnut woodland, oak riparian woodland, southern willow scrub, coastal sage scrub, and alluvial fan scrub.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	All of the plant communities and habitats mentioned as being restricted in distribution on a regional basis, are also restricted in distribution within the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The major canyons within this SEA support well-developed and diverse riparian woodlands, as well as a source of perennial water. These represent important stopover and overwintering areas for a wide variety of migratory birds, as well as essential habitat for resident species of fauna and flora. These canyons also support seasonal and more frequent movement for wide-ranging mammals, which must move over large areas to fulfill their habitat requirements. The federally threatened California gnatcatcher has been sighted (2010) in the Glendora foothills, and probably maintains a small population along the lowest slopes of the San Gabriel Mountains.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Not Met	The SEA does not contain biotic resources that are clearly an extreme in physical/geographical limitations, or represent unusual variation in a population or community, and therefore does not meet this criterion. However, the extreme localization of several species of plants in the SEA may indicate geographical processes that are not well understood at this time that merit scientific inquiry.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Virtually all of the native biotic communities within this SEA are relatively undisturbed over most of their extent. Because urbanization throughout much of the County's foothill regions has removed large expanses of these communities, those in the SEA are particularly important to the County's natural heritage.

In conclusion, the area is an SEA because it contains B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; and F) areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.

San Gabriel Canyon SEA

Boundary and Resources Description

The San Gabriel Canyon Significant Ecological Area (SEA) is located along the cismontane foothills of the eastern section of these mountains. Generally, the SEA is centered on the mouths of three major canyons, which flow from the mountains and interconnecting terrain. From west to east these include, Santa Anita, Monrovia and Sawpit, and San Gabriel canyons, which are located above the cities of Sierra Madre, Arcadia, Monrovia, Duarte, Bradbury, Irwindale, and Azusa. A substantial part of the

DRAFT

~~eastern and southern part of the SEA along the San Gabriel River is in the Audubon California–designated State Important Bird Area (IBA) of the Los Angeles Flood Control Basin IBA. The San Gabriel River has largely been dammed and channelized, but with infrequent clearing of the detention basins and wash areas, substantial parts of the San Gabriel River have reverted to riparian habitat or the even more rare alluvial fan habitat, and this attracts many resident birds, as well as numerous spring and winter migrants.~~

~~The SEA is found within the, Mount Wilson, Azusa, San Dimas, and Glendora U.S. Geological Survey (USGS) 7.5' California Quadrangles.~~

~~Over most of its boundaries (north, east, and west), the SEA is bordered by open space within the Angeles National Forest. However, generally to the south, the borders are defined by the edge of urban development within the San Gabriel Valley. The SEA begins in the west at the peak of Mount Wilson within the Angeles National Forest. Traveling east, the northerly boundary follows a major east-west trending ridgeline to Pine Mountain. This ridgeline defines the separation between the watershed of the San Gabriel River West Fork to the north, and the Santa Anita, Sawpit, and lower San Gabriel canyons to the south. These front range canyons are tributaries of the San Gabriel River.~~

~~At Pine Mountain, the boundary turns south to follow the ridgeline that is the western border of the San Gabriel River, and turns east onto a secondary ridge, and descends towards the San Gabriel River near the Morris Reservoir Dam. This easterly boundary crosses the San Gabriel Canyon at Morris Dam and climbs the adjacent ridgeline to Glendora Ridge and the Glendora Ridge Motorway. The southerly boundary follows the motorway to the west, to the point near the mouth of the San Gabriel Canyon where the motorway leaves the ridgeline. The SEA boundary turns north towards the San Gabriel River, and descends to the opening of the San Gabriel Canyon into the Los Angeles Basin. This is between the Glendora Ridge and the mountains near Fish Canyon. The boundary turns along the southeast side of the San Gabriel River floodplain and follows the east side of the San Gabriel River flood control channel. A development near the mouth of Roberts Canyon that is just north of the river mouth has been excluded from the SEA.~~

~~In the mouth of the San Gabriel Canyon is a population of the San Gabriel Mountains live forever (*Dudleya densiflora*), which is unusual in that it has multiple dense flower clusters, whereas other live-forevers have one or several flower stalks with spaced blooms. This live forever is extremely limited in range and occurs only on the slopes of granitic rubble and canyon walls in the nearby south face of the San Gabriel Mountains. Another population is on private land about one mile upstream of the canyon mouth, on the north side slope of the Glendora Ridge. Another live forever population is upstream in nearby Fish Canyon, which is a little downstream of the Fish Canyon Falls. Collections have been made from Mystic Canyon to the east, and Van Tassel Canyon to the west.~~

~~The mouth of San Gabriel Canyon and nearby canyons are the principle area for the San Gabriel bedstraw (*Galium grande*), which is another local endemic. The only known populations of the bedstraw and the San Gabriel Mountains live forever on the planet occur in the County in this small area of the San Gabriel Mountains.~~

~~The Los Angeles Flood Control Basin IBA covers all of the SEA in the San Gabriel River and downstream at the Santa Fe Dam Recreation Area. Furthermore, the IBA extends upstream beyond the SEA to the confluence area of the West, North, and East forks of the San Gabriel River in the Angeles National Forest, and it extends downstream beyond Santa Fe Dam to the Whittier Narrows Dam.~~

DRAFT

A finger of the SEA extends along the San Gabriel River, south of its confluence area with Fish and Van Tassel canyons to pass under the Interstate-210. The finger boundary enlarges around the Santa Fe Flood Control Basin and Recreation Area to include one of the last remaining natural alluvial fan habitats in the County. The Santa Fe Flood Control Basin is one of the most unusual vegetation habitats in the County, and has special sensitive species.

The main SEA boundary continues just west of the Van Tassel Canyon confluence along the north side of the Encanto Equestrian Center, along the northern extent of development in the City of Duarte. A lobe of the SEA encloses the natural habitat of the steep watershed areas of Spinks and Maddox canyons, extending to the edge of development in the City of Bradbury. The ridge bordering the southeast side of Bliss Canyon is the western edge of the lobe, and the boundary crosses Bliss Canyon at its upper end near the Van Tassel Truck Trail. At this point the boundary of the SEA has reentered the Angeles National Forest. After crossing Bliss Canyon, the boundary follows the southern ridgeline of Spanish Canyon westward to cross out of the Angeles National Forest, tracking around the northern arm of the City of Monrovia. The Sawpit Debris Basin is included in the SEA as is the undeveloped part of Monrovia Canyon Park. To the west of Monrovia Canyon, a lobe of the SEA extends along the undeveloped ridges of the San Gabriel Mountains bordered by the urban edges of the City of Monrovia and City of Arcadia. These communities extend into the mountains where the cities have municipal water rights. The southern boundary skirts the edge of development in Santa Anita Canyon, but includes the Santa Anita Debris Basin, Arcadia Natural Park, Big Santa Anita Dam and Reservoir, and the Santa Anita Canyon stream course above the Dam, which has numerous lease hold cabins north of the 1600 foot elevation contour. The boundary reenters the Angeles National Forest just north of Arcadia Natural Park.

The southern ridge of Sawpit Canyon, from its dam to about a 0.5 mile upstream has a population of the endangered San Gabriel bedstraw (*Galium grande*), which is an endemic species of highly restricted distribution. It occurs only on the south slopes of the western section of the San Gabriel Mountains.

Within the SEA, just to the south of Arcadia Natural Park is a Santa Anita Canyon tributary, Clamshell Canyon. On the south banks and ridge of Clamshell Canyon is critical habitat for the federally-endangered Braunton's milk vetch (*Astragalus brauntonii*), which is a locoweed that prefers interbedded sandstone and carbonate substrate, probably deposited near the coastline of former oceans. Very limited areas of this substrate occur at the boundary of the San Gabriel Mountains in this area. Most of the rocks of the San Gabriel Mountains are igneous granites and metamorphic rocks.

Santa Anita Canyon has some stands of Pacific madrone (*Arbutus menziesii*), which is a plant known elsewhere from the Pacific coast north of Santa Barbara to British Columbia. The Santa Anita stands are isolated occurrences, which is one of the few places madrone is found between Santa Barbara and Baja California.

Near the confluence with Winter Creek in the vicinity of Chantry Flats, the southern boundary of the SEA turns west and climbs the southern ridgeline of Winter Creek, including Winter Creek watershed in the SEA and excluding San Olene Canyon on the south. The boundary follows the ridgeline, marking the southern limits of the Winter Creek watershed to Mount Harvard, and then travels along the Harvard ridgeline to Mount Wilson.

The SEA is comprised of three major canyons: San Gabriel, Sawpit, and Santa Anita. In general, the topography of the SEA is severe, consisting of steep walled canyons and narrow ridgelines. Elevations range from a high of approximately 5,710 feet above mean sea level (MSL) at Mount Wilson, to a low

DRAFT

of approximately 660 feet above MSL in San Gabriel Canyon. Numerous drainages and tributaries of the main canyons are included in the SEA and exit the San Gabriel Mountains into the Los Angeles Basin through this SEA.

The wide range of elevation, topography, slope aspect, and geology represent a wide array of physical habitats within this SEA. Consequently, a number of plant communities exist, including grasslands, riparian, shrublands, woodlands, and forests. Within these major community types, there are many sub-communities, which vary according to plant species dominance. Of particular note, this SEA contains the last remaining relatively well-developed lower montane riparian habitats in the eastern County and dammed drainages that have created significant reservoirs or flood control basins in Sawpit and Santa Anita canyons. Enclaves of sensitive plant species and vegetation habitats are found here. Other jurisdictions within the SEA include the unincorporated area of the County, the City of Arcadia, City of Monrovia, City of Bradbury, City of Irwindale, City of Duarte, City of Azusa, and the City of Glendora.

Wildlife Movement

Wildlife movement within the SEA takes on two major forms. First, due to the extreme intervening topography, it is logical to expect considerable movement of wildlife up and down the sizeable drainages, which course through this SEA to connect the forest interior with foothill areas. Consequently, this type of movement occurs on a seasonal and more frequent basis, particularly for large mobile mammals whose full range of habitat needs are typically met over broad areas, including American black bear, mountain lion, coyote (*Canis latrans*), mule deer (*Odocoileus hemionus*), gray fox (*Urocyon cinereoargenteus*) and other medium-sized mammals.

The second major type of movement occurs across the flanks of the foothills and lower mountains, in an east-west direction. Particularly for riparian favoring migratory birds, a corridor linking lower elevation riparian habitats in the SEA is of high use and importance. In addition to providing essential habitat for resident riparian birds, this SEA contains some of the best developed riparian habitat for birds, which are seasonal visitors to the cismontane region of the County.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE SAN GABRIEL CANYON SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	The SEA contains a core habitat area for the endangered plant Braunter's milkvetch. The upper San Gabriel River is a core habitat of several native fishes, one of the last areas where three of five original natives occur together: federally threatened Santa Ana sucker, and the arroyo chub and Santa Ana speckled dace, which is of state concern. All three live in the San Gabriel River in the SEA area. A local population of the speckled dace is known from the mouth of Fish Canyon. The very rare San Gabriel bedstraw and San Gabriel Mountains live-forever only

DRAFT

	Criterion	Status	Justification
			occur in this area of the world.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The SEA contains habitat of extremely rare plants: San Gabriel bedstraw and the San Gabriel Mountains dudleya. In addition, several plant communities within this SEA are CDFW highest priority communities due to their restricted distribution in the Southern California region. These communities include walnut woodland, oak riparian woodland, southern willow scrub, coastal sage scrub, and alluvial fan scrub. The federally endangered California gnatcatcher has been recently sighted in the Glendora foothills, and probably maintains a small population along the lowest slopes of the San Gabriel Mountains.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	All of the plant communities and habitats mentioned as being restricted in distribution on a regional basis, are also restricted in distribution within the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The three major canyons within this SEA support well developed and diverse riparian woodlands, as well as year-round water sources. These represent important stopover and overwintering areas for a wide variety of migratory birds, as well as essential habitat for resident species. These canyons also support seasonal and more frequent movement for wide-ranging mammals, which must move over large areas to fulfill their habitat requirements.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The SEA contains biotic resources that are of scientific interest for their very restricted distributions: Braunton's milkvetch San Gabriel bedstraw, San Gabriel Mountains live-forever, and a local isolated population of Pacific madrone. The population of Santa Ana speckled dace in Fish Canyon may be the remaining extreme western extent of its population.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Virtually all of the native biotic communities within this SEA are relatively undisturbed over most of their extent. Because urbanization throughout much of the County's foothill regions has removed large expanses of these communities, those in the SEA are particularly important to the County's natural heritage.

In conclusion, the area is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, and migrating grounds, which are limited in availability in the County; E) populations of scientific interest because of very restricted distributions and isolated populations; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Santa Clara River SEA

Boundary and Resources Description

The Santa Clara River SEA extends along the entire County reach of the Santa Clara River, primarily within unincorporated areas of the County. The SEA encompasses a wide variety of topographic features and habitat types, as well as major tributaries—all of which contribute to this diversity. It is a major biotic corridor for the County (and Ventura County). The orientation and extent of the SEA depends upon the surface and subsurface hydrology of the Santa Clara River, from its headwaters, tributaries, and watershed basin, to the point at which it exits the County's jurisdiction. Nearly all of the SEA is designated by Audubon California as a Globally Important Bird Area (IBA). The Santa Clara River IBA extends beyond the SEA in both upstream and downstream directions (across Soledad Pass to the Barrel Springs area in the Antelope Valley and through Ventura County to the mouth of the River at the Pacific Ocean).

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Pacifico Mountain, Acton, Agua Dulce, Sunland, San Fernando, Mint Canyon, Oat Mountain, Newhall, and Val Verde.

The SEA covers a wide variety of topographic features and habitat types, including parts of the watershed tributaries. The biological and ecological functionality of the SEA is integrally linked to the Santa Clara River basin for its entire length. The bio-geographic limits of the SEA would extend downstream through Ventura-Los Angeles County line to its mouth at the Pacific Ocean, and encompass significant tributary drainages of Ventura County (Piru Creek, Sespe Creek, Santa Paula Creek, Wheeler Creek, etc.).

The eastern portion of the SEA follows natural contours at the headwaters of the watershed to incorporate much of upper watershed of Soledad Canyon (which becomes the Santa Clara River), the Kentucky Springs and the Aliso Canyon basins, and the downstream unnamed tributaries of the Santa Clara River to Arrastre Creek. This includes the watershed southern headwater areas within the Angeles National Forest. The headwaters of both Kentucky Springs and Aliso Canyon are in the Angeles National Forest, in semi-arid chaparral and desert scrub habitat; however, the drainages themselves support vegetation of desert and interior riparian habitat, which ranges from Great Basin sagebrush in Kentucky Springs Wash to dense, mature, willow cottonwood-sycamore woodlands along permanent streams in Aliso Canyon. The surrounding uplands in the basins support pinyon-juniper woodlands, chamise, mountain mahogany, and manzanita-dominated chaparral, buckwheat scrub, and ruderal lands. The alluvial plain formed along the southern margin of the Santa Clara River basin below these canyons supports intact, high diversity xeric alluvial fan sage scrub. Alluvial terraces within both drainages have been extensively cultivated for orchard crops and dryland agriculture, and in more recent years, rural and urban type residential developments have encroached on the watersheds. The Kentucky Springs basin has a large population of Parish's Great Basin sagebrush (*Artemisia tridentata* ssp. *parishii*), which is considered rare and sensitive in the County. A population of the federally threatened red legged frog (*Rana draytonii* FT, SC) is known to inhabit and breed in the Aliso Canyon watershed. Blum Ranch and another area on Aliso Canyon Road are disturbed, with farming development, but important to continuity of the SEA. The Santa Clara River IBA extends in a branch upstream to include Blum Ranch.

The boundary follows the Santa Clara River channel downstream through the Acton basin, paralleling Soledad Canyon Road on the north side, following the toe of the slope of the San Gabriel Mountains to the south. Boundaries continue along the channel margins to the southwest from Acton to Arrastre

DRAFT

~~Creek, where the southern boundary follows watershed contours to take in four upper tributary channels (Arrastre, Moody, and Bootleggers). Downstream from Acton, there are developed areas as along the Santa Clara River. From a little upstream of the Arrastre Creek confluence to a little downstream in the vicinity of the railroad stop of Lang (about 13 miles of river), the floodplain of the Santa Clara River is designated critical habitat for the federally endangered arroyo toad (*Anaxyrus californicus*). Some of the confluence area of Mill Canyon is also critical habitat for the arroyo toad. Part of the area of critical habitat for the toad was also proposed as critical habitat for the state and federally endangered unarmored threespine stickleback (*Gasterosteus aculeatus williamsoni*), which is a small three-inch fish that essentially only occurs in the County. It once was widespread throughout the Los Angeles Basin and beyond, but is now restricted to the upper Santa Clara River. The proposal for critical habitat was never approved, and this is now referred to as “essential habitat” for the fish. The type area for the fish is the Arrastre Creek, where it was first collected and described with a museum specimen.~~

~~The habitat along the Santa Clara River supports the largest community of riparian obligate birds between Santa Ynez River in Santa Barbara County and the Prado Basin in Riverside County. In the Soledad Canyon stretch are breeding summer tanager (*Piranga rubra*) and other desert species, along with some instances of least Bell's vireos (*Vireo bellii pusillus*), coastal cactus wrens (*Campylorhynchus brunneicapillus sandiegensis*), and southwestern willow flycatchers (*Empidonax traillii extimus*) from the coastal influence areas. The area is notable for having a combination of species that are characteristic of the desert and characteristic of coastal influence.~~

~~Just west of the confluence with Arrastre Creek the northern boundary loops up to the slopes of Parker Mountain and the eastern watershed of Hughes Canyon around the basal contours of significant rock outcroppings above the river basin, and on the south side, around the Mill Canyon tributary basin. The rocky buttes on the north side of the river, while only a minor part of the watershed of the river, provide important nesting, roosting, and sheltering habitat values for bats, birds of prey, and other sensitive species foraging along the river corridor. The boundaries stay at the river margins west to the watersheds of two northern tributaries, Nellus and Bobcat canyons. These drainages were identified by the South Coast Wildlands Project as important to connectivity across the Santa Clara River between the western and eastern highland areas of the San Gabriel Mountains.~~

~~At the Agua Dulce Canyon drainage, the northern boundary loops around the watershed, including the Vasquez Rocks County Natural Area. Agua Dulce Canyon has a permanent stream and supports high quality riparian habitat from the confluence with the river to the intersection with State Route 14. The Santa Clara River IBA extends upstream to include about one mile of the Agua Dulce Canyon.~~

~~The Agua Dulce underpass of State Route 14 is an important crossing of the highway barrier for wildlife. From that point, north riparian areas exist where the creeks (Agua Dulce and Escondido) pass through Vasquez Rocks County Natural Area. The Agua Dulce Canyon extension was included in the SEA for its value as a wildlife corridor to provide connectivity across the Santa Clara River between the western and eastern highland areas of the San Gabriel Mountains. The extension includes the watershed of Bee Canyon, which is a downstream tributary of the Santa Clara River. Bee Canyon has an important population of the federally endangered slender horned spineflower (*Dodecahema leptoceras*) in its broad, floodplain area. In the Bee Canyon slopes of coastal sage chaparral, the federally threatened coastal California gnatcatcher (*Polioptila californica californica*) is sometimes resident. The Bee Canyon area has some underpasses of the State Route 14 that could be used by smaller wildlife if maintained unlogged. The extension includes upper watersheds of Spring and Tick canyons to enhance the connective area. Beyond upper areas of Tick Canyon, the SEA boundaries cross Mint Canyon into the Angeles National Forest and the watershed of Rowher Canyon. The SEA~~

DRAFT

~~continues to the upper reaches of Rowher Canyon onto the main ridgeline of the Sierra Pelona. At the Mint Canyon crossing, just southwest of the community of Sleepy Valley, a lobe of the SEA extends along Mint Canyon to capture riparian woodlands of coast live oak, with a number of heritage trees (diameters greater than 36 inches). Residences are scattered and the natural communities of chaparral are intact on the canyon slopes.~~

~~The southern boundary of the SEA opposite the confluence with Agua Dulce Canyon includes the flood plain. The SEA dips southward into the lower portion of Bear Canyon (tributary of Santa Clara River) and includes undeveloped alluvial terrace slopes of the river downstream of Bear Canyon. The flood plain is a narrowed part of the SEA in the vicinity of Lang, which is a railroad stop on the transcontinental railroad line that runs the length of the Soledad Canyon. Downstream from Lang, the SEA expands to the southern slopes between Lang and Oak Spring Canyon, adjacent to the river channel. Downstream of Oak Canyon, the SEA narrows to the flood plain, passes Sand Canyon, and reaches the west ridge of Sand Canyon. A broad finger of the SEA goes south along the ridgeline of the Sand Canyon watershed, where the finger expands when it reaches the watershed of Placerita Canyon.~~

~~The alluvial fans of Oak Springs Canyon and Sand Canyon are important recharge grounds for the river aquifer. Surface flows from both canyons enter the Santa Clara River basin through natural, unconfined channels. Recognizing the importance of the Sand Canyon drainage, the SEA boundaries are drawn to encompass the entire upper Sand Canyon watershed, which is largely natural with scattered residences, as well as the Sand Canyon tributary, Bear Canyon. Most of the upper Sand Canyon and its Bear Canyon tributary are within the Angeles National Forest, and Sand Canyon originates on the peak of Magic Mountain. These canyons form a natural movement zone for wildlife traversing among the western end of the San Gabriel Mountains, the eastern end of the Santa Susana Mountains, and the Santa Clara River basin. Together, they encompass a spectrum of significant and unique habitat, vegetation and wildlife resources. The major habitat linkage zones and watersheds between the river basin and the Angeles National Forest, and the protected areas of the County (Placerita Canyon Natural Area), have also been included within the SEA boundary. Near the peak of Magic Mountain, the boundary contours to the southwest, and then proceeds west along the Santa Clara Divide to its intersection with the junction of Interstate 5 and State Route 14. Natural areas of the Sand Canyon watershed, along with the major topography of ridgelines, earthquake escarpments, grasslands, and canyon habitat features and watersheds of Bear, Placerita, Whitney, and Elsmere canyons are the important features of the wildlife linkage. Existing rural residential developments are excluded from the SEA, but the remaining natural highland areas of the western banks of the Sand Canyon watershed are included. These are integral parts of the river basin recharge system and functional ecosystem.~~

~~Parts of this area have coastal sage scrub and are critical habitat for the threatened coastal California gnatcatcher. The watershed of Placerita Canyon southeast of the State Route 14 is generally critical habitat for the federally threatened coastal California gnatcatcher. An area of development surrounding the Placerita Creek near State Route 14 is excluded from the critical habitat. The critical habitat area for the gnatcatcher extends along the east side of State Route 14 beyond Placerita Creek and envelops watersheds into the Angeles National Forest along Whitney Canyon, Elsmere Canyon, and southward over the main ridge of the San Gabriel Mountains, into Grapevine Canyon in its upper natural watershed. Upper areas of these canyons with oaks and big cone Douglas fir are habitat for the California spotted owl (*Strix occidentalis*).~~

~~The eastern half of the Los Piñetos undercrossing of State Route 14 on old oil development roads is included, and focuses on a major wildlife conduit connecting the Santa Susana Mountains to the San~~

DRAFT

Gabriel Mountains, and to the Santa Clara River. The adjacent part of the Santa Susana Mountains and Simi Hills SEA includes the west half of the Los Piñetos undercrossing of State Route 14, connecting through the natural oak woodlands and drainages adjacent to the San Fernando Pass. This area, once called "San Francisco" or "Newhall Wedge," is north and west of the junction of Interstate 5 and State Route 14 with The Old Road running through it. The Newhall Wedge area is nearly all critical habitat for the coastal California gnatcatcher. This critical habitat of the Newhall Wedge is adjacent to the gnatcatcher critical habitat across State Route 14 in the SEA, but is in the Santa Susana Mountains and Simi Hills SEA.

The SEA boundary borders State Route 14 from the north ridge of Grapevine Canyon and heads northeast from the Los Piñetos undercrossing, on the natural side of existing development east of State Route 14. The area around development along Running Horse Road off Placerita Canyon has been excluded from the SEA. The movie shoot ranch at the junction of State Route 14 and Placerita Canyon has much area with development or staging excluded, but there is a connected finger of the SEA in Placerita Canyon that leads to the Placerita Canyon watercourse underpass. Much of the watercourse underpass is used by wildlife to transition between the natural areas of Placerita Canyon and the oil field area on the west side of State Route 14. The SEA narrows to the western hills of Sand Canyon beyond the movie shoot ranch, to avoid developed areas, and continues back to the river margin at Humphreys railway stop, about a 0.4 mile west of its previous point of departure from the river channel. The boundary was drawn to avoid existing major development, but connect the uplands to the river basin. The narrow aperture for the linkage at the Santa Clara River reflects the remnant nature of the last unobstructed terrestrial passageway between the upland areas and the river.

West of Sand Canyon, the river has been intermittently armored to allow for development within flood hazard zones. From Sand Canyon westward through the residential neighborhoods of Santa Clarita, the SEA boundary continues on the margins of the flood plain to the confluence with San Francisquito Canyon. The segment of the Santa Clara River passing through the City of Santa Clarita is a dry channel, except during seasonal runoff flows. Some irregular extensions go north into tributaries that have remnant riparian habitat and probable outflows from irrigation runoff that flows into neighborhood storm drains. Regardless of the intermittent nature of water, the river bed elevated areas among braided channels support relatively intact stands of alluvial sage scrub, riparian woodland, and southern riparian scrub. The dry zones are essential to the continued genetic isolation and integrity of the unarmored three-spine stickleback population in the upper reaches of the Santa Clara River.

The boundary extends northward upstream into the reaches of San Francisquito Creek (formerly a separate SEA, but now included with the SEA), following the approved development setback limits, north into the Angeles National Forest (Santa Clara/Mojave Rivers District). The SEA continues nearly the length of the San Francisquito Creek to beyond the junction with South Portal Creek in the vicinity of the community of Green Valley. The Santa Clara River IBA extends in a branch upstream in close proximity to the crossing of Copper Hill Drive.

As the channel enters the Angeles National Forest, flows become less seasonal, and riparian resources expand and diversify. San Francisquito Creek supports dense and mature southern riparian scrub and riparian woodland formations, along with small areas of freshwater marsh, which provide essential wintering areas and resident habitat for waterfowl, wading birds, marshland birds, and a variety of other vertebrate species. The headwaters of San Francisquito Creek are on a low ridge that bounds the San Andreas Fault Zone, and this is an important connective element of the SEA, in that it completes the path from the Pacific Ocean through the mountains to the Mojave Desert. The sub-watershed and flood plain of the San Francisquito Creek perennial flow in the Angeles National Forest jurisdiction is designated critical habitat for the federally threatened red-legged frog, which extends

DRAFT

from about the Angeles National Forest southern boundary to about one mile south of the junction with Bee Canyon. Much of the San Francisquito Creek is considered essential habitat (one of three areas) for the endangered unarmored threespine stickleback, although the fish has not been found in the San Francisquito Canyon in recent years.

The boundaries west of the confluence with San Francisquito Creek follow the river margins under the Interstate 5 to the Castaic Creek confluence, at which point the northern setback line has been drawn around the lower portion of Castaic Creek, which embraces the riparian habitat areas around and above the confluence. Castaic Creek is the tributary with the largest watershed for the Santa Clara River in the County. The SEA boundaries go upstream about four miles along the watercourse of Castaic Creek to the crossing of Lake Hughes Road, which is just downstream of Castaic Lagoon. The Santa Clara River IBA extends in a branch upstream into Castaic Creek for approximately one mile.

Relatively extensive areas of willow cottonwood forest and southern riparian scrub occur west of San Francisquito Creek and within the junction zone of Castaic Creek and the Santa Clara River. These river forests support numerous sensitive species and provide multi-layered riparian habitat for a wide diversity of wildlife species, particularly birds of prey and riparian-obligate song birds, such as the federally endangered least Bell's vireo (*Vireo bellii pusillus*) and the southwestern willow flycatcher (*Empidonax traillii extimus*).

Federally designated critical habitat for the endangered arroyo toad extends from the east side of Interstate 5, from the junction of the Santa Clara River with San Francisquito Creek, under the Interstate 5, about 5.8 miles to the confluence, with an unnamed drainage just upstream of the confluence of the river with San Martinez Chiquito. The critical habitat area for the toad also includes the flood plain of Castaic Creek as far upstream as the Interstate 5 undercrossing (about 2.5 miles), and for about one mile upstream into the natural area of Hasley Canyon, a tributary of Castaic. Coincident with the critical habitat for the toad is critical habitat for the endangered least Bell's vireo (FE, SE). Critical habitat for the vireo extends along the floodplain from the Rye Canyon undercrossing of the river (west side of Interstate 5), over the Ventura Los Angeles County line, to about a mile short of the confluence of the Santa Clara River with Piru Creek in Ventura County (about 9 miles). The river area from near Interstate 5 towards the Ventura Los Angeles County line is "essential habitat" for the threespine stickleback. A disjunct SEA area is on a ridge south of the river bend at Castaic Junction (interchange of Interstate 5 and State Route 126). This area supports a population of the federal candidate and state endangered San Fernando Valley Spineflower (*Chorizanthe parryi* var. *fernandina*, FC, SE), which is a diminutive, once common flower of slopes within the San Fernando Valley and adjacent passes and mountain ranges. The plant became so rare that it was believed to be extinct until it was rediscovered during required surveys for development.

Beyond the confluence with Castaic Creek, the boundaries of the SEA follow the margins of the Santa Clara River channel to the Ventura Los Angeles County line. The Santa Clara River IBA has a lobelike expansion opposite the confluence with San Martin Chiquito, extending south to cover diverse topography from river cliffs to confluence flood plains in the area around Potrero Canyon.

The Santa Clara River channel and its alluvial terraces and tributary creeks together form the single most important and natural wildlife movement zone through the County. Mobile species can enter the river basin anywhere along its length (outside of developed areas) and proceed in either direction without having to pass through narrow culverts or blind channels, with continuous vegetative cover and only short stretches of dry substrates. The overall drainage course provides a continuum of aquatic and terrestrial movement opportunities, shelter, forage, and resident habitat from the mouth

DRAFT

of the river at Ventura County and the Pacific Ocean, to the Antelope Valley. The drainage course connects to both districts of the Angeles National Forest, and links together three large public resource preserves (Vasquez Rocks and Placerita County Natural Areas and the Angeles National Forest).

Wildlife Movement

Historically (and prehistorically) the riparian corridor along the Santa Clara River has served as the primary east-west linkage between the Pacific coastline, coast ranges, interior ranges, high desert and southern Sierra (via the Tehachapi Range). Animals moving through the Santa Clara drainage had unobstructed passage along the river and within the riparian systems between the coastal lowlands of Ventura County and the Mojave Desert. The tributary routes extend south into the Santa Susana Mountains, south and north into the San Gabriel Mountains, northward via Castaic, Bouquet and San Francisquito tributaries (over the coastal ranges and San Gabriel Mountains of the Transverse Ranges and into the San Joaquin Valley), west into the central coast ranges, or east through the Tehachapi Mountains, and into the southern Sierra Nevada. The present configuration of the tributary drainages has impinged upon connectivity from the Santa Clarita Valley to the north, but the Santa Clara River remains relatively intact and open. The SEA embraces the river corridor and the linkage zones that are considered essential to ensuring connectivity and resource values within the historic movement zones for all of the wildlife species present within the County portion of the Santa Clara River, including mountain lion, coyote, bobcat, and several medium-sized mammals, as well as birds, reptiles, amphibians, and fishes.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE SANTA CLARA RIVER SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	The only existing natural population of the federally endangered unarmored three spine stickleback is within the Santa Clara River and its tributaries, and all of its essential habitat is in this SEA. The federally threatened Santa Ana sucker occurs in the river, as does the state species of concern, the arroyo chub. The population of state and federally endangered slender horned spineflower in Bee Canyon is one of fewer than seven known occurrences for this species, one of only two known occurrences in the County, and one of its largest populations. San Francisquito Creek has a breeding area for the endangered red-legged frog. The San Fernando Valley spineflower (at Newhall Ranch in Interstate 5 vicinity) is found in only a few nearby places. Some of the critical habitat for the threatened California coastal gnatcatcher is included in this SEA. Western spadefoot, which is a species of concern, is extremely rare and local in the County away from this SEA. One of the largest, if not largest populations of least Bell's vireo in the County occurs along the river in the vicinity of the crossing of Interstate 5 near Newhall Ranch. Many RPR listed rare plants occur within the SEA. Critical habitat occurs in the SEA for the

DRAFT

	Criterion	Status	Justification
			listed arroyo toad, the red legged frog, the coastal California gnatcatcher, and the least Bell's vireo.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The low elevation bigcone Douglas fir canyon oak forests above Placerita Canyon, the vernal pool in the Placerita Canyon Sand Canyon divide, the native grassland on the Golden Valley Ranch (upper Placerita Canyon), and the alluvial fans with sage scrub in lower San Francisquito Canyon, Kentucky Springs and Acton are unique and regionally restricted biotic communities. Additionally, the riparian forests and woodlands along the Santa Clara River are among the most extensive, diverse and intact vegetative stands of this type in Southern California. Rare aquatic species, such as the unarmored three spined stickleback, Santa Ana sucker, red legged frog, least Bell's vireo, summer tanager, spineflower, and many others represented within the SEA are found nowhere else in the region.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The cottonwood willow forests and woodlands, alluvial fan sage scrub, and coast live oak riparian forest are best represented in the County within the SEA. The lower elevation examples of bigcone Douglas fir canyon oak forest communities where they mix with low elevation biota are restricted to the edges of mountain habitat communities, which are regionally rare and also designated in this SEA.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The Santa Clara River is simultaneously an oasis running through a dry landscape and an extension of coastal conditions into the dry interior. For this reason, it supports unique populations of aquatic and amphibious species, as well as aridlands species extending towards the coast and coastal species' extension inland. It is a principle migratory route for the County plants and animals and a center of diversity for the County. The Santa Clara River and its tributaries provide breeding opportunities for numerous species otherwise not known to breed within the County, including California red legged frog, summer tanager, southwestern willow flycatcher, and the unarmored three spined stickleback. The extensive riparian areas shelter dozens of migrant songbird species during winter, including high concentrations of white-crowned and golden-crowned sparrows, fox sparrow, yellow-rumped warbler, dark-eyed junco, and sharp-shinned hawk. The SEA embraces the river corridor and the linkage zones that are considered essential to ensuring connectivity and resource values for many of the wildlife species that are present within the County portion of the Santa Clara River.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The Santa Clara River represents a unique example of a drainage that stretches from the desert to the coast through the mountains. Its resources are, by definition, present at their geographic extremes. Plants such as western juniper, snake cholla, basin sagebrush, and birds, such as summer tanager are at the southwestern edges of their ranges along the river. Coastal taxa extend to the headwaters in the Acton area. High elevation

DRAFT

	Criterion	Status	Justification
			species, such as bigcone Douglas fir, spotted owl, and Steller's jay occur at fairly low elevations at the edges of Santa Clara River valley, on north facing slopes that remain cool all summer.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The SEA encompasses some of the highest quality, least disturbed and biotically intact acreage of bigcone Douglas fir canyon oak forest, riparian forest and woodland, coastal sage scrub, and alluvial fan sage scrub that remains in the County, and one of the three known vernal pools along the river. Vernal pools are rare everywhere in California.

In conclusion, the area is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) numerous examples of species at their habitat extremes as the coastal and desert influences meet; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Santa Felicia SEA

Boundary and Resources Description

The Santa Felicia SEA is located northwest of the City of Santa Clarita within unincorporated area of the County. Some of the SEA extends into the Angeles National Forest. The area is west of the Interstate 5, north of State Route 126 and encompasses almost the entire County portion of the Santa Felicia watershed that drains into Lake Piru and Piru Creek. Piru Creek has the largest watershed of any tributary of the Santa Clara River. The SEA is largely composed of natural coastal slopes of the western San Gabriel Mountains, with south facing slopes of coastal sage scrub and grasslands, north-facing slopes of oak woodland and chaparral, and canyons of riparian oak forest and other riparian habitats. This habitat has been diminished by development, and the SEA is one place in the County where the natural habitat remains.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Whitaker Peak and Val Verde.

Starting at the north end of the SEA boundary on the Ventura Los Angeles County line, the northern SEA boundary is about one quarter to one half mile north of the boundary of Angeles National Forest, with private property as it follows along the northern ridgeline of Devil Canyon. Most of the SEA captures the natural, coastal sage scrub covered south facing slopes, oak forests and mixed chaparral of the north facing slopes, and deep ravines with riparian oak forest in the canyons. Devil Canyon is a major tributary of Piru Creek (at Lake Piru) in Ventura County. Where the Devil Canyon watershed contacts the southern ridgeline off Townsend Peak, the SEA boundary turns south along a ridge that separates Santa Felicia Canyon and its tributaries on the west side, from Palomas Canyon on the east side. The boundary goes south out of the Angeles National Forest into private lands about a 0.75 mile south of Townsend Peak. To capture the watershed tributaries, the eastern boundary follows the dominant ridgeline between Palomas Canyon and Santa Felicia Canyon, and then between

DRAFT

~~Santa Felicia Canyon and Violin Canyon after Palomas Canyon joins Violin Canyon. There is a triple divide for the watersheds of Violin Canyon (which joins Castaic Creek), the watershed of Romero Canyon, which joins Hasley Canyon before it joins Castaic Canyon, and Santa Felicia Canyon, which is a tributary of Piru Creek. At the triple divide, the SEA boundary turns west to follow the ridge of the Santa Felicia watershed. The boundary crosses Loma Verde (peak), where it separates Santa Felicia Canyon from Romero Canyon, and the unnamed headwater creeks of Hasley Canyon, excluding the rapidly developing areas. About a mile southwest of Loma Verde Canyon, the SEA boundary turns south and west to encompass the watershed of Oak Canyon, with coast live oaks and riparian forest. The SEA boundary goes north along the Ventura Los Angeles County line, crossing riparian forest of Santa Felicia Canyon and Devil Canyon into the Angeles National Forest, where it joins the north ridge of Devil Canyon. Most of the SEA is included in the wildlife movement linkage Sierra Madre Castaic Connection that was outlined by the South Coast Wildlands study of linkages (Penrod, et al. 2005).~~

~~The SEA includes a wide variety of topographic features and habitat types. The orientation and extent of the SEA encompasses the surface and subsurface hydrology of the Santa Felicia watershed, from its headwater, tributaries, and basin to the point at which it exits the County jurisdiction.~~

~~The SEA encompasses most of the County portion of the Santa Felicia watershed that drains into Lake Piru. This watershed is largely undeveloped and contains vast stands of coastal sage scrub and chaparral communities on south and north-facing slopes. In addition to the undisturbed upland habitats, the watershed includes examples of mixed riparian (sycamore willow), oak riparian and coast live oak forests and alluvial scrub in the bottomlands. Grasslands occur in areas where grazing may have taken place; however, there is little invasion of these ruderal taxa into the native communities.~~

Wildlife Movement

~~The SEA provides riparian corridors, which serve as linkages between the Pacific coastline, coast ranges, interior ranges, the high desert and southern Sierras (via the Tehachapi Range). Animals move through the Santa Felicia watershed along and within the riparian systems between Piru Lake in Ventura County and the San Gabriel Mountain range and beyond. The tributary drainages in this SEA appear fully intact and open and support regional movement by many wildlife species. Most of the SEA was designated as an important wildlife movement linkage Sierra Madre Castaic Connection which was outlined by the South Coast Wildlands study of linkages (Penrod, et al. 2005).~~

Regional Biological Value

~~The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.~~

CRITERIA ANALYSIS OF THE SANTA FELICIA SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Not Met	The SEA does not include known core habitat.
	On a regional basis, biotic		The natural mountainous terrain and vegetation habitats of

DRAFT

	Criterion	Status	Justification
B)	communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	the SEA, which is a coastal drainage, have been diminished elsewhere within Southern California through development. The SEA encompasses a fine example of vegetation that has not been impacted very much by development.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	This is a somewhat remote area where natural flora and natural bottoms remain in the canyons, invasive predators are nonexistent, and human intrusion is minimal, which is a rare habitat in the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The SEA encompasses important tributaries (Santa Felicia Canyon and Devil Canyon) of Piru Creek. Piru Creek is the major tributary of the Santa Clara River. Nearly all of the drainages within the SEA are natural with sensitive habitats of all kinds of riparian forest. These drainages are all migratory corridors for both plants and animals that connect the San Gabriel Mountains with the Santa Clara River and the coast, the Sierra Madre-Castaic Connection.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Not Met	The Santa Felicia watershed is not known as a physical/geographical extreme habitat.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The Santa Felicia watershed is an excellent example of the inland watercourses and their vegetation on the coastal side of the Transverse Ranges. It is largely undisturbed, and a number of the drainages are still perennial.

In conclusion, the area is an SEA because it contains B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, and migrating grounds, which are limited in availability in the County; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Santa Monica Mountains SEA

Boundary and Resources Description

The Santa Monica Mountains SEA is located within the Santa Monica Mountains in a mostly unincorporated area of the County. Much of the area is in the Santa Monica Mountains National Recreation Area, but is privately owned. Many of the federal lands under the jurisdiction of the National Park Service (NPS) are included in the SEA designation. Many of the state parklands, notably Malibu Creek State Park and Topanga State Park, are also included in the SEA. The SEA includes nearly all of the canyons and ridges from the Ventura-Los Angeles Countyline, and east to Sullivan Canyon, which is near the communities of Pacific Palisades Brentwood to the south, and Encino to the north.

DRAFT

~~From south to north, the SEA extends from the Pacific Ocean shoreline or urban-wildland interface of Malibu, through the unincorporated area of the Santa Monica Mountains proper, to the northern edge of the SEA extending along the undeveloped southern edge of the San Fernando Valley or irregularly along the Ventura Los Angeles County line. It should be stated that this SEA recognizes the rare habitat of a small regional mountain range with a high diversity of topography and moisture regimes, and with vegetation adapted to a Mediterranean climate, which is globally rare, existing elsewhere only along western portions of continents at 30-40° latitude². Although the habitats may seem common within the Santa Monica Mountains, in terms of limited indigenous global ranges of the constituent species, their special adaptations to climate, the relatively intact character of the habitats, and the plant assemblage of the Santa Monica Mountains are unique. Development within the SEA that extends the nearby expansive urban development of the Los Angeles Basin and San Fernando Valley needs to be carefully considered to preserve these special resources.~~

~~The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Thousand Oaks, Calabasas, Canoga Park, Triunfo Pass, Point Dume, Malibu, and Topanga.~~

~~Within the SEA boundary, there are a number of areas that are not a part of the SEA due to dense development. For example, La Sierra, Malibu Lake, and most of the City of Malibu have been excluded.~~

~~The majority of the SEA lies within the unincorporated area of the County. Other jurisdictions include the City of Malibu, the City of Los Angeles, City of Calabasas, the City of Agoura Hills, the City of Hidden Hills, and the City of Westlake Village.~~

~~From the Ventura Los Angeles County line to the vicinity of Topanga Beach, the southern boundary of the SEA irregularly contacts the boundary of the Malibu Coastline CRA, which extends offshore about a 0.8 mile to capture the kelp beds and rich natural marine communities of and off the coastline of the Santa Monica Mountains. The sandy beaches along this SEA are the least disturbed beaches of the County, some of them completely natural. This is a remnant of the typical rock and sand shoreline that once occurred along the coast of Southern California. From Mugu Lagoon to Latigo Point (the County portion starting at the Ventura Los Angeles County line and Leo Carrillo State Beach), this is an ASBS (Area of Special Biological Significance), which is a marine area designated by the State Water Resources Control Board as having exceptionally good water quality and natural community features. Populated and disturbed areas along the shore are not included in the SEA.~~

~~Beginning at the intersection of the Ventura Los Angeles County line and State Route 1, the southern area of the SEA includes all the terrestrial area of Leo Carrillo State Beach (a state park that has much beautiful upland acreage of grasslands, coastal chaparral, and an excellent riparian area of sycamore-oak forest along the Arroyo Sequit). The Arroyo Sequit from the coast to the junction of West and East forks and about a 0.5 mile of the West Fork are federally designated critical habitat for the southern steelhead (*Oncorhynchus mykiss irideus*), which is a salmon that runs upstream for breeding, but~~

² Six regions on Earth have a coastal climate where ocean currents mediate an equable climate yet cause most precipitation to occur outside the summer period of most intense sunlight and most vegetative growth for many plants. These are California, the Mediterranean Basin, portions of South Australia, South Africa, and Chile

DRAFT

spends most of its life in the ocean. The young fish, however, spend the first year of life in fresh water.

~~At the eastern boundary of Leo Carillo State Beach, an unnamed canyon is excluded from the SEA, but San Nicolas Canyon with its riparian woodland is included all the way to the shore. Urban areas to the east of San Nicolas Canyon are also excluded. The Los Alisos and Lechuza Canyon immediate drainages are included up to the highway. From Lechuza Canyon, the excluded area extends north to the southern border of Charmlee Natural Area (a County park supporting grasslands and chaparral). Much of the Encinal Canyon drainage is included in the SEA north of State Route 1, and the Steep Hill Canyon drainage is included north of State Route 1. Trancas Canyon is densely settled in the area near the coastline; only the steeper, natural section is included in the SEA. To the east, a large excluded area encompasses the dense habitation of the Malibu Riviera.~~

~~The SEA includes the riparian forest of Zuma Canyon as it exits onto the coastal plain, with a small sub-watershed designated critical habitat for the federally endangered Braunton's milkvetch (*Astragalus brauntonii*). This rare plant is narrowly endemic to four mountain ranges of Southern California that have a particular marine-originated rock stratum of interbedded carbonates and sandstones. The SEA also includes portions of the riparian forest of Ramirez Canyon, where it has not been impacted by development in the Malibu Riviera. The natural area of Escondido Canyon and its 150-foot waterfall is included. Latigo Canyon is included north of State Route 1, where chaparral intergrades with walnut and oak woodlands along the drainages and north-facing slopes. East of Latigo Canyon, the boundary of the excluded area follows the ridges immediately north of State Route 1 to join the highway west of Solstice Canyon. Solstice Canyon and its County Park are included in the SEA. These support grasslands, chaparral, and a fine riparian area of sycamore-oak forest along Solstice Canyon. Puerco and Marie Canyon drainages are included north of State Route 1.~~

~~In the vicinity of Winter Canyon, the boundary follows Malibu Canyon Road to exclude the central community area of Malibu in another large island. A finger of the SEA extends through central Malibu along Malibu Creek in the area of Malibu Creek State Park and Malibu Lagoon State Beach to include the Creek and Malibu Lagoon south of Pacific Coast Highway. This is one of the points of contact between the Santa Monica Mountains SEA and the Malibu Coastline CRA. Malibu Creek from the shore to its major bend to the west in Malibu Canyon is federally designated critical habitat for the southern steelhead. Coincident with the steelhead critical habitat along Malibu Creek and all brackish portions of the Malibu Lagoon are designated critical habitat for the endangered tidewater goby (*Eucyclogobius newberryi*). This little two-inch fish was once common in brackish water of coastal lagoons of streams the length of the State of California. This pristine habitat is disappearing (the fish is apparently gone from San Francisco Bay), and any location where the tidewater goby is still found is a unique place.~~

~~The Malibu Lagoon is the only remaining intact coastal lagoon in the County³, and still has its perennial stream to maintain the brackish conditions that are essential to the diverse and unique group of species of coastal lagoons. The federally endangered southern steelhead passes through this lagoon on its upstream and downstream spawning runs, and its young fish (smolts) may spend some time in the lagoon before entering their oceanic habitat. The passage is critical habitat for the steelhead, as well as for the endangered tidewater goby (*Eucyclogobius newberryi*). This lagoon has one of the~~

³ Ballona Creek lagoon is also in Los Angeles County, but exists in a disturbed condition.

DRAFT

County's three ⁴ remaining saltmarshes, with dominant species of pickleweeds (*Arthrocnemum subterminale* and *Salicornia pacifica*). The Malibu Lagoon supports a good representation of coastal strand community (now groomed off many of the recreational beaches), is a resting stop on the Pacific Flyway (over 200 species of birds reported), and has many resident avian species. From fall to spring, it supports a flock of the federally-threatened western snowy plover (*Charadrius alexandrinus nivosus*), and there are recent indications that some may over-summer to breed (after a hiatus in breeding of many decades, Ryan Ecological Consulting, 2010). The pickleweed is valuable non-breeding habitat for the state-endangered Belding's savannah sparrow (*Passerculus sandwichensis beldingi*).

East of central Malibu, the excluded area borders primarily follow the less-inhabited ridges that border Carbon and Las Flores canyons and about a 0.5 mile north of State Route 1. East of Las Flores Canyon, the border of the excluded area follows a transmission line to skirt an old oil field. At the western ridge of Piedra Gorda, the SEA boundary follows the ridgeline southeast to State Route 1. From Piedra Gorda, the SEA boundary closely tracks State Route 1 along the substantial sea cliffs, avoiding two small developed areas north of the highway near the mouths of Pena and Tuna canyons. The SEA includes all of the meandering Topanga Canyon Creek at Topanga Beach. Upstream nearly to the community of Fernwood, Topanga Canyon is federally-designated critical habitat for the southern steelhead. Near the shore in Topanga Canyon Creek is critical habitat for the tidewater goby. This is the eastern point of contact between the Malibu Coastline CRA and SEA. It is also at this point where the shoreline where there are no more excluded areas.

The SEA includes Parker Mesa, north of State Route 1, and is coterminous with the boundary for the City of Los Angeles towards the north. About one mile north of the coast, the boundary extends to the east, contouring around the ridgelines that border the populated areas of Santa Ynez, Pulga, and Temescal canyons, with some residential areas on the ridges excluded from the SEA. These three canyons contain riparian oak and sycamore forests, which are a good variety of canyon habitats that include rocky outcrops and small waterfalls, are preserved in their upper reaches by inclusion in the Topanga State Park. The upstream areas of Santa Ynez, Pulga, and Temescal canyons that are uphill from the community of Palisades Highlands include much of federally-designated critical habitat for the Braunton's milkvetch. The boundary skirts Will Rogers State Park, but includes much of the unpopulated part of Rustic Canyon.

At Rustic Canyon, the boundary follows the eastern ridge to the north. Tracing the edge of development along the eastern slope of the canyon, the boundary continues north and eventually crosses over the ridgeline and into the undeveloped part of Sullivan Canyon. The boundary then follows the eastern slope of Sullivan Canyon and continues north along the ridgeline between Sullivan Canyon and Mandeville Canyon. After crossing the main ridge of the coastal zone and Mulholland Drive near San Vicente Mountain, the SEA boundary reaches and includes the sub-watershed of Encino Reservoir.

On the west side of Encino Reservoir, the SEA boundary turns west and follows the edge of development on the northern slopes of the Santa Monica Mountains that form the southern border of the San Fernando Valley. Some development is excluded on the east side of Topanga Canyon and the Calabasas Highlands area. The upper reaches of Topanga Canyon are included, and the SEA crosses the ridge to include the upper slopes of a natural drainage that is opposite to Topanga Canyon

⁴ The three areas with salt marsh in Los Angeles County are Alamitos Bay, Ballona Lagoon, and Malibu Lagoon.

on the San Fernando Valley side.

West of Calabasas Park, the SEA boundary extends northeast and contours along the upper slopes of McCoy Canyon to include a finger of SEA on the north ridge of McCoy Canyon, which is part of state park land. On the west side of Calabasas Park and Hidden Hills, the boundary crosses State Route 101 along the western edge of development in Hidden Hills north to the County line. The undeveloped portion of Gates Canyon within the Simi Hills and its watershed is included north to the Ventura-Los Angeles County line, excluding a ridgetop island and developed portions of the City of Calabasas. A narrow finger of connection joins the portion of the SEA east of the City of Calabasas to the area west of the City, north of State Route 101 along a tributary of Las Virgenes Creek. Las Virgenes is an important tributary of Malibu Creek, and although this area is not in the coastal zone, it represents a portion of the upper coastal watershed. Further to the west, the boundary extends north again to encompass the open space from Las Virgenes Road along the Ventura-Los Angeles County line to the edge of development within the City of Agoura Hills. Within this portion of the SEA, Chesebore Canyon contains fine natural, undeveloped sections of canyon just east of the City of Agoura Hills that are administered by both the NPS and the State Park system. The SEA includes all of undeveloped Chesebore Canyon and the undeveloped western ridge of Chesebore's important tributary, Palo Comado Canyon. This is the western terminus of the SEA that lies north of State Route 101.

An important feature of this area is a small tributary of Las Virgenes Creek (and ultimately Malibu Creek) named Liberty Canyon. The underpass of State Route 101 at Liberty Canyon Road along the drainage conveys relatively less vehicular traffic than other freeway crossings within several miles, and is one of the few active wildlife passage areas along the entire extent of State Route 101 through the Santa Monica Mountains. All other watercourse and street crossings of State Route 101 are very constrained or else entirely impassible for wildlife. There are widely documented concerns for the consequences of genetic isolation for the small band of mountain lions (*Puma concolor*) of the Santa Monica Mountains. Mountain lions that are known to use this area of the SEA to transit back and forth between the Santa Monica Mountains and the greater lion populations of the Simi Hills and north. The area north of the Liberty Canyon underpass is natural chaparral, watercourses, and grassland administered by the Santa Monica Mountains Conservancy. South of State Route 101, development is present with a very narrow corridor of natural habitat that includes valley oak (*Quercus lobata*) along the creek of Liberty Canyon.

A small island of developed area south of State Route 101 and along Liberty Canyon is excluded from the SEA. From the Liberty Canyon underpass south of State Route 101 the SEA boundary follows the north-facing side of an unnamed ridge at the southern end of the City of Agoura Hills towards the northern slopes of Ladyface Mountain. This is a chaparral-dominated area with numerous canyons and ridges supporting federally-threatened Agoura Hills dudleya (*Dudleya cymosa* ssp. *agourensis*), and one of the nine critical habitat areas in the County of Los Angeles and in the SEA for the state and federally-endangered Lyon's pentachaeta (*Pentachaeta lyonii*), as well as other uncommon and rare plants, such as Fish's milkwort (*Polygala cornuta* var. *fishiae*), Sonoran maiden fern (*Thelypteris puberula* var. *sonorensis*), and the southern-most known occurrence of the California juniper (*Juniperus californica*) within the County. The SEA northern boundary crosses Kanan Dume Road in the area where Lindero Canyon formerly joined Medea Creek and continues its contour along the north-facing ridge of Ladyface Mountain, which also supports Agoura Hills dudleya, Lyon's pentachaeta, Ojai navarretia (*Navarretia ojaiensis*), and other rare and sensitive plants on its chaparral-covered slopes. A small area with development and fill along the south side of Kanan Dume Road has been excluded from the SEA.

DRAFT

~~The SEA boundary veers to the southwest to avoid the developed areas of Westlake Village, but includes Las Virgenes Reservoir in the SEA and three more of the critical habitat areas for the Lyon's pentachaeta. A finger includes an undeveloped section of Triunfo Canyon. A short distance southwest of Las Virgenes Reservoir, the boundary crosses Decker Canyon road and extends west to the Ventura-Los Angeles County line, which is another area of critical habitat for the Lyon's pentachaeta. Most of the critical habitat areas for Lyon's pentachaeta in the County are covered by the North Area Plan regulations for the Santa Monica Mountains, but some critical habitat areas also occur in the coastal zone. The boundary then follows the Ventura-Los Angeles County line to the southwest all the way to the coast and State Route 1 where the northern and the southern boundary join at Leo Carillo State Beach.~~

Wildlife Movement

~~The SEA contains major features that are important to wildlife movement. For the mammals and other large terrestrial wildlife, it is important to recognize the crossing area of State Route 101 freeway at Liberty Canyon, and this has been detailed in the General Boundary and Resources Description section. Future highway modifications in this far western area of the County may take the need for wildlife connectivity into account. On the eastern end of the mountains (separated from the SEA by the communities of Brentwood and Encino in the City of Los Angeles) the California Department of Transportation, or Caltrans, has undertaken widening conversion of one of the bridges to incorporate a vegetated area on the bridge that will encourage wildlife transit between the mountain sections west and east of Interstate 405. This is the first such endeavor in County.~~

~~The many natural drainages and ridgelines of the Santa Monica Mountains connect populations in a netlike web, and development should always take account of the importance of these natural areas to wildlife connectivity.~~

~~The major coastal drainages with natural habitat shading and cooling the waters support a very unique and rare wildlife movement, which is the spawning runs of the federally endangered southern steelhead. Spawning run areas and young fish habitat are in critical habitat areas of Arroyo Sequit, Malibu, and Topanga Canyons. Historically, steelhead were known from Solstice and Zuma canyons as well, and it is likely that all the major drainages that once had perennial water and extended to the shore in the rainy season supported this species.~~

~~Although wildlife movement is hampered by rural development in the SEA, animals are still able to move through the Santa Monica Mountains in many areas. Due to its large size and topographic complexity, linkages in many directions occur within the SEA. However, there are also various bottlenecks. Edelman (1990) identifies Malibu Creek State Park as the central core habitat area in the Santa Monica Mountains, serving as a connective hub between the Simi Hills to the north and the open space preserves of Topanga State Park to the east, and Mugu State Park to the west (which is near the Ventura-Los Angeles County line, but within Ventura County). These linkages allow movement among large open space areas within the SEA as well as between areas outside the SEA, such as the Simi Hills and the western extent of the Santa Monica Mountains in Ventura County. The genetic flow through these areas is crucial in maintaining the diversity and viability of the species within the Santa Monica Mountains. Open space linkages between Kanan Road and Calabasas Parkway along State Route 101, as indicated by the National Park Service, are of particular importance for continued connectivity of wildlife populations, due to a lack of alternative routes and encroachment of development (Nelson, 2000). Although there are significantly large open spaces within the SEA, maintaining habitat linkages between them is critical in providing for long term sustainability. A wide~~

DRAFT

variety of wildlife use linkages throughout the SEA, including mountain lion, coyote, mule deer, bobcat, and a number of medium-sized mammals.

Regional Biological Value

The SEA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE SANTA MONICA MOUNTAINS SEA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	<p>The SEA provides habitat for the following listed species: Lyon's pentachaeta (FE, SE); beach spectaclepod (ST); Blochman's dudleya (FT); marcescent dudleya (FT); Santa Monica dudleya (FT); Braunton's milk vetch (FE); salt marsh bird's beak (FE, SE); tidewater goby (FE); steelhead – Southern California ESU (FE); western snowy plover (FT); western yellow-billed cuckoo (SE); southwestern willow flycatcher (FE, SE); American peregrine falcon (SE); bald eagle (FT); bank swallow (ST); California least tern (FE, SE); least Bell's vireo (FE, SE).</p> <p>The SEA contains designated critical habitat for Lyon's pentachaeta, Braunton's milk vetch, tidewater goby, steelhead, and western snowy plover.</p>
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Upper La Sierra Canyon contains an unusually rich and diverse stand of canyon flora, including marcescent dudleya, creek dogwood, and many unusually large specimens of other rare plant species; Malibu Lagoon is the only intact natural lagoon between Point Mugu in Ventura County and Anaheim Bay in Orange County; Malibu Canyon contains a unique mix of floral species that are uncommon in the region, such as black cottonwood and leather leaf ash, as well as a regionally unique mixture of inland and coastal species; regionally rare volcanic rock formations create unique communities where they occur.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Malibu Lagoon is the only natural lagoon in the County; upper La Sierra Canyon contains an unusually rich and diverse stand of canyon flora, including marcescent dudleya, creek dogwood, and many unusually large specimens of other rare plant species; and Malibu Canyon contains a regionally unique mix of floral species that are uncommon in the County, such as black cottonwood and leather leaf ash, as well as a unique mix of inland and coastal species.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or	Met	The Malibu Lagoon and the upstream riparian woodland in Malibu Creek is an important migrating bird refuge, with over 200 species recorded. Tuna and Pena canyons are an important area to migratory birds due to their combined qualities of healthy vegetation, riparian woodland, surface moisture, undeveloped land, and an unobstructed opening to the coast. The SEA also contains habitat linkages between large open space areas within the SEA and areas outside

DRAFT

	Criterion	Status	Justification
	in the County.		the SEA, such as the Simi Hills and the western extent of the Santa Monica Mountains in Ventura County. Such linkages are crucial in maintaining regional plant and animal population health and viability.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The SEA includes a myriad of unique and pristine natural areas that are important for nature study and scientific research; the range of extremes of many species, such as the California juniper, linear-leaved goldenbush, splendid mariposa lily (<i>Calochortus venustus</i>), and valley oak; and disjunct and unique populations of redshank, island mountain mahogany, lyre snake, mountain quail, hirsute rain-beetle, and the Jerusalem cricket.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Zuma Canyon is one of the last major drainages in the Santa Monica Mountains with a year-round stream that supports a rich riparian community and remains in an undeveloped state; Cold Creek includes an excellent example of an undisturbed natural sandstone basin with springs and a perennial stream; Tuna and Pena canyons are the last drainages in the central and eastern Santa Monica Mountains that have no development between the canyon mouth at the coast and upper areas of the watershed; Palo Comado and Chesebore canyons support one of the last examples of an oak woodland savannah of any significant size in the County; Temescal, Rustic, and Sullivan canyons represent contiguous, self-contained watersheds that are large enough to support representative samples of native flora and fauna; the area surrounding Encino Reservoir supports the best undisturbed stand of an inland chaparral, coastal sage scrub, and streamside vegetation remaining on the inland slope of the Santa Monica Mountains.

In conclusion, the area is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Santa Susana Mountains and Simi Hills SEA

Boundary and Resources Description

The Santa Susana Mountains and Simi Hills SEA is located northwest of the San Fernando Valley within unincorporated areas of the County and the City of Los Angeles, west of Chatsworth. The area is south of State Route 126 and the Santa Clara River, west of the Interstate 5, and includes much of the Santa Susana Mountains to the north, the Santa Susana Pass, Chatsworth Reservoir, and the eastern portion of the Simi Hills to the south. This SEA encompasses much of the natural area of the Santa Susana Mountains in the County. The north face of the Santa Susana Mountains is the southwestern watershed of the Santa Clara River in the County, and on the south face, the Santa

DRAFT

~~Susana Mountains are part of the direct coastal watershed as well as part of the watershed of the Los Angeles River. The Simi Hills are part of the direct coastal drainage in their southern area. The variations in vegetation communities are extensive. The area in the Santa Susana Mountains covered by the SEA is considered an important connective wildlife corridor among the San Gabriel Mountains, the Santa Clara River, and the Santa Monica Mountains.~~

~~The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Calabasas, Canoga Park, Oat Mountain, Santa Susana, Val Verde, and Newhall.~~

~~The entire western boundary of the SEA follows the Ventura Los Angeles County line from El Escorpion Park, west of the intersection of Valley Circle Boulevard and Vanowen Street in the West Hills community of the San Fernando Valley, and north to an area just south of the Santa Clara River near Salt Canyon Road. El Escorpion Park is adjacent to state park land just across the Ventura Los Angeles County line. The small ridgeline where the SEA begins just south of El Escorpion Park separates the coastal drainage of Las Virgenes (tributary of Malibu Creek) from drainages that flow into the San Fernando Valley and the Los Angeles River. El Escorpion Park is at the southern end of Bell Canyon Park, which is also on Ventura Los Angeles County line. The watercourse of Bell Canyon flows through the park. The SEA continues north, including the natural watershed of Dayton Canyon. Here on the southern side of Dayton Canyon is designated critical habitat for the Braunton's milk vetch (*Astragalus brauntonii*), which is alocoweed that prefers a substrate of intermixed sandstone and carbonate beds, which is probably deposited at the margins of a former seacoast. The SEA circles around the community of Lakeside Park, which is excluded from the SEA, and then encircles and includes the Chatsworth Reservoir along boundaries of private property. The western boundary of the SEA follows the Ventura Los Angeles County line, but bends to exclude development in upper Woolsey Canyon and Chatsworth Lake Manor.~~

~~From Chatsworth Reservoir, the SEA continues north with the west side along the Ventura Los Angeles County line and the east side tracing natural habitat at the edge of the Simi Hills and the San Fernando Valley. The SEA extends eastward to include all of the Santa Susana Pass area, much of which is preserved in the Santa Susana Pass State Historic Park. Just across the Ventura Los Angeles County line near State Route 118 in Ventura County are Corriganville Park, a former and current natural movie production area, and Rocky Peak Park. Corriganville is a regional park of the City of Simi Hills; Rocky Peak Park is administered by the Santa Monica Mountains Conservancy. Rocky Peak Park is the former Runkle Ranch and stretches from State Route 118 five miles northward to Las Lajas Canyon. The Park is a vital wildlife habitat linkage between the Simi Hills and the Santa Susana Mountains. Spectacular sandstone boulders, outcroppings, oak savannahs, and perennial water sources provide diverse habitat for vertebrates and a number of rare plants. The Ventura Los Angeles County line and the SEA boundary cross directly over Rocky Peak here.~~

~~A very important wildlife passage between the Santa Susana Mountains and the Simi Hills is just west of the Ventura Los Angeles County line connecting Corriganville Park and the Runkle Ranch. It is a broad tunnel under the freeway, which enabled the connection of the property that was divided by constructing State Route 118. This tunnel connects dirt roads and trails on either side, and is regularly used by mountain lions and other wildlife.~~

~~Most of the SEA from State Route 118 northward is designated critical habitat for the coastal California gnatcatcher (*Polioptila californica californica*), which is a diminutive bird that is becoming rare due to loss of its preferred habitat, which is coastal sage scrub. The critical habitat for this bird extends across the Ventura Los Angeles County line along the extent of the Santa Susana Mountains into Ventura~~

County.

~~North of State Route 118, the SEA excludes development in the area of Hialeah Springs (but includes the springs), and circles round the development to include the more sparse settlement in the Deer Lake Highlands area. The SEA boundary goes north along the watercourse of Browns Canyon to the confluence with Mormon Canyon. Here the SEA boundary climbs the ridgeline that separate Browns Canyon and Mormon Canyon to include Browns Canyon and the Michael D. Antonovich Regional Park at Joughin Ranch, and exclude the Mormon Canyon. Continuing up the ridge to its origin on the crest of Oat Mountain, the SEA boundary turns eastward along the ridgeline, excluding the oil fields in the upper reaches of Mormon Canyon and including the extensive natural areas of the north slope of the Santa Susana Mountains.~~

~~Along the Ventura-Los Angeles County line north of Rocky Peak, the SEA boundary crosses Blind Canyon (draining to San Fernando Valley), then Lajas Canyon (draining to Simi Valley and ultimately Mugu Lagoon), then El Toro and Chivo canyons (also draining to Mugu Lagoon). Northwest of Chivo, the crest of the Santa Susana Mountains is crossed, and drainages are tributaries of Salt Canyon and the Santa Clara River. The north edge of the critical habitat for the coastal California gnatcatcher is crossed near the crest, where the south-facing slopes that favor coastal sage scrub give way to the ridgeline and north-facing slopes that promote denser chaparral and oak woodlands. At the northern boundary of the SEA, the Santa Clara River SEA is contiguous as is the critical habitat for the state and federally endangered least Bell's vireo (*Vireo bellii pusillus*), which is a small bird that usually nests next to perennial water.~~

~~From the northwesterly corner, the boundary travels east along the north side of Salt Canyon, and then along the northern side of the Salt Canyon East Fork. Where the East Fork turns south, the SEA boundary continues east to encompass the steeper southern areas along Potrero Canyon and all of the Pico Canyon drainage south of Potrero. The SEA boundary is truncated at the Stevenson Ranch development, including the Wickham Canyon tributary of Pico in the SEA, but excluding most of Dewitt Canyon. This boundary of the SEA is essentially following the northern edge of the Salt Creek open space that was approved with the Newhall Ranch Specific Plan. In the vicinity of Pico Canyon, the boundary continues eastward to encompass the Lyon Canyon watershed and an unnamed watershed just north of Lyon up to the west side of Interstate 5. Along Interstate 5, the SEA boundary continues along the line of natural vegetation (west and southwest side of Gavin Canyon) including the watersheds of tributaries Towsley, Wiley, Leaming, Rice, and East canyons. The boundary continues east along the western edge of Interstate 5 to an area just west of the City of Los Angeles line, near the interchange with State Route 14. Here the boundary excludes the drainage of Sunshine Canyon, which is involved in the Sunshine Canyon Landfill used by both the City of Los Angeles and the County. Critical habitat for the coastal California gnatcatcher is in the watersheds of Towsley, Wiley, Rice, and East canyons south of the Interstate 5 below about the 2400 foot elevation contour.~~

~~North across the Interstate 5 is the Newhall Wedge. The Newhall Wedge is a very rugged part of the Santa Susana Mountains, with substantial natural vegetation of oak woodlands, chaparral, and coastal sage scrub. The Newhall Wedge is truncated by the flood plains of tributaries of the South Fork of the Santa Clara River to the north, east and west, and these flood plains have had extensive development as the City of Santa Clarita (Gavin Canyon is one of these tributaries.). Important connective areas are the road crossings of the Interstate 5 and State Route 14. The connection to the Santa Clara River SEA is the Los Pinetos Road underpass of the State Route 14. Consistent wildlife movement has been recorded with motion activated cameras there. The Weldon Canyon Road overpass of the Interstate 5 is another connection for the Newhall Wedge with the main part of the SEA. The Old Road underpass of the Interstate 5 is a broad connection. The Calgrove underpass is another broad~~

DRAFT

connection, but busy with traffic. Natural areas are adjacent to all these under- and overpasses. The South Fork of the Santa Clara River is formed by the junction of Towsley, Wiley and East canyons in the northeast corner of Michael D. Antonovich Open Space. Its underpass of Interstate 5 has a natural bottom that is used frequently by wildlife, but on the east side of Interstate 5 there is a series of 15 foot drops and channeled sides, which make it unlikely that terrestrially tied wildlife would continue into the populated parts of the City of Santa Clarita along the South Fork. Critical habitat for the coastal California gnatcatcher is in most of the Newhall Wedge part of the SEA between the Interstate 5 and the Sierra Highway that is just west of State Route 14.

On the west side of Sunshine Canyon, a broad lobe of the SEA extends along the ridgeline, which separates Sunshine and Bee Canyon to include Bee Canyon Park and Mission Point of O'Melveny Park. This is an area with walnut woodlands, oak woodlands, grasslands, and chaparral including coastal sage scrub, which is the diverse and green vegetation typical of the Santa Susana Mountains. Critical habitat for the coastal California gnatcatcher covers O'Melveny Park, except for the ridgeline of Mission Point, which is above the 2400 foot contour. The SEA boundary travels west from Mission Point along the ridgeline above the Aliso Canyon Oil Field and turns south at the western edge of the Aliso Canyon Oil Field, along the ridgeline between Mormon and Browns canyons. Critical habitat for the coastal California gnatcatcher extends below about the 2400 foot contour (including Mormon and Browns canyons) and roughly is within the SEA north of State Route 118.

The SEA includes a variety of topographic features; the northern portion of the SEA encompasses Oat Mountain and much of the Santa Susana Mountains from the Ventura-Los Angeles County line east to Interstate 5. Portions of many of the canyons associated with the Santa Susana Mountains and Oat Mountain are also included, such as Salt Canyon, Potrero Canyon, Pico Canyon, Towsley Canyon, El Toro Canyon, Sulphur Canyon, Devil Canyon, Ybarra Canyon, Browns Canyon, Bee Canyon, and Mormon Canyon. Several perennial stream areas occur within these canyons, and there are many natural springs. The north slopes of the Santa Susana Mountains are within the Santa Clara River watershed, which drains the Los Padres National Forest to the north, the Angeles National Forest to the northeast and east, and the Santa Susana Mountains to the south and southeast. The remainder of the SEA is within the Los Angeles River watershed. The majority of the land in the SEA is natural open space with very sparse disturbances in the form of ranches, oil wells, and unimproved access roads. The SEA consists of east-west and northwest trending primary ridges and north-south trending secondary ridges.

The peak of Oat Mountain represents the highest point in the SEA at 3,747 feet above mean sea level (MSL). From Oat Mountain, one can appreciate the diverse influences that create extremely diverse habitat within this SEA. One can see downstream along the Santa Clara River to the mouth of the Pacific Ocean and to the northern Channel Islands. Across the San Fernando Valley are the Santa Monica Mountains, and the Simi Hills enclose the west end of the Valley. To the east are the ascending ridges of the San Gabriel Mountains, and to the northeast the Santa Clara River continues upstream towards the Antelope Valley and the Mojave Desert. Coastal, valley, montane, and desert influences all meet within this small mountain range.

Open space within the SEA supports this great variety of communities, but is dominated by chaparral, oak woodlands, coastal sage scrub, bigcone Douglas-fir-canyon oak woodland, and grasslands; however, there are numerous examples of special vegetation. Not uncommon are cherry woodlands, which are dominated by hollyleaf cherry (*Prunus ilicifolia*). These mountains are a meeting area of the (regular) Douglas fir (*Pseudotsuga menziesii*) and the bigcone Douglas fir (*Pseudotsuga macrocarpa*). There are a number of special endemic plants, such as the Santa Susana tarplant (*Deinandra minthornii*), which is a tarplant like few others since it is perennial. Its distribution spreads through the

DRAFT

Simi Hills and into the Santa Monica Mountains, but it is primarily at home among the sandstone boulders and terraces, which prevail in the Santa Susana Mountains. The Santa Susana Mountains are the only known place in the County with individuals of the uncommon Palmer's oak (*Quercus palmeri*). This desert oak can be very long lived. A clone found in Riverside County was judged to have started from an acorn in the last Ice Age, over 10,000 years ago. Other oaks with groves in the Santa Susanas include coast live oak (*Quercus agrifolia*), valley oak (*Q. lobata*), canyon live oak (*Q. chrysolepis*), scrub oak (*Quercus berberidifolia*), and interior live oak (*Q. wislizenii*). The numerous creeks and canyons support riparian scrub and woodland communities with oaks, sycamores, and willows. There are walnut woodlands of the California black walnut (*Juglans californica*) mixed with flowering ash (*Fraxinus dipetala*) and western blue elderberry (*Sambucus nigra ssp. caerulea*) and coast live oak. Flowering ash may be a tree up to 60 feet tall in the Santa Susana Mountains, whereas it usually is a low tree or even spindly shrub. The woodlands dominated by walnuts and flowering ash appear to be unique to the Santa Susana Mountains. The bigcone Douglas fir canyon live oak forest at higher elevations represents one of the northwesternmost examples of this community. At its southern end, the SEA includes the eastern portion of the Simi Hills, including the east-facing slopes descending from Chatsworth Peak. Chatsworth Reservoir forms a portion of the south boundary and is currently dry, except for a small detention basin north of the reservoir.

Chatsworth Reservoir is now dry, having been taken out of service in the early 1970s due to seismic safety concerns. However, it retains a variety of interesting habitats, with avian communities of songbirds and waterfowl that benefit from the protection provided by fencing that still surrounds the reservoir. Hence, it is valuable for bird study by students, researchers, and naturalists. North of the reservoir proper and south of Valley Circle Boulevard is a pond and associated freshwater marsh, which supports numerous kinds of waterfowl during the spring and fall migration periods. The periphery of the reservoir is grassland and savannah, with a mixture of valley and coast live oaks (*Quercus lobata* and *Q. agrifolia*), some in small stands. The surrounding grassland and savannah are also important forage areas for migrating birds that roost in or near the wetlands, as well as year-round forage for resident species.

The majority of the SEA is within the unincorporated area of the County.

Wildlife Movement

The SEA includes several important linkages for wildlife movement. The Simi Hills and Santa Susana Mountains provide a vast open space corridor to foster wildlife movement between the Santa Monica Mountains to the south, San Gabriel Mountains to the east, and Los Padres National Forest to the north in the western San Gabriel Mountains of the Transverse Ranges. Dense, natural habitat associated with the majority of the SEA provides excellent opportunities for concealment and water sources, while the grasslands provide an abundance of prey. Examples of wildlife that use these linkages include mountain lion (*Puma concolor*), mule deer (*Odocoileus hemionus*), coyote (*Canis latrans*), bobcat (*Lynx rufus*), and a number of medium-sized animals.

Regional Biological Value

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met described below.

CRITERIA ANALYSIS OF THE SANTA SUSANA MOUNTAINS AND SIMI HILLS SEA

DRAFT

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	Most of the SEA has critical habitat for the coastal California gnatcatcher. A population of the Braunter's milkvetch has critical habitat in the Simi Hills part of the SEA. The SEA has robust populations of rare plants, such as the Plummer's mariposa lily and the Santa Susana tarplant.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The SEA contains habitat of the extremely rare Santa Susana tarplant. In addition, several plant communities within the SEA are CDFW highest inventory priority communities due to their restricted distribution in the Southern California region. These communities include coastal sage scrub, alluvial scrub, valley oak woodland, valley oak savannah, mainland cherry woodland, native grassland, southern willow scrub, and cottonwood willow riparian forest.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	All of the plant communities and habitats mentioned above as being restricted in distribution on a regional basis are further restricted in distribution within the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The open space of the SEA allows for connectivity between the Santa Monica Mountains and the San Gabriel Mountains (both the eastern and western sections). Due to the development within the San Fernando Valley and the valley of the Santa Clara River, this is an important corridor for gene flow and species movement.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The SEA contains several populations that are unusual or at the extreme ends of their distributions: Douglas fir, both big cone and the Douglas fir common to the north, and coastal California gnatcatcher at its western extent. Several unusual vegetation alliances are in the Mountains, for example groves of walnuts and flowering ash. Flowering ash within this SEA are uncommonly tall. The Santa Susana Mountains contain some representatives of the desert Palmer's oak, which is unusual in the County.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The relatively undisturbed nature and large size of the plant communities within the Santa Susana Mountains and Simi Hills provides many undisturbed examples of native, natural communities within the County.

In conclusion, the area is an SEA because it contains A) core habitats of listed species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are restricted in distribution in the County and regionally; D) an essential habitat linkage, which is limited in availability in the County; E) unusual populations at the extreme ends of their distributions that are of scientific interest; and F) areas that provide for the preservation of relatively undisturbed examples of original

natural biotic communities in the County.

Tujunga Valley and Hansen Dam SEA

Boundary and Resources Description

The Tujunga Valley and Hansen Dam SEA is located on the northern edge of the San Fernando Valley. The SEA consists of the Tujunga Valley and Wash, starting in the riparian areas of the Big Tujunga, which is the main tributary of the Los Angeles River, within the Angeles National Forest and stretching to include Hansen Dam, Hansen Dam Flood Control Basin, Hansen Dam Park, Hansen Dam Golf Course, Tujunga Wash, and industrial areas downstream of the Hansen Dam. The SEA is entirely in the City of Los Angeles. Most of the part of the SEA upstream, including the Hansen Dam, is an Audubon California designated State Important Bird Area (IBA), which is part of the Los Angeles Flood Control Basin IBA. The Big Tujunga area is recognized for its great importance to migrating birds on the Pacific Flyway as well as the very rare habitat of alluvial fan scrub, which has uncommon resident birds. The Tujunga Wash above Hansen Dam and into the Angeles National Forest beyond the SEA is designated critical habitat for the federally threatened Santa Ana sucker (*Catostomus santaanae*). Two other fishes of the original native four for the Los Angeles River also occur in the Wash and upstream in the Big Tujunga: arroyo chub (*Gila orcuttii*) and an unnamed subspecies of the speckled dace group (*Rhinichthys osculus* ssp. 3).

The SEA is located within portions of the United States Geological Survey (USGS) 7.5' California Quadrangles: San Fernando, Sunland and Van Nuys.

The SEA boundary encompasses the Tujunga Valley Wash and the Hansen Dam Recreation area. The SEA begins in the Angeles National Forest downstream of the confluence of Pipe Canyon with the Big Tujunga, and generally includes the Wash area for much of its extent. The Wash most of the time has water, and from downstream at Hansen Lake (in the Hansen Dam Park) to beyond the upstream area of the SEA, is critical habitat for the federally threatened Santa Ana sucker (*Catostomus santaanae*). This critical habitat extends beyond the SEA and beyond the Big Tujunga Dam to near the headwaters in both Mill Creek and the Upper Big Tujunga Canyon. The state fish species of concern, speckled dace (*Rhinichthys osculus* ssp. 3) and arroyo chub (*Gila orcuttii*) also occur here. A finger extends off the Wash, up a ridge with native chaparral habitat towards Mount Lukens. The boundary travels east to west, and the Wash receives multiple tributaries from the north and south as it flows west. The northeastern end of the Los Angeles Flood Basins IBA begins at about the crossing of Oro Vista Avenue over the Big Tujunga Wash. The clubhouse part of the Angeles National Golf Club next to Foothill Boulevard is excluded from the SEA, but most of the golf course, which has the natural braids of the wash running among its greens, is included in the SEA. East and adjacent to the golf course is a state reserve area for the state and federally endangered slender-horned spineflower (*Dodecahema leptoceras*). This diminutive wash plant is known locally from Santa Clarita to the east end of the San Bernardino Mountains, and south to the Santa Ana Mountains. It is endangered due to all kinds of development that alters wash flood plains. The golf course area once had and may still have some plants.

The Wash, golf course, and the reserve area have an unusual remnant vegetation of alluvial fan scrub, with California junipers (*Juniperus californica*), cactus patches of prickly pear (*Opuntia littoralis*) and cholla (*O. parryi*), in conjunction with the usual riparian and scrub plants. The very sensitive coastal cactus wren (*Campylorhynchus brunneicapillus sandiegensis*) is in residence in this area. The natural area of riparian vegetation is on the banks of the Wash, golf course, and reserve next to the Wash, as well as with a remnant native forest in the stream course of Haines Canyon Creek, which joins the

DRAFT

Wash at the golf course. This kind of habitat, which once covered the bajada of coalescing coastal alluvial fans next to the mountains of the County has been nearly extirpated by development and flood control. It is only represented in a few places in the County. The San Gabriel Canyon SEA has another example at the Santa Fe Dam Recreation Area. Migratory waterfowl often use the ponds of the golf course that are filled by Haines Canyon Creek and the Big Tujunga. The alluvial fan habitat mixed with riparian forest continues across the Interstate 210 as the Big Tujunga Wash approaches and flows into Hansen Dam Park.

The SEA has a broad undercrossing of the Interstate 210, which includes a braid of its tributary Haines Canyon Creek crossing under Foothill Boulevard next to Wentworth Street. The Wentworth Street undercrossing is important to wildlife, as it connects to somewhat dispersed horse ranchettes in the Shadow Hills and from there to the natural areas of the Verdugo Mountains and the Verdugo Mountains SEA. Mountain lions may live in the Verdugo Mountains, which would be an important connection for that population as well as for other animals and plants of the Verdugo Mountains.

The Hansen Dam Park is a meeting area for migratory birds with its perennial water and riparian forest that includes some giant sycamores (*Platanus racemosa*) and cottonwoods (*Populus fremontii*). It is a favorite of bird watchers, and the species list has about 260 species. Unusual species have been observed here, such as the federally endangered coastal California gnatcatcher (*Polioptila californica californica*). Downstream of the Dam, the Hansen Dam Golf Course is included in the SEA, since the pools and mix of native riparian vegetation along with the ornamental vegetation attracts wildlife. The downstream side of the earthen Hansen Dam has been planted with coastal sage scrub, and may be attractive to the gnatcatcher. The quarry and recharge areas have many spots of willow scrub and occasional other native vegetation. The quarry and the recharge pool areas are included in the SEA up to the undercrossing of San Fernando Boulevard.

The area southwest of the Dam is used as a spreading ground. This has created several freshwater marsh areas that are used by marsh birds, migratory waterfowl, and shore birds. The area is also valuable as a wildlife corridor. The vegetation in the Tujunga Valley runs nearly uninterrupted from the foot of the Verdugo Mountains well up into the San Gabriel Mountains. The area has been recognized for its importance, and is used by the Audubon Society and local universities and colleges as a sample of a rapidly disappearing habitat type. As a result, the resources of the area are well known.

Wildlife Movement

Although wildlife movement is hampered by development surrounding the SEA to the south, animals are still able to move through the adjacent hills and valleys, and through the Verdugo Mountains, well up into the San Gabriel Mountains. Due to its large size and topographic complexity, many linkages are certain to occur within the SEA at various bottlenecks. These linkages allow movement between large open space areas within the SEA. Although there are significantly large open spaces within the SEA, contiguous habitat linkages between them are critical in reducing bottlenecks and providing for long term sustainability. A wide variety of wildlife use linkages throughout the SEA, including mountain lion (*Puma concolor*), coyote, mule deer, bobcat (*Lynx rufus*), and a number of medium sized mammals. The Wentworth Street underpass of Interstate 210 is a conduit for wildlife through the dispersed residential neighborhoods of the Shadow Hills. The Shadow Hills are the westernmost extent of the Verdugo Mountains, which provide a large area of natural habitat for wildlife.

Regional Biological Value

The SEA meets all SEA designation criteria and supports many regional biological values. Each

DRAFT

criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE TUJUNGA VALLEY AND HANSEN DAM SEA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Met	The Tujunga Valley Wash supports populations of the federally endangered plants Nevin's barberry and slender-horned spineflower. Most of the SEA is critical habitat for the federally threatened Santa Ana sucker. Two other sensitive native fishes, speckled dace and arroyo chub, co-occur with the sucker. Coastal California gnatcatcher may be a resident of the area. The coastal cactus wren nests in the alluvial fan vegetation of the SEA.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The stream is naturally perennial, but largely controlled by the upstream Big Tujunga Dam. Similar wash and floodplain habitat of these plants and fishes is under considerable pressure from development and from flood control hard channelization throughout Southern California. The rarity of the vegetation extends to other biota that use these areas, and these plants are indicators for a widespread loss of this kind of habitat. Several birds considered species of special concern occur in the SEA. They typically occupy alluvial scrub areas, such as the outwash fans formerly found where mountain canyons exit onto the plain of the Los Angeles Basin, and are in this habitat in the SEA.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	The wash and floodplain habitat of these plants, fishes, and birds is under considerable pressure from development and for flood control hard channelization throughout Los Angeles and is much diminished from the continuous habitat it once was.
D) Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	This SEA is located on one of the main tributaries of the Los Angeles River, Tujunga Canyon and connects with the Hansen Dam flood and recharge area. In spite of the channelization of the Los Angeles River, this area is still an important connecting and migration area for plants and wildlife between the San Gabriel Mountains, the Verdugo Mountains, and the San Fernando Valley. It is an important rest area for aerial fauna traveling between the Santa Monica Mountains and the San Gabriel Mountains.
E) Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The scarcity of natural alluvial wash and fan habitats in Los Angeles ensures that the remaining ones are good areas for scientific study of birds and other organisms that were once more common in the Los Angeles area.
F) Areas that would provide for the	Met	The Tujunga Valley Wash is a relatively undisturbed

DRAFT

<p>preservation of relatively undisturbed examples of the original natural biotic communities in the County.</p>	<p>example of the alluvial washes and fans that once lined the mountains of Los Angeles. It is near a fresh water marsh area used as a spreading ground that is southwest of the dam. A fresh water marsh near the stream exit from the mountains would have been typical of the former configuration with faults along the mountain base, creating uneven ground that would contain marsh pockets. Now most of this type of area has been developed for residences, and most of the washes have been altered as flood control projects. The Wash, therefore, is important to preserve.</p>
--	---

In conclusion, the area is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C), biotic communities, vegetative associations, and habitat of plant or animal species that are restricted in distribution on a regional basis and limited in availability in the County; D) habitat for breeding, feeding, resting, and migrating that is limited both in the County and regionally; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Valley Oaks Savannah SEA

Boundary and Resources Description

The Valley Oaks Savannah SEA is located north of the Santa Susana Mountains, approximately one mile south of the Santa Clara River and one mile north of Pico Canyon. The SEA is bordered on the east by Interstate 5 and is situated between Valencia Boulevard and Stevenson Ranch Parkway. To the west, the SEA is bordered by the foothills of the Santa Susana Mountains. The habitat within the SEA was once the emblem of Spanish California, with rolling hills, grasslands and spaced giant valley oaks. This was a chief habitat in the San Fernando Valley when it was filled with wildlife, such as herds of pronghorn. Under the influence of European settlers, these areas were at first cattle range, and then gradually developed into residential neighborhoods.

The SEA is located in the Newhall United States Geological Survey (USGS) 7.5' California Quadrangle.

The SEA is bordered to the east by Interstate 5 and is situated between Valencia Boulevard on the north and Stevenson Ranch Parkway on the south of the SEA. To the west, the SEA is bordered by the West Ridge residential area of Valencia to the north, and the foothills of the Santa Susana Mountains to the south. The SEA boundary outlines an irregular area of undisturbed grassland savannah with hundreds of valley oaks (*Quercus lobata*). In the north, the SEA includes a small part of the TPC of Valencia (a private golf club) that has retained a number of its valley oaks on the ridges between sections of the golf course. The topography is rolling ridges that trend northeast-southwest along the edge of development. The West Ridge development is on graded areas of these rolling ridges.

The SEA is almost completely undisturbed, except for a few dirt roads. This area contains one of the last remaining stands of valley oak (*Quercus lobata*) in the Santa Clarita Valley. The site consists of specimen trees scattered over the SEA. The adjoining natural hills to the southwest of the SEA have a mixture of plants from the coastal sage chaparral scrub and chaparral communities, which are typical of those found in the Santa Clarita Valley. Other vegetation on the SEA in the southwest area includes

DRAFT

coastal sage chaparral scrub and non-native grasses.

The majority of the SEA consists of undisturbed open space bordered by a few high density residential developments. Open space that adjoins the SEA to the southwest is mostly vegetated with dense stands of chaparral. Other types of vegetation, such as woodlands and grasslands, occur in smaller portions that are scattered throughout the adjacent land on moist or north-facing slopes and canyon bottoms. Lesser amounts of coastal sage chaparral scrub are also present, chiefly as an early successional community in areas that have been previously disturbed.

Wildlife Movement

Wildlife movement within the SEA is limited to local movement of foraging animals. Although the SEA does not support regional corridors itself, adjacent lands to the west and northwest may be important linkages for wildlife movement to and from the Santa Susana Mountains and the Santa Clara River. The location of the SEA, therefore, may be important as a corridor buffer and/or adjacent foraging grounds.

Regional Biological Value

The SEA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE VALLEY OAKS SAVANNAH SEA

Criterion		Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	The County considers oaks as indicators of the presence of important biological communities for preservation, and the uncommon valley oaks of the western areas of the County are especially valued.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The County has the southern boundary of occurrence for the valley oak, which has its core population in the California Central Valley. This makes the valley oak areas important for Southern California — they are very uncommon for the region as a whole.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The valley oak is uncommon in the County, and the western areas with this species are scattered.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in	Met	Oaks are the basic structure for complex communities of organisms. They form shelter and provide many ecosystem functions that facilitate breeding, feeding, resting, and migration. As the basis of the community, it is important to conserve this habitat.

Criterion	Status	Justification
availability either regionally or in the County.		
E) Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The valley oaks of the County are on the southern edge of the species' range. This makes them important as the extreme in physical and geographical limits for the scientific study of the species.
F) Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Although the SEA area is small, this savannah is natural and representative of a natural biotic community that has chiefly been lost in the County.

In conclusion, the area is an SEA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) a natural savannah of valley oaks, once a common habitat on the west side of the County.

Verdugo Mountains SEA

Boundary and Resources Description

The Verdugo Mountains SEA is located within the Verdugo Mountains. This SEA encompasses the Verdugo Mountains south of Interstate 210 and east of the Interstate 5, as well as a portion of these mountains north of Interstate 210.

The Verdugo Mountains are a wilderness island in the middle of the urbanized metropolitan area of the County, surrounded by the cities of Los Angeles, Burbank and Glendale. This area is cherished by the local communities, much of which are designated agricultural with many equestrian properties. The Verdugo Mountains have retained a rural atmosphere despite their proximity to urban Los Angeles. The Verdugo Mountains currently encompass wilderness area, which ranges through various chaparral, coastal sage chaparral scrub, southern willow scrub, coast live oak woodland and forest ecosystems, and many riparian areas with seasonal waterfalls. It is one of the few remaining natural regions in the Los Angeles area that supports abundant native wildlife and habitats, and also contains several rare and sensitive plant and animal species. The geographic location of the Verdugo Mountains makes them important for scientific study, genetic interchange between otherwise isolated populations, and recreation for urban residents.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Burbank, Sunland, and Pasadena.

DRAFT

The SEA is an island refuge, providing what remains of a link between plant and animal populations found in the Santa Monica and San Gabriel mountains. Genetic interchange, by way of this linkage is important in perpetuating the genetic variability in isolated populations, and the maintenance of healthy ecosystems. Chaparral and coastal sage chaparral scrub cover the hillsides of the mountains, with lush riparian vegetation, including California bay (*Umbellularia californica*), western sycamore (*Platanus racemosa*), coast live oak (*Quercus agrifolia* var. *agrifolia*), ferns, and ocellated Humboldt lily (*Lilium humboldtii* ssp. *ocellatum*), which are found in most of the stream drainages. These plant communities provide habitat that is essential to the diverse and abundant fauna that are found in the area. The mountains are also home to the northernmost population of mission manzanita (*Xylococcus bicolor*).

The proximity of the mountains to urban areas provides an excellent opportunity to study the interaction between wild animal populations and humans. The area has already been used for studies concerned with public health. The Interstate 210 crosses the northern edge and there is some intrusion of development along La Tuna Canyon Road. Residential development is beginning to trim the area and has been excluded from the SEA. However, present human use of the natural, native area has been low and has not significantly affected the natural resources found in the Verdugo Mountains.

The northernmost point of the SEA is in the Shadow Hills district of the City of Los Angeles at the west side of the undercrossing of Sunland/Foothill Boulevard beneath the Interstate 210. The freeway frontage is native vegetation and connective to the north with the Wentworth Street undercrossing of Interstate 210. The connection is through native vegetation along the freeway frontage and among dispersed equestrian ranchettes in the Shadow Hills. Wentworth Street ends a short distance east of Interstate 210 at the Angeles National Golf Club, which has its greens interspersed among braided stream courses of the Big Tujunga Canyon alluvial fan. The golf course is part of the SEA. The native vegetation along the freeway, at the restored County Public Works area on the north side of Wentworth, and the native vegetation among the residences of Shadow Hills is all considered an important conduit for wildlife traversing among the San Gabriel Mountains, the SEA, and the Verdugo Mountains.

The SEA boundary follows southward along the southwest side of the Interstate 210 for about a half mile and then crosses to the northeast side of Interstate 210, along a ridge that connects to slopes with natural, native habitat that continues east, along the southern border of the communities of Sunland and Tujunga. In the vicinity of Pasko Peak, the SEA border is drawn around development in the southern edge of the community of Tujunga. The SEA border recrosses the Interstate 210 and then La Tuna Canyon Road. The crossing boundary goes along the east side of a northern tributary of La Tuna Canyon.

A paved road goes along the streamcourse in La Tuna Canyon, and the SEA includes about two miles of native, natural vegetation bordered on the eastern, undeveloped section of the road. The developed western end of La Tuna Canyon Road is excluded from the SEA in a "cherry stem" configuration.

After crossing Interstate 210 and La Tuna Canyon Road, the SEA tracks the southern side of natural habitat along the Interstate 210 for about a mile. The boundary trends irregularly southeast excluding development in the flatter areas that border the Verdugo Wash in the Verdugo City District of Glendale. The SEA includes the incised canyons of Sheep Corral, Cunningham, Henderson, Engleheard, and unnamed others, which are all tributaries of the channelized Verdugo Wash.

In the relatively narrowed area around the State Route 2 (Verdugo Canyon) between the San Rafael Hills on the west and the Verdugo Mountains on the East, the SEA continues irregularly south along the natural, native vegetation of the steep hillsides of the Verdugo Mountains that border Glendale. The SEA

DRAFT

~~includes many unnamed canyons and also Ayars Canyon and Deer Creek at the end of Beaudry Boulevard. The unnamed canyon north of Dead Horse Canyon has a possible wildlands connection with the San Rafael Hills. Its northeast facing slope along Sunshine Drive has chiefly native vegetation and ends in the Verdugo Park on Verdugo Boulevard. Across Verdugo Boulevard is Glendale Community College. The College's eastern border is a natural ridge that connects with the Mountain Avenue overpass of State Route 2 and native vegetation of the San Rafael Hills. Verdugo Boulevard and its development is a substantial block to terrestrial wildlife movement, but aerial fauna and plant seeds can connect fairly easily along this path. In the vicinity of Dead Horse Canyon, the SEA boundary turns generally westward and proceeds irregularly around development in Glendale, including natural parts of Toll, Hillcrest, Sherer, Idelwood, Pomeroy, Brand, and Childs canyons.~~

~~The lobe of the Verdugo Mountains between Hillcrest and Brand canyons is the area with the shortest possible traverse, which is a little less than two miles, to the Santa Monica Mountains in Griffith Park. Aerial fauna and plant seeds can easily make the journey, and the Los Angeles River channel at the base of the Santa Monica Mountains has developed a natural bottom and riparian habitat that must be inviting to migrants. A city park and a cemetery are on the route. The Verdugo Mountains are often viewed as one of the principle connections between the Santa Monica Mountains and the San Gabriel Mountains. However, it must be noted that this corridor is highly fragmented and probably impassable for most terrestrial wildlife.~~

~~Curving to the northwest in Burbank, the SEA boundary includes the natural, native vegetation of Elmwood, Story, Deer, and Sunset canyons. Sunset Canyon excludes a cherry stem shaped area around a development. The Wildwood Canyon Park is included in the SEA with its many stately sycamore and coast live oak trees along the narrow stream course. In Stough Canyon the DeBell Municipal Golf course is excluded, as is the paved section of Stough Canyon Road and the buildings of the Stough Canyon Nature Center; however, the hillsides around Stough Canyon Road are included with a lobe of the SEA including the west ridge of Stough. On the west side of this ridge are the excluded Starlight Bowl and a nearby landfill. Continuing northwest, the SEA includes the natural, undeveloped uppermost elevations of McClure, Brace, Cabrini canyons in Burbank, and Fisher, Jeffries, and Chandler canyons of Sun Valley.~~

~~North of Chandler is the developed south side of La Tuna Canyon Road, where the SEA boundary turns west along natural vegetation, using lot lines in part, which delineate the extent of fuel modification in this area of high fire hazard. The SEA boundary includes the north-facing south side of La Tuna Canyon for a distance of about three miles. The boundary crosses the road at the point where development stops and natural vegetation is on both sides of the road. From this point, the boundary continues the cherry-stem exclusion westward along the border of natural vegetation on the south-facing slope of La Tuna Canyon. Near the northwestern end of the Verdugo Mountains and La Tuna Canyon, the La Tuna stream joins the West Burbank Flood Control Channel. The SEA boundary includes the McDonald Creek drainage (tributary of La Tuna) and loops around the north ridge of McDonald Creek, changing direction to the northeast. The SEA boundary includes the natural area along the northern edge of the Verdugo Mountains by again following the edge of development in the Shadow Hills district of the City of Los Angeles. The boundary joins the northernmost point of the SEA near the undercrossing of Interstate 210 for Sunland/Foothill Boulevard.~~

~~The SEA is wholly within incorporated boundaries (cities of Los Angeles, Glendale, and Burbank), but much is preserved in conservation easements under the guidance of the Santa Monica Mountains Conservancy.~~

Wildlife Movement

Although wildlife movement is hampered by adjacent rural development in proximity to the SEA, animals are still able to move through the Verdugo Mountains in many areas. Due to its large size and topographic complexity, many linkages occur within the SEA at various bottlenecks. These linkages allow movement between large open space areas within the SEA, as well as between areas outside the SEA toward the Angeles National Forest. The genetic flow through these areas is crucial in maintaining the diversity and viability of the species within the Verdugo Mountains. Some areas of probable, possible, and perhaps future connection have been indicated in the General Boundary and Resources section. Although there are significantly large open spaces within the SEA, contiguous habitat linkages between them is critical in reducing bottlenecks and providing for long term sustainability. A wide variety of wildlife use linkages throughout the SEA, including mountain lion (*Puma concolor*), coyote (*Canis latrans*), mule deer (*Odocoileus hemionus*), bobcat (*Lynx rufus*), as well as a number of medium-sized mammals.

Regional Biological Value

The SEA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE VERDUGO MOUNTAINS SEA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Not Met	No critical habitats are designated within this SEA.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The Verdugo Mountains combined is an extensive, relatively undisturbed island of natural vegetation in an urbanized area, which is very rare in Southern California.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	The Verdugo Mountains combined is an extensive, relatively undisturbed island of natural chaparral and canyon vegetation. It is an important island refuge for migration among the mountain ranges of the northern portion of the County. Aerial animals and plant seeds can easily traverse the distances between the Verdugo Mountains and Santa Monica, Santa Susana, and the San Gabriel mountains. The Verdugo Mountains serve as the centerpiece of these connections. The hillsides are covered by chaparral and coastal sage chaparral scrub. The canyons' riparian vegetation includes California bay, sycamores, ferns and tiger lilies. These plant communities provide habitat that is essential to the diverse and abundant fauna found in the area.
D) Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds	Met	The Verdugo Mountains serve as an arm of the San Gabriel Mountains, extending towards the eastern end of the Santa Monica Mountains in Griffith Park—only two miles distant. Aerial animals and plant seeds easily

DRAFT

	and is limited in availability either regionally or in the County.		cross the gap. The Verdugo Mountains are exceedingly important for connections among the Santa Monica, Santa Susana, and the San Gabriel mountains. Genetic interchange, by way of this linkage, is important in perpetuating the genetic variability in isolated populations, which maintains healthy ecosystems and resilience to climate change.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The geographic location of the Verdugo Mountains makes them important for scientific study, genetic interchange between otherwise isolated populations, and recreation for urban residents. The area has already been used for studies concerned with public health.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	The Verdugo Mountains have been impacted by dirt roads, one paved road in Tuna Canyon, fuel breaks, transmission lines and isolated buildings for houses, radio towers, and water tanks. Because of their extent, however, the Verdugo Mountains are still considered largely natural and little impacted—a prime example of the chaparral and coastal sage chaparral scrub once prevalent in the County coastal areas.

In conclusion, the area is an SEA because it contains B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Altadena Foothills and Arroyos SEA

General location and landform

The Altadena Foothills and Arroyos SEA is located in the westernmost portion of the San Gabriel Valley. This SEA includes both incorporated and unincorporated areas. The SEA represents the lower elevation/urban interface portions of Millard, Alzada, Chiquita, Las Flores, Rubio, and Eaton canyons from the urban edge to undeveloped wildland areas of the lower elevations of the San Gabriel Mountains.

The SEA is located within the Mount Wilson and Pasadena United States Geological Survey (USGS) 7.5' California Quadrangles.

The SEA recognizes one of the principal ecotones of the southern California coastal areas, the area where the sediment of the coastal alluvial fans from the mountain streams and drainages exits the abrupt upthrust rock of the mountains. Here one finds the biotic communities of the mountains meeting the communities of the coastal plain areas, combining with the organisms that are only found at the junction. The natural habitats of this kind of biological area have largely been lost as urban communities expand to the limits of easily buildable space. The chief attribute of this SEA is a high diversity of species within a relatively small area, due to the presence of significant riparian habitat as well as the SEA's position between the mountain biome and the valley biome, caused by an abrupt

DRAFT

change of slope formed by the San Gabriel Mountains. The SEA has as its center the dividing ridge between the two principal rivers of the Los Angeles Basin, the Los Angeles River and the San Gabriel River.

Biological Resources

Vegetation and communities

The wide range of elevation, topography, aspect, and geology represent a diverse array of physical habitats within this SEA. In general, the topography of the SEA is moderately steep to very steep, which results in very narrow drainages with elevations ranging from a high of approximately 2,400 feet above mean sea level (MSL) to a low of approximately 1,200 feet above MSL. Consequently, a variety of plant communities exist, including riparian and upland shrublands and woodlands. Within these major community types, there are many vegetation series that vary according to plant species dominance. The SEA supports significant foothill formations of riparian forest, as well as Diverse coastal sage scrub and chaparral associations, extensive and diverse stands of oak woodland, and in upper reaches, dense bigcone Douglas-fir forest.

This SEA lies at the extreme northwestern extent of the Peninsular floristic province, with representative plant species such as Engelmann oak (*Quercus engelmannii*) reaching the limit of their ranges here. The gneiss soil present supports the San Gabriel oak (*Quercus durata* var. *gabrielensis*), as well as a restricted population of *Lupinus formosus*, formerly described as Pasadena lupine (*Lupinus pasadenensis*). Other rare plants known from the upper-elevation portions of the SEA include San Gabriel manzanita (*Arctostaphylos glandulosa* ssp. *gabrielensis*), San Gabriel bedstraw (*Galium grande*), San Gabriel linanthus (*Linanthus concinnus*) and Greata's aster (*Symphotrichum greatae*).

California spotted owl (*Strix occidentalis occidentalis*) still maintains a resident population in the more remote canyons of the SEA, and Mountain lion (Southern California/Central Coast ESU) are commonly detected on wildlife cameras.

Several wildlife species approach their lowland range limit here, including Merriam's chipmunk and American black bear.

Wildlife Movement

Wildlife movement within the SEA takes on two major forms. First, due to the extremely steep intervening topography, considerable movement of wildlife up and down the drainages, which course through this SEA to connect the forest interior with foothill areas, is expected. This type of movement occurs year round, and is particularly important for large mobile mammals such as mountain lion, mule deer and black bear that meet their full range of habitat needs over broad areas. The second major type of movement occurs across the flanks of the foothills in an east-west direction. Particularly for riparian-obligate and riparian-favoring migratory birds, the lower elevation riparian habitats in the SEA such as the Arroyo Seco at Hahamongna Watershed Park and the adjacent Arroyo Seco are of high importance and heavily utilized.

The SEA serves as the only corridor to provide interacting component habitat areas for species to move from low basin and foothill elevations to the sub-alpine elevations of the high San Gabriel Mountains. In addition, the foothill zone of the San Gabriel Mountains represents a key migratory pathway for a wide variety of songbirds and raptors (e.g., Swainson's hawk, Turkey Vulture). A long-term banding station at Bear Divide intercepts birds moving northwest from this SEA toward the Sierra Nevada.

DRAFT

With conservation planning, there is potential for north-south wildlife movement between the Angeles National Forest and the Verdugo Mountains via numerous small (urbanized) drainages, as well as via the Arroyo Seco into the San Rafael Hills. The Arroyo Seco is the eastern limit of this link and creates a potential movement corridor from the Angeles National Forest, under Interstate-210. Across Interstate-210, the linkage enters the San Rafael Hills, where blocks of habitat remain in the cities. Some are conserved in natural open space, such as Cherry Canyon Park and Open Space Preserve of the City of La Cañada-Flintridge, just south of the County Descanso Gardens. These open spaces are interspersed with residential development and have been recommended for inclusion into the existing Verdugo Mountains SEA. From the San Rafael Hills, linkage potential may be traced to the west across State Route 2 and Verdugo Wash, past enclaves of residential development to access the Verdugo Mountains.

Designation Criteria

The SEA meets important SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE ALTADENA FOOTHILLS AND ARROYOS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	<u>Met</u>	<ul style="list-style-type: none">- <u>Mountain lion (Southern California/Central Coast ESU)</u>- <u>San Gabriel manzanita</u>- <u>San Gabriel bedstraw</u>- <u>San Gabriel linanthus</u>- <u>Greata's aster</u>- <u>Palmer's mariposa-lily</u>- <u>Golden eagle</u>- <u>California spotted owl</u>- <u>Least Bell's vireo</u>

DRAFT

Criterion		Status	Justification
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>San Gabriel oak</u> – <u>Engelmann oak</u> – <u>Plummer’s mariposa-lily</u> – <u>Chaparral rein orchid</u> – <u>Coast Range newt</u> – <u>Coast horned lizard</u> – <u>Coast patch-nosed snake</u> – <u>Yellow-breasted chat</u> – <u>Significant riparian woodlands and forest</u> – <u>Diverse coastal sage scrub and chaparral associations</u> – <u>Extensive and diverse stands of oak woodland</u> – <u>Bigcone Douglas-fir forest.</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Only available low-elevation corridor for species to feed, rest, and migrate from low basin and foothill elevations to the high San Gabriel Mountains.</u> – <u>Key migratory pathway for a wide variety of songbirds, raptors, and other bird species.</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Extreme northwestern extent of the Peninsular floristic province (e.g., northernmost Engelmann oak).</u> – <u>Edaphic and geologically restricted San Gabriel oak and Pasadena lupine (<i>Lupinus formosus</i> / <i>L. pasadenensis</i>)</u> – <u>Low-elevation limit of Merriam’s chipmunk and black bear.</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Extensive stands of low-elevation foothill scrub, chaparral, and canyon woodland communities</u>

Criterion		Status	Justification
G)	Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.	Met	– East-West and North-South movement opportunities across the San Gabriel foothills.

Antelope Valley SEA

General location and landform

The Antelope Valley SEA is located in the eastern portion of the Antelope Valley, primarily east of the cities of Palmdale and Lancaster, within a predominantly unincorporated area of the County. The SEA was originally focused on two principal watercourses of the area, Little Rock Wash and Big Rock Wash and its tributaries, such as Mescal Creek. However, recognizing the continuing loss of open desert habitat within the County (and the western Mojave Desert), we recommend expanding the boundaries to better include habitats outside these drainages, taking in a) the entire alkali sink scrub ecosystem north and east of Lancaster, b) extensive and largely undisturbed creosote bush flats in the far northeastern corner of the County, and c) various formations of Joshua tree woodland at different elevations and soil types.

The SEA is located, at least partially, in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Rosamond, Rosamond Lake, Redman, Rogers Lake South, Jackrabbit Hill, Lancaster East, Alpine Butte, Hi Vista, Adobe Mountain, Palmdale, Littlerock, Lovejoy Buttes, El Mirage, Pacifico Mountain, Juniper Hills, Valyermo, and Mescal Creek.

Biological Resources

Vegetation and communities

Watercourses and water features, such as dry lakes and alkali sinks, alluvial washes, and springs, are the focus for desert wildlife and central to connectivity and biodiversity in this region. Audubon California recognizes the area of Edwards Air Force Base as a Globally Important Bird Area (IBA), which is visited by tens of thousands of migrant birds during the spring and fall migratory seasons, and supports the breeding of rare and endangered birds during the spring and summer months. The SEA traverses the Antelope Valley from the foothills of the San Gabriel Mountains, to the low elevations of the dry lake basins, and its expanse and considerable topographical relief is reflected in its relatively high floral and faunal diversity. The SEA includes playa lake, alkali marsh, alluvial fan scrub, a mosaic of xeric desert scrubs, Joshua tree woodland, desert riparian woodlands, juniper scrub, pinyon pine, chaparral and mixed conifer, oak, and riparian communities of higher elevations. Transitional zones (ecotones) between these communities often contain unusual species compositions, such as pinyon pine, juniper and Joshua trees together, or Joshua trees adjacent to cottonwood forest.

While the SEA was originally delineated to emphasize the importance of the Little Rock Wash and Big Rock Wash watersheds to the surface and subsurface hydrology of the Antelope Valley and to the dry

DRAFT

lakes and sinks in the region, significant upland habitats may also be found in this SEA, including extensive Joshua Tree woodland. Here in the eastern Antelope Valley, these trees are larger than those toward the west, and are more spread out, with native shrubs and wildflower patches filling the gaps between trees. The SEA also includes several major rocky buttes and numerous minor ones, which have highly diverse biota along with diverse desert habitats, which range from sand dunes formed from the wind-blown dust that the buttes collect, to rocky crags, which are home to nesting and roosting raptors and bats, and a distinctive flora.

The origins of the watercourses in the high San Gabriel Mountains are an important aspect of their diversity and connectivity, enabling the free movement of wildlife upslope and downslope, and allowing for long term migration of its plant communities in response to a warming, drying climate.

The SEA includes the County's portion of the watershed basin for dry lakes, the ultimate destination for the watercourses. There are three dry lakes and their adjacent plains (part of Edwards Air Force Base) included in the SEA: Rosamond Dry Lake with the adjacent Piute Ponds, Buckhorn Lake, and Rogers Lake. These lakes and ponds are often flooded during the rainy winter-spring seasons, and are a key resting area in the region for migrant waterbirds on the Pacific Flyway. The northern portion of the SEA encompasses some agricultural cropland (an increasing portion of which is fallow) and dispersed rural residential uses; however, the underlying hydrology of the washes remains intact throughout the entire SEA.

All of Edwards Air Force Base that is in the County is included in the SEA because the restricted entry and use protect the dry lakes and their neighboring areas. Many desert plants and wildlife species once found broadly across the Antelope Valley are now found only or primarily within Edwards Air Force Base. The ponds and dry lakes have distributed habitat of marshy alkali grassland, alkali flats, and cattail and bulrush marsh augmented by wastewater treatment facilities that have additional ponds. Some of the nesting rare and uncommon birds include white-faced ibis (*Plegadis chihi*), tricolored blackbird (*Agelaius tricolor*), redhead (*Aythya americana*), gadwall (*Anas strepera*), yellow-headed blackbird (*Xanthocephalus xanthocephalus*), least bittern (*Ixobrychus exilis*), and federally-threatened western snowy plover (*Charadrius alexandrinus nivosus*).

Little Rock Wash flows north from Little Rock-Palmdale Dam. Upstream from the reservoir is critical habitat for the endangered arroyo toad (*Anaxyrus californicus*).

The South Fork of Big Rock Wash is part of the federally-designated critical habitat of the mountain yellow-legged frog (*Rana muscosa*). This frog is known in the County from only a few high-mountain streams in the San Gabriel Mountains. A fungal pathogen is principally responsible for its decline; however, climate change, air pollution and non-native predators are also likely contributing factors, and it is now wholly dependent on active restoration/captive-rearing efforts coordinated by federal agencies and Los Angeles Zoo.

This northeastern part of the SEA includes designated conservation areas (including several County sanctuaries) that support the last remaining populations of the federally-threatened Mojave desert tortoise and state-threatened Mohave ground squirrel in the county. In this area of creosote-bush flats and cactus-dotted rocky buttes, many desert species reach their westward distributional limit in the county (and for some, the state). This habitat is essentially continuous east to the Colorado River, broken only by long stretches of freeways and associated clusters of development. However, within Los Angeles County, it is becoming increasingly fragmented and disturbed by human uses, some of which were not envisioned 20 years ago (e.g., industrial solar, cannabis grow operations).

DRAFT

Edwards Air Force Base supports the last large stands of mesquite (*Prosopis glandulosa*) remaining in the County. It also supports rare flora such as the Mojave spineflower (*Chorizanthe spinosa*) and alkali mariposa lily (*Calochortus striatus*). Saltbush (*Atriplex* spp.) diversity is among the highest in the state here, and Rosamond Lake has the best example of the shadscale scrub and alkali sink biotic communities in the County. Shadscale (*Atriplex confertifolia*) scrub needs heavy soil with underlying hardpan between 3000-6000 feet elevation, which is unusual in the County, and more common in the north Mojave Desert and Owens Valley. In addition, the area represents the southernmost extension of elements of the Great Basin ecosystem, including the Great Basin kangaroo rat (*Dipodomys microps*), which occurs as a geographically isolated population.

The areas to the south in the SEA support some of the most floristically diverse habitats in the county, including multi-species oak and conifer woodlands that are characteristic of the middle-elevation zones on the north face of the San Gabriel Mountains. The creeks are higher energy systems at those elevations, as they collect water from the surrounding terrain, and are typically lined with woodlands of alder, willow, sycamore and cottonwood.

As the creeks drop north of the pressure ridges of the San Andreas Fault Zone, they lose gradient and widen, and most of the flow becomes sub-surface, except during high energy storms or in the spring (depending upon rainfall totals in the watersheds). The vegetation here becomes sparser and less evenly distributed along the channel margins. Crossing the lowlands of the Antelope Valley, the channels support a variety of desert scrub vegetation within the alluvial plains. Where the alluvial plains are wide and shallow (e.g., at Valyermo), cottonwood-willow woodland and sycamore woodland vegetation communities occur within the overall floodplain on stable terraces; around oxbow flow zones in the Antelope Valley; or where the groundwater table is replaced or augmented by agricultural runoff. Intact Joshua tree woodland, with native understory present, supports a relatively high diversity of native annual wildflowers, reptiles and mammals (as evidenced by abundant tracks and burrows just about anywhere in this SEA). The Joshua trees also provide nest sites for many resident and migratory bird species.

Lovejoy, Alpine, Piute, Black and Saddleback buttes, along with other, smaller unnamed buttes, form most of the topographical relief within the SEA. These areas offer different ecological conditions that are associated with rock shelter, perching sites, nesting sites, denning areas, wind protection and sand sheet accumulation areas. Local and migratory bat species roost and reproduce in the caves and crevices of the butte formations. The higher buttes provide local nesting sites for owls and other birds of prey.

Alpine Butte is the least disturbed butte in the County, with excellent stands of Joshua tree woodland and creosote bush scrub, and impressive wildflower displays when rainfall creates appropriate conditions. Lovejoy Butte has Joshua tree woodland and creosote bush scrub, with a central wind-blown sand community for a good mixture of rock and sand habitats. In addition, the close proximity of Lovejoy Butte to Big Rock Wash increases the diversity of habitats in the area. Nevertheless, it also suffers from impact from the Lake Los Angeles community, which borders the butte on three sides. The clustering of buttes in the SEA may be important to the abundant, diverse wildlife that inhabits the various vegetation communities around and in the buttes. Saddleback Butte and Piute Butte together are protected as a state park, but Saddleback Butte is also subject to development for campsites and hiking trails. Piute Butte has a prehistoric site that may protect it from much future recreational development. All of the buttes harbor diverse wildlife and flora. Most of them are critical habitat for the state and federally-threatened Mojave desert tortoise. Some buttes within the desert tortoise's critical habitat are not included in the SEA.

DRAFT

The active and fallow open agricultural lands support a diversity of bird species typical of grassland habitats, such as Swainson's and ferruginous hawks, Short-eared and burrowing owls (both now very rare), Western Meadowlark, and Vesper sparrow. A variety of local and migratory bat species feed over the irrigated fields in the spring and summer, when insect numbers are the highest, and at least one sensitive bat species, the pallid bat, forages in open scrub or ruderal desert habitats.

The northern portion of the SEA contains several unique habitat types, including mesquite bosque (threatened locally by lowering water tables and harvest for firewood), clay pan pools, vernal pools, alkali grasslands, alkali and freshwater marshes, and permanent ponds. Hundreds of bird species have been recorded from the pond and marsh habitats around the dry lakes and ponds, and numerous species nest on the playa margins or in the associated riparian habitats. The open creosote scrub and other xeric habitats on the slopes surrounding the lake playas serve as important wintering areas for many raptor species, as well as large numbers of songbirds.

Wildlife Movement

The geographical features of the SEA serve as a major habitat linkage and movement corridor for all wildlife species within its vicinity, allow for both lateral movement (east-west) and well as elevation movement into the San Gabriel Mountains. Ecologically generalist species (mountain lion, bobcat, coyote, gray fox, etc.) have the ability to move across such vast areas and through changing habitat types. For such species, the SEA may serve as an important system for long-term and genetic exchange among populations. For smaller or less-mobile species or taxa, which are narrowly restricted in their habitat needs, the SEA can serve as a broad linkage zone, in which individual movement can take place during seasonal population dispersal or over generations. This provides essential genetic exchange within and between metapopulations. The two drainages (Big Rock Creek and Little Rock Creek), combined with the upland terrestrial Desert-Montane transect portion of the SEA, ensure linkage and direct movement areas for essentially all of the wildlife species present within the County portion of the Antelope Valley.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

DRAFT

CRITERIA ANALYSIS SUMMARY OF THE ANTELOPE VALLEY SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>Critical habitat for the only known Antelope Valley population of arroyo toad (adjacent to Little Rock Reservoir, upstream in Little Rock Creek)</u> – <u>Mojave desert tortoise</u> – <u>Mohave ground squirrel</u> – <u>Mountain yellow-legged frog (South Fork of Big Rock Creek).</u> – <u>Alkali mariposa lily</u> – <u>Habitat for up to two dozen endangered or threatened plants and wildlife, including some of the rarest plant species in the County, such as Rosamond eriastrum and Lancaster milkvetch.</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>More than 50 plant and wildlife species considered rare at the regional or County level.</u> – <u>Unique mesquite bosque, sand sheet, rocky butte, desert riparian woodland, and alluvial fan sage scrub habitats regionally restricted biotic communities encompassed by the SEA.</u> – <u>Dozens of desert species not found (or rarely found) elsewhere in the County are still common here.</u> – <u>Regionally-significant ponds and other riparian and wetland systems in the northern portion of the SEA (on Edwards AFB) that support breeding populations of water birds and raptors not found elsewhere in the County (e.g., white-faced Ibis and burrowing owl).</u>

DRAFT

	Criterion	Status	Justification
D)	<p><u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, migrating grounds and is limited in availability either regionally or in the County.</u></p>	Met	<ul style="list-style-type: none"> – <u>Freshwater habitats that support waterbird nesting colonies not found elsewhere in the region, and more typical of Great Basin wetlands to the northeast.</u> – <u>Rocky desert buttes are unique roosting, sheltering, perching and nesting sites for birds of prey and bats.</u> – <u>Centered on migratory routes for both plants and animals along principal desert washes and buttes that connect the mountains to freshwater playas.</u>
E)	<p><u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u></p>	Met	<ul style="list-style-type: none"> – <u>Mesquite bosque at geographical range extreme, including associated biota (e.g., mesquite borer).</u> – <u>Joshua Tree across elevational gradient, including mixed stands with montane forest species along Big Pines Highway.</u> – <u>Northwestern range limit of multiple Colorado Desert ecosystem elements.</u>
F)	<p><u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u></p>	Met	<ul style="list-style-type: none"> – <u>Biotically intact Joshua tree woodland, desert riparian woodland, and desert alluvial fan sage scrub.</u> – <u>Among the most intact remaining desert community examples in Los Angeles County.</u> – <u>Remnant mesquite bosque representing formerly widespread Antelope Valley vegetation.</u>

Criterion	Status	Justification
G) <u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Major east-west and elevational habitat linkage into the San Gabriel Mountains.</u> - <u>Big Rock Creek and Little Rock Creek drainages functioning as movement corridors.</u> - <u>Desert-montane transect providing cross-habitat connectivity.</u> - <u>Facilitates genetic exchange within and between regional wildlife populations.</u>

Baldwin Hill SEA

General location and landform

The Baldwin Hills extends over roughly two square miles on the floor of the Los Angeles Basin, just east of Los Angeles International Airport. It is located within the lower Ballona Creek Watershed south of the creek channel, which historically formed a large complex of sloughs and marshes (“ciénegas”) centered east of the hills. The hills themselves were uplifted from the sedimentary soil of the basin floor, and form the northern terminus of a series of low hills extending north along the Newport – Inglewood Fault. The northern face of the hills is steep, while the southern edge features low ridges and incised canyons.

Coastal scrub vegetation within the 1,000-acre Inglewood Oil Field has been free of most human disturbance for more than a century, as this entire property is fenced. From the outside, one may see large cactus patches, *Dudleya* (cf. *lanceolata*), and other elements that have been lost in other open space patches on the Los Angeles Basin floor. So, while oil operations have continued, most other impacts, such as trampling and human disturbance, have been minimal.

The Baldwin Hills SEA is located within the Beverly Hills, Hollywood, Venice, and Inglewood United States Geological Survey (USGS) 7.5' California Quadrangles.

Surrounded by dense urban development, the Baldwin Hills represent a mosaic of ownership and land uses, including the following areas with native/natural vegetation:

- Inglewood Oil Field (c. 1000 acres, a mix of active oil extraction areas and pockets of native coastal sage scrub and seasonal wetland vegetation).
- Kenneth Hahn State Regional Park (401-acre recreational area with considerable reserves of native coastal sage scrub habitat, east of La Cienega Blvd.).
- Baldwin Hills Scenic Overlook State Park (8.5-acre park with active habitat restoration program, south of Jefferson Blvd.).
- Stoneview Nature Center (5-acre park west of La Cienega Blvd.).
- Holy Cross Cemetery (200-acre, largely manicured area of lawn and planted trees, but with a portion of undeveloped habitat toward the back/northern end).

Biological Resources

Vegetation and communities

The Baldwin Hills represent the most significant reserve of low-elevation coastal scrub habitat in Los Angeles County away from the Palos Verdes Peninsula. Although this scrub contains little sage (*Salvia* spp.), it does have a strong component of California buckwheat (*Eriogonum fasciculatum*) and a prickly-pear cactus (*Opuntia* cf. *littoralis*). A more mesic scrub is present on north-facing slopes and at seeps, notably containing blue elderberry (*Sambucus mexicana*), giant wildrye (*Elymus condensatus*) and western poison-oak (*Toxicodendron diversilobum*). Notable plant occurrences include relict stands of purple needlegrass (*Stipa pulchra*), greenbark ceanothus (*Ceanothus spinosus*), and the only documented occurrence of Nuttall's scrub oak (*Quercus dumosa*), a CNPS Rare plant, in Los Angeles County.

While the hills still support isolated populations of many lowland wildlife species, including gray fox (*Urocyon cinereoargenteus*), regionally-rare wildlife species have been extirpated over time, with notable losses including San Diego black-tailed jackrabbit (*Lepus californicus bennettii*)—last recorded in 2000—and cactus wren (*Campylorhynchus brunneicapillus*), last seen in 1996. However, the Federally Threatened coastal California gnatcatcher (*Polioptila californica californica*) appears to be established now, with regular sightings of multiple individuals. The red coachwhip (*Masticophis flagellum piceus*) also maintains a small population in the hills based on photos uploaded to iNaturalist, completely isolated from nearest populations in the Long Beach and Whittier Narrows area. The Trask shoulderband snail (*Helminthoglypta traskii*) appears to be common here, which has a NatureServe Rank of S2 (Imperiled, at high risk of extirpation in the state due to restricted range, few populations or occurrences, steep declines, severe threats, or other factors), as is Crotch's bumblebee (*Bombus crotchii*), a Candidate for State listing. Southern California black walnut (*Juglans californica*, CNPS Rank 4) also occurs, within mesic scrub, isolated from nearest occurrences by several miles.

Many raptor nests are present, with all the known species found in the Los Angeles area present and either nesting, or suspected of doing so (e.g., American kestrel (*Falco sparverius*), American barn owl (*Tyto furcata*)).

Wildlife Movement

Major connectivity opportunities exist along the Ballona Creek channel to and from the Ballona Wetlands/El Segundo Dunes (both SEAs), which otherwise would be completely isolated. In addition, the entire Baldwin Hills are the largest patch of habitat between the Palos Verdes peninsula and the Santa Monica Mountains, so function as a “stepping stone” between these two largely protected areas.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE BALDWIN HILLS SEA

DRAFT

<u>Criterion</u>		<u>Status</u>	<u>Justification</u>
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> - <u>Coastal California gnatcatcher</u> - <u>Nuttall’s scrub oak</u>
B/ C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> - <u>Low-elevation coastal scrub on basin floor</u> - <u>Nuttall’s scrub oak occurrence (county-unique)</u> - <u>Trask’s shoulderband snail (S2)</u> - <u>Crotch’s bumble bee (Candidate)</u> - <u>Southern California black walnut (isolated population)</u> - <u>Red coachwhip (isolated population)</u> - <u>Relict purple needlegrass stands</u> - <u>Greenbark ceanothus</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Not Met	
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> - <u>Isolated basin-floor reptile populations</u> - <u>Isolated mammal populations</u> - <u>Urban-edge remnant coastal scrub system</u> - <u>High raptor nesting density</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> - <u>Extensive remnant low-elevation coastal scrub</u> - <u>Relatively intact basin-floor scrub habitat</u> - <u>Relict native grassland (purple needlegrass)</u>

<u>Criterion</u>	<u>Status</u>	<u>Justification</u>
<p>G) <u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u></p>	<p>Met</p>	<ul style="list-style-type: none"> - <u>Linkage along Ballona Creek corridor</u> - <u>Stepping stone habitat between Palos Verdes Peninsula and Santa Monica Mountains</u> - <u>Largest habitat block between coastal and inland open space</u>

Colorado Lagoon SEA

General location and landform

Historically a component of the greater Los Cerritos Wetlands, Colorado Lagoon is located in the southeastern corner of the City of Long Beach, where it feeds into upper Alamitos Bay at Marine Stadium. The existing 18-acre lagoon and six acres of associated natural uplands are currently being expanded by another six acres through an ambitious project to daylight and restore an open channel to Alamitos Bay that was filled in decades ago. In addition to expanding the area of natural habitat by 25 percent, the intent of the open channel project is to improve tidal circulation in the lagoon to better support stands of eelgrass that were restored as part of the above-mentioned improvements. At the time of the last SEA update, in 2000, Colorado Lagoon consistently ranked among the most polluted and impaired coastal water bodies in Los Angeles County, and it was not known to have high ecological values. The lagoon has since undergone extensive improvements to remove toxic sediments, filter and re-route urban runoff before it enters the lagoon, and restore coastal marsh and native upland scrub around the water’s edge, actions that have dramatically improved habitat conditions during the last decade.

The SEA is located within the Downey United States Geological Survey (USGS) 7.5' California Quadrangle.

Although not a formal SEA Selection Criterion, it is relevant that wetlands historically associated with the San Gabriel River, including Colorado Lagoon, are considered crucial to the Gabrieleño-Kizh/Tongva and Juaneño/Acjachemen peoples, for whom the permanent coastal settlement of Puvunga, including Motuucheyngna, is recognized as a major, sacred site of emergence. As such, formal recognition of the ecological significance of Colorado Lagoon through SEA designation would be consonant with and respectful of TEK.

The first phase of community-based restoration at Colorado Lagoon lasted from 2008 to 2011, when volunteers manually removed 25,000 pounds of non-native invasive ice plant (various species) and installed 3,500 native plants in about two acres of intertidal wetlands and low-lying uplands on the lagoon’s eastern shore. The second and third phases of restoration, addressing the lagoon’s western arm, continued through 2014.

Biological Resources

Vegetation and communities

In its improved incarnation, Colorado Lagoon is widely recognized as an important area for avian life. Birders have documented more than 240 bird species there, including regular sightings of Endangered California least terns (*Sterna antillarum browni*) during the summer, when adults bring fish from the lagoon to their colony at nearby Anaheim Bay to the south. On occasion, the terns will bring recently fledged young to the lagoon.

Piscivorous species found at the lagoon year-round include California brown pelicans (*Pelicanus occidentalis californicus*), double-crested cormorants (*Nannopterum auritum*), snowy egrets (*Egretta thula*), and pied-billed grebes (*Podilymbus podiceps*). During winter, dozens of Lesser Scaup (*Aythya affinis*) can be found bobbing on the surface, often accompanied by such species as American wigeon (*Mareca americana*), redhead (*Aythya americana*), bufflehead (*Bucephala albeola*), and ruddy duck (*Oxyura jamaicensis*), and flocks of several dozen gulls congregate along the southern shore daily. Shorebirds utilize the lagoon daily, mainly during migration, when least and western sandpipers pass through along with such larger species as the willet (*Tringa semipalmata*), marbled godwit (*Limosa fedoa*), and long-billed curlew (*Numenius americanus*).

Below the water's surface, the lagoon functions as a natural fish hatchery. It supports approximately 21 species of marine fish, such as the California halibut (*Paralichthys californicus*), yellowfin croaker (*Umbrina roncadore*), and gray smooth-hound (*Mustelus californicus*), as well as bat rays (*Myliobatis californica*) and Haller's round rays (*Urobatis halleri*). These populations benefit from the lagoon's eelgrass beds, which provide essential nursery grounds and hunting territory for predatory species, and which are now very rare in southern California. The benthic environment is equally rich, inhabited by a variety of invertebrates including striped shore crabs, clams, and sea slugs. Visitors often observe moon jellies drifting near the shoreline or washing up on the sandy beaches, a testament to the lagoon's direct tidal exchange with Alamitos Bay. The beaches provide habitat for s-banded (*Cicindela trifasciata*) and wetsalts tiger beetles (*Cicindelidia hemorrhagica*). Also resident is the wandering [saltmarsh] skipper (*Panoquina errans*), with a NatureServe Rank of S2 (Imperiled, at high risk of extirpation in the state due to restricted range, few populations or occurrences, steep declines, severe threats, or other factors).

Rimming the lagoon is a band of intertidal wetland (a habitat widely lost in the region), including at least 17 native salt marsh plant species. Restoration projects have replaced invasive flora with such southern California natives as coastal goldenbush (*Isocoma menziesii*), California sunflower (*Encelia californica*), dune buckwheat (*Eriogonum parvifolium*), and giant coreopsis (*Leptosyne gigantea*), which provide habitat for a variety of butterflies, bees, wasps, spiders and other invertebrates that provide prey for resident songbirds such as common yellowthroats (*Geothlypis trichas*). Sensitive and noteworthy plant species include woolly seablite (*Suaeda taxifolia*) and southwestern spiny rush (*Juncus acutus* ssp. *leopoldii*), both California Rare Plant Rank 4.2 (watch list).

Wildlife Movement

Colorado Lagoon is not located along any terrestrial wildlife movement corridor or habitat linkage, but it does function as an important stopover site for a variety of shorebirds during spring and fall migration periods.

DRAFT

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE COLORADO LAGOON SEA

	<u>Criterion</u>	<u>Status</u>	<u>Justification</u>
A)	The habitat of core populations of endangered or threatened plant or animal species.	Not Met	
B/ C)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	<ul style="list-style-type: none"> - eelgrass beds - intertidal salt marsh - sandy beach habitat - saltmarsh skipper (S2) - woolly seablite (CRPR 4.2) - southwestern spiny rush (CRPR 4.2)
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	<ul style="list-style-type: none"> - wintering waterfowl concentrations - migratory shorebird stopover habitat - waterbird foraging habitat - fish nursery habitat associated with eelgrass beds
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Not met	
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Not met	- restored eelgrass and intertidal marsh habitats; expected to meet Criterion F as restoration matures

G)	<p><u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u></p>	<p><u>Not met</u></p>	
----	---	-----------------------	--

Cruzan Mesa Vernal Pools SEA

General location and landform

The Cruzan Mesa Vernal Pools SEA lies north of the Santa Clara River southeast of Bouquet Canyon. The SEA boundaries encompass the watershed and drainages of the Cruzan Mesa and Plum Canyon vernal pools, which are considered as a single ecosystem within the SEA. Vernal pools are a rare habitat. They are floored by clay or other substrates that retain the water of spring rains for a period of weeks to months. The pools dry primarily by evaporation. Specialized fauna and flora are adapted to this regimen, and often form propagules that can last through years of desiccation. The pools are ephemeral, and may not fill every year. The fauna may be specific to an area of pools. Most biota is spread from pool to pool in mud, which adheres to the feet of birds and other animals that visit the pools.

The SEA is located within the unincorporated area of the County and lies entirely within the United States Geological Survey (USGS) 7.5' California Mint Canyon Quadrangle.

Biological Resources

Vegetation and communities

The SEA includes mesas, canyons and interior slopes, with Plum Canyon Creek running east-west through the southern portion of the SEA. The extent of the SEA encompasses the watershed, which supports both regionally unique vernal pools, including their immediate watersheds and the corridor between them. Plum Canyon forms the major drainage running east-west through the southern portion of the SEA, which drains west toward Bouquet Canyon. Uplands within the SEA are comprised of slopes and canyons, which support coastal sage scrub or scrub-chaparral vegetation. The Cruzan Mesa vernal pool complex lies within an elevated, topographically enclosed basin atop an eroded foothill between Mint and Bouquet canyons. Plum Canyon vernal pool, which is situated in a landslide depression on a hillside terrace, is smaller than the Cruzan Mesa pools, but possesses the same essential vernal pool characteristics as the larger system, and the two areas together form an ecologically functional unit.

This SEA area includes most of the local federally-designated critical habitat for the threatened vernal pool plant, spreading navarretia (*Navarretia fossalis*). The seasonally wet vernal pools and surrounding open coastal sage scrub and chaparral slopes support migrant and resident birds, other native sage scrub vertebrate species, and a number of sensitive taxa, including the County's only occurrences of both California Orcutt grass (*Orcuttia californica*) and Riverside fairy shrimp (*Streptocephalus woottoni*) and is one of a handful of known locales for western spadefoot toad (*Spea hammondi*). The steep cliffs that surround the Cruzan Mesa vernal pools, especially along the

DRAFT

southeast and north margins, provide protected sites for perching, roosting and nesting by a variety of birds of prey, including great horned owl, barn owl, and red-tailed hawk. The Plum Canyon vernal pool is hidden from view from the Plum Canyon roadway, and receives little human attention. Trash dumping, shooting and off-road vehicle activities have occurred within a few meters of the margin of the pool basin, but the pool shows only limited evidence of human intrusion.

Wildlife Movement

This SEA is located between the urban boundary of Santa Clarita and the vast undeveloped Sierra Pelona, and thus allows lateral movement for wildlife between lowland and montane environments. The vernal pools situated within this SEA serve as isolated, high-quality, habitat linkage sites for migratory waterfowl. The vernal pools teem with arthropod and amphibian activity, and provide essential feeding grounds for local species of reptiles, birds and mammals in an otherwise arid environment.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE CRUZAN MESA VERNAL POOLS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>Riverside fairy shrimp</u> – <u>California Orcutt grass</u> – <u>Spreading navarretia</u> – <u>Western spadefoot toad (sensitive; limited local occurrences)</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>Regionally unique vernal pool ecosystem (Cruzan Mesa and Plum Canyon)</u> – <u>Vernal pool flora and fauna restricted to seasonal clay basin habitats</u> – <u>County’s only known occurrences of Riverside fairy shrimp and California Orcutt grass</u> – <u>Federally designated critical habitat for spreading navarretia</u>

DRAFT

Criterion		Status	Justification
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Ephemeral vernal pools supporting breeding amphibians and invertebrates</u> – <u>Seasonal wetlands providing feeding habitat for migratory waterfowl</u> – <u>Concentrated breeding habitat for vernal pool-dependent species</u> – <u>Limited availability of vernal pool habitat within Los Angeles County</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Vernal pools occurring in isolated clay basin atop mesa landform</u> – <u>Unusual hydrologic regime (seasonal inundation and desiccation)</u> – <u>Highly specialized biota adapted to intermittent wetland conditions</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>One of few remaining vernal pool complexes in Los Angeles County</u> – <u>Intact vernal pool watershed supporting functional ecosystem processes</u> – <u>Limited human disturbance at Plum Canyon vernal pool basin</u> – <u>Preserved examples of rare seasonal wetland habitat</u>

Criterion		Status	Justification
G)	Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.	Met	<ul style="list-style-type: none"> – Located between Santa Clarita urban edge and Sierra Pelona open space – Lateral movement between lowland and montane environments – Vernal pools serving as linkage sites for migratory waterfowl – Connectivity along Plum Canyon drainage

Eastern San Gabriel Foothills SEA

General location and landform

The Eastern San Gabriel Foothills SEA (formerly the San Dimas Canyon and San Antonio Wash SEA), extending south from the larger San Gabriel Mountains SEA to the northern edge of urban development in the eastern San Gabriel Valley, includes approximately 12,353 acres in the Angeles National Forest, the unincorporated area of the County, the City of Claremont, the City of Glendora, the City of La Verne, and the City of San Dimas. From San Antonio Wash, which runs along the County line and is tributary to the Santa Ana River, the SEA extends westward through the Padua Hills/Claremont Wilderness Park, incorporating the lower parts of Williams, Burbank, Live Oak, Marshall, San Dimas, Ham, Sycamore, Shuler, Spring, Wildwood, Morgan, Mull, and Big Dalton canyons, all of which are tributary to the San Gabriel River.

The SEA is found within the Glendora, Mount Baldy, San Dimas, and Ontario U.S. Geological Survey (USGS) 7.5' California Quadrangles.

Biological Resources

Vegetation and communities

With elevations ranging between 1,200 and 4,400 feet above MSL, the topography tends toward the severe, especially in the northern part of this SEA, which includes steep-walled canyons and narrow ridgelines. The varied topography and geology of the Eastern San Gabriel Foothills SEA have allowed for development of diverse natural communities that include grasslands, shrublands, riparian woodlands, and forests, with many sub-associations represented. This area contains the last remaining relatively well-developed lower montane riparian habitat in eastern Los Angeles County. Dammed drainages have created significant reservoirs or flood control basins in the SEA.

The San Dimas Canyon watershed is part of the Experimental Forest section of the Angeles National Forest. Experiments conducted here during the latter half of the 20th century helped to elucidate the relationships among rainfall, topography, vegetation, and runoff, which influenced flood control practices in the Los Angeles Basin and other areas of the U.S. The area was carefully protected

DRAFT

through very limited and monitored access. The terrain chiefly includes undisturbed natural habitats of rocky canyon walls and canyon forest, riparian areas, coniferous and oak forest, chaparral, and grassland. Many years after the experimental removal of woody vegetation from selected slopes, some of those areas remain grassland.

Unusual soil types in the Eastern San Gabriel Foothills SEA include the Glendora Volcanics, which provide habitat for many-stemmed dudleya (*Dudleya multicaulis*), and clay inclusions that provide habitat for the County's only populations of the State- and federally listed thread-leaved brodiaea (*Brodiaea filifolia*).

Downstream of the San Antonio Dam, and extending onto the adjacent alluvial plain, is the best example of arroyo or wash vegetation remaining in the County. The vegetation is a dry form of coastal sage scrub, with included desert plants that are adapted to coarse substrate. The vegetation is much more dense and stable than that growing on alluvial fans behind Santa Fe Dam (San Gabriel Canyon SEA) and Hansen Dam (Tujunga Valley-Hansen Dam SEA).

The foothills to the west support not only thread-leaved brodiaea, but also the federally and state-listed Nevin's barberry (*Berberis nevinii*), the federally threatened coastal California gnatcatcher, and numerous other special-status species, including San Gabriel oak (*Quercus durata* var. *gabrielensis*), Engelmann oak (*Q. engelmannii*), oak hybrids, coastal cactus wren, patch-nosed snake, two-striped gartersnake, San Gabriel chestnut and Trask shoulderband.

Wildlife Movement

Extensive riparian, oak, and walnut woodlands/forests in this SEA constitute important stopover and overwintering areas for a wide variety of migratory birds. The major canyons provide conduits for wildlife to move north and south between foothill areas and the extensive wildlands of the San Gabriel Mountains SEA. This type of movement occurs on a seasonal and more frequent basis, particularly for large mobile mammals, such as American black bear, mountain lion, coyote, bobcat, and mule deer, whose full range of habitat needs are typically met over broad areas. Preserving a band of natural habitat along the lower-elevation foothills facilitates east/west movement for many terrestrial species.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

DRAFT

CRITERIA ANALYSIS SUMMARY OF THE EASTERN SAN GABRIEL FOOTHILLS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>Coastal California gnatcatcher</u> – <u>Thread-leaved brodiaea</u> – <u>Slender mariposa lily</u> – <u>San Gabriel linanthus</u> – <u>Greata’s aster</u> – <u>Golden eagle</u> – <u>Mountain lion (Southern California/Central Coast ESU)</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>Lower montane riparian habitat (last well-developed example in eastern Los Angeles County)</u> – <u>Arroyo/wash vegetation downstream of San Antonio Dam</u> – <u>Walnut woodland and oak riparian woodland</u> – <u>Alluvial fan scrub and coastal sage scrub mosaic</u> – <u>Habitat associated with Glendora Volcanics and clay inclusions (localized endemic flora)</u> – <u>Concentration of foothill-restricted species (e.g., coastal cactus wren, coast range newt)</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Not Met	
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Not Met	

DRAFT

Criterion		Status	Justification
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Extensive relatively undisturbed foothill canyon habitats</u> – <u>Intact lower montane riparian and woodland communities</u> – <u>Undisturbed experimental forest landscapes in San Dimas Canyon</u> – <u>Preserved foothill mosaic of forest, chaparral, and grassland</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>North-south linkage between foothills and San Gabriel Mountains</u> – <u>East-west movement corridor along lower foothill band</u> – <u>Canyon corridors facilitating wildlife movement between habitats</u> – <u>Documented movement of wide-ranging mammals (mountain lion, black bear, mule deer, bobcat)</u>

East San Gabriel Valley SEA

General location and landform

The East San Gabriel Valley SEA is located in the eastern end of the San Gabriel Valley, between the Puente Hills and the base of the San Gabriel Mountains (“Glendora Foothills”). The SEA includes some of the most extensive low-elevation habitat in the County, and is comprised of undeveloped ridgelines, hilltops and drainages with a mix of oak and walnut woodland, coastal sage scrub and riparian vegetation.

Generally, the topography within this SEA consists of moderate to steep hillsides with north, south, east and west slope aspects. Ridgelines vary in width, from narrow to broad, with well-defined drainages in between. One major undeveloped drainage, Walnut Creek, and a large, marsh-rimmed man-made reservoir, Puddingstone Reservoir, are found within this SEA. Elevations are very low for open space in the County, ranging from a low of approximately 560 feet above Mean Sea Level (MSL) in the Walnut Creek drainage, to a high of approximately 1,375 feet above MSL at Buzzard Peak.

Broadly, the open space within this SEA may be thought of as six separate, but ecologically connected,

DRAFT

units. These include 1) South Hills Park and the surrounding undeveloped land in the City of Glendora north of Interstate-210; 2) The natural riparian section of Walnut Creek County Park and, disjunct from this drainage by residential development, undeveloped hillsides in the Via Verde area of San Dimas; 3) Frank G. Bonelli Regional County Park (Bonelli Park) and adjacent open space; 4) Buzzard Peak and associated undeveloped open space mainly within, the cities of Walnut and West Covina; 5) Elephant Hill and an adjoining ridgeline in the City of Pomona; and 6) Parkland and other undeveloped habitat within City of Diamond Bar.

In its entirety, the SEA is located within the San Dimas, Glendora and Baldwin Park U.S. Geological Survey (USGS) 7.5' California Quadrangles. The SEA encompasses several different local jurisdictions, including the unincorporated area of the County, City of Covina, City of Glendora, City of La Verne, City of Pomona, City of San Dimas, City of Walnut, and City of West Covina. Diamond Bar is recommended for inclusion within this SEA, as it supports some of the most intact and significant natural habitats in the area.

Biological Resources

Vegetation and communities

Large parts of this SEA are designated critical habitat for the federally-threatened coastal California gnatcatcher (*Polioptila californica californica*), restricted to the remaining coastal sage scrub habitat. Other associated wildlife species include an isolated coastal population of cactus wren. More than a dozen regionally-rare species are found within this SEA, including some of the County's only breeding grasshopper sparrows, and extensive stands of southern California black walnut woodland.

The plant and wildlife community found on low hills within this SEA include species widely extirpated in the County, such as Hubby's phacelia and southern California legless lizard.

The natural areas of the east San Gabriel Valley serve as a migration and habitat connection between the San Gabriel Mountains and the Puente Hills, and large numbers of songbird migrants are encountered near the peaks and ridges of this SEA during spring migration.

This SEA may be thought of as a mosaic of interconnected, or nearly connected units, sharing similar habitat and resources:

Unit 1: The South Hills is an area of chaparral and grasslands, and serves as a stepping stone for wildlife (chiefly aerial, including birds and insects) that move along the spine of the San Jose Hills between the San Gabriel Mountains of the Transverse Ranges, and the Puente and Chino Hills of the Peninsular Ranges on the southern side of the San Gabriel Valley. In South Hills Park, there are fine intact stands of walnut woodland in the upper elevations; ravines with oak and walnut woodland in the upper elevations; what remains of the coast live oak woodland that once bordered the Big Dalton Wash; and extensive stands of prickly pear cactus covering some of the slopes. There are also grasslands (though most now have a dominant component of introduced mustards and other Mediterranean forbs) and some rocky outcrops with uncommon flora. Coastal sage scrub with dominant components of California sagebrush (*Artemisia californica*), sumacs (*Rhus* spp.) and elderberries (*Sambucus mexicana*) cover many of the slopes. Big Dalton Wash is channelized, and much development has been inserted into what was formerly Big Dalton's riparian oak woodland. This residential area forms a border for the northeast area of the SEA.

Unit 2: Walnut Creek runs along the northern base of the San Jose Hills and is a riparian interface between the rolling hill habitat and the coastal plain of the San Gabriel Valley. This segment

DRAFT

encompasses the undeveloped creek bottom and flood plain, which is a relatively natural area along and adjacent to Walnut Creek, most of it preserved as Walnut Creek Park. The Creek and lower slopes of the San Jose Hills contain one of the best riparian oak woodlands in the County, with a mixture of ash (mostly the introduced Shamel ash, *Fraxinus udhei*), walnut, willows, and sycamores. The upper areas of the oak woodland transition into excellent examples of walnut woodland.

Unit 3: Walnut Creek Park is connected to Bonelli Park unit at Interstate 210 where the drainage and the SEA underpass along with San Dimas Avenue. There is a fenced footpath on the south side of the underpass that wildlife and humans use extensively. The upper edges of Walnut Creek have business parks and industrial properties, as well as residential areas.

Bonelli Park includes the 250-acre Puddingstone Reservoir, which is a flood control basin created by damming the main course of Walnut Creek. Puddingstone Reservoir is home to many resident and migratory water birds, and its marsh and riparian edges support birds typical of these habitats in southern California, including sensitive species such as least Bell's vireo. Although subjected to boating activities and shoreline recreational use, Puddingstone Reservoir serves as an important habitat for breeding, wintering, and migrating water birds (including rookeries of herons, egrets, and double-crested cormorant).

Another species of note is the many-stemmed dudleya (*Dudleya multicaulis*), which is an uncommon plant found on the picturesque rocky outcrops of the Glendora volcanics formation. This plant is at its northernmost extent in this area of the San Gabriel Valley, and is more common in Orange County.

Unit 4: Buzzard Peak represents the highest elevation of a natural area that extends the habitats of Bonelli Park westward (i.e., across State Route 57). Over the past two decades, much of the walnut woodland and coastal sage scrub here has been set aside as designated open space, either via easements following residential development, or as protected lands purchased by land conservancies. Thus, while some areas are still threatened by expansion of residential and commercial development, the majority of undeveloped land here is now parkland, critical for the maintenance of biodiversity in the east San Gabriel Valley.

Unit 5: Elephant Hill, within the City of Pomona, and the adjacent undeveloped ridge extending southeast across Mission Blvd. is bordered chiefly by development but has retained an undisturbed remnant of the original habitat: north-facing slopes of oak and walnut woodland. South-facing slopes consist of a mixture of dispersed chaparral, coastal sage scrub, and grasslands. The ravines in the area have fine examples of walnut woodland and the upper slopes have grassland, often dominated by mustards, with some areas of prickly-pear scrub.

Unit 6. The final unit in this SEA falls within the city limits of Diamond Bar and supports similar habitat to the others (walnut and oak woodland, cactus-rich coastal sage scrub, etc.). However, this unit is notable due to its adjacency to the extensive open space of upper Tonner Canyon / Dos Hermanos Ranch across the San Bernardino County line. This area supports an extremely high diversity of plant and particularly wildlife species otherwise extirpated from the Los Angeles area, such as breeding golden eagle, grasshopper sparrow, and tricolored blackbird. Because so much of the habitat in the area is private, or otherwise inaccessible, the true ecological significance has yet to be fully explored.

Wildlife Movement

The SEA represents the only feasible wildlife linkage opportunity between the Transverse and Peninsular Ranges within the County via the San Gabriel Mountains and the Puente-Chino Hills. Unlike a traditional wildlife corridor, however, this SEA contains a series of discontinuous (or tenuously

DRAFT

connected) habitat blocks and patches, rather than an unbroken corridor for movement. This SEA facilitates movement and exchange between larger habitat areas by permitting terrestrial "island-hopping" between the SEA components.

Recent research has shown that this SEA represents a key connectivity zone for golden eagle movement, and years of repeated observation suggests that a variety of migrant songbirds (in addition to raptors) move through the SEA, touching down as weather or food conditions warrant.

This same function probably does not apply to other taxonomic groups. It is highly doubtful that amphibian, reptile and most mammal populations use this corridor as effectively as birds, if at all. Mule deer (*Odocoileus hemionus*), for example, do not occur within Bonelli Park, but are common in the San Gabriel Mountains and the Puente Hills. However, some mammals that tolerate urban environments, such as Virginia opossum (*Didelphis virginiana*), raccoon (*Procyon lotor*), and striped skunk (*Mephitis mephitis*), use the corridor in the manner previously described. Even bobcat (*Lynx rufus*) and mountain lion (*Puma concolor*) periodically enter Bonelli Park and Walnut Creek Park from the outside, by way of routes related to SEA components. The localized western gray squirrel (*Sciurus griseus*) maintains small populations within this SEA, where it manages to persist in patches of oak and walnut woodland, and within larger riparian corridors.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE EAST SAN GABRIEL VALLEY SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none">- <u>Coastal California gnatcatcher critical habitat</u>- <u>Multi-stemmed dudleya (northern range extent)</u>- <u>Intermediate mariposa lily</u>- <u>Sensitive coastal sage-scrub-dependent species</u>

DRAFT

Criterion		Status	Justification
B/C)	<p><u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u></p>	Met	<ul style="list-style-type: none"> – <u>Extensive southern California black walnut woodland</u> – <u>Low-elevation coastal sage scrub largely extirpated elsewhere in County</u> – <u>Isolated coastal cactus wren population</u> – <u>Occurrences of Hubby’s phacelia and southern California legless lizard</u> – <u>Regionally rare grasshopper sparrow breeding habitat</u>
D)	<p><u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u></p>	Met	<ul style="list-style-type: none"> – <u>Puddingstone Reservoir supporting breeding, wintering, and migratory water birds</u> – <u>Migratory songbird stopover habitat along ridgelines and peaks</u> – <u>Riparian oak woodland along Walnut Creek providing breeding and foraging habitat</u> – <u>Seasonal concentration areas for raptors and waterbirds</u>

DRAFT

Criterion		Status	Justification
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Northern range limit for several Peninsular Range species</u> – <u>Multi-stemmed dudleya at northernmost extent in San Gabriel Valley</u> – <u>Low-elevation open space uncommon within Los Angeles County</u> – <u>Transitional habitat between Transverse and Peninsular Ranges</u>
E)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Remnant walnut and oak woodland on undeveloped ridgelines</u> – <u>Relatively intact riparian oak woodland along Walnut Creek</u> – <u>Mosaic of coastal sage scrub and chaparral on minimally disturbed slopes</u> – <u>Preserved patches of historic low-elevation habitats</u>

DRAFT

	Criterion	Status	Justification
G)	<p><u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u></p>	Met	<ul style="list-style-type: none"> – <u>Only feasible linkage between San Gabriel Mountains and Puente-Chino Hills</u> – <u>Island-hopping connectivity across discontinuous habitat blocks</u> – <u>Movement pathway between Transverse and Peninsular Ranges</u> – <u>Connectivity zone for golden eagle movement</u> – <u>Riparian and ridgeline linkages facilitating bird migration</u> – <u>Terrestrial movement through Walnut Creek and Bonelli Park system</u>

El Dorado Regional Park Lakes SEA

General location and landform

El Dorado Regional Park, covering more than 600 acres in the City of Long Beach, features a series of natural lakes historically associated with swamps of the lower San Gabriel River, which were extensive prior to its channelization in the mid-twentieth century. The 135 acres of the park designated as SEA include six hydrologically connected lakes in Areas 1, 2, and 3 of the park, which constitute rare and especially valuable freshwater wetland habitats in coastal Los Angeles County. The surrounding developed parts of the park are not recommended for SEA designation.

The SEA is located within the Downey United States Geological Survey (USGS) 7.5' California Quadrangle.

Although not a formal SEA Selection Criterion, it is relevant that wetlands historically associated with the San Gabriel River, including those in El Dorado Regional Park, are considered crucial to the Gabrieleño-Kizh/Tongva and Juaneño/Acjachemen peoples, for whom the permanent coastal settlement of Puvunga, including Motuucheyngna, is recognized as a major, sacred site of emergence. As such, formally recognizing the ecological significance of these natural lakes and connecting waterways through SEA designation would be consonant with and respectful of TEK.

DRAFT

Biological Resources

Vegetation and communities

Considered as a whole, El Dorado Regional Park is known to provide important habitat for birds, with eBird records documenting more than 240 species. Along with large numbers of resident waterfowl, ducks found at the lakes during fall and winter include American wigeon (*Mareca americana*), gadwall (*M. strepera*), northern shoveler (*Spatula clypeata*), and ruddy ducks (*Oxyura jamaicensis*). Three California Species of Special Concern, the least bittern (*Botarus exilis*), Clark's marsh wren (*Cistothorus palustris clarkae*), and northern yellow warbler (*Setophaga aestiva*), nest in and around the lakes. Other wetland-dependent nesting species include great blue (*Ardea herodias*) and green (*Butorides virescens*) herons, and tree swallows. Lucky morning-glory (*Calystegia felix*), a species thought to be extinct until being rediscovered in Chino, San Bernardino County, in 2011, has since been found at Area 1 of El Dorado Regional Park (the Nature Center), where a few small populations persist as reminders of the park's heritage as part of an extensive natural swamp.

The lakes support some extensive stands of emergent freshwater marsh habitat characterized by native California bulrush (*Schoenoplectus californicus*) and broad-leaf cattail (*Typha latifolia*), mixed with smaller amounts of various herbs, such as salt marsh fleabane (*Pluchea odorata*). Native trees growing around the margins of some of the lakes include black and arroyo willow (*Salix gooddingii*, *S. lasiolepis*).

Wildlife Movement

El Dorado Regional Park is contiguous with the paved channel of the lower San Gabriel River. This channel and that of the Los Angeles River, several miles to the west, are the only geographic features connecting the southern coast of Los Angeles County to SEAs and other natural areas located 15 or more miles inland.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE EL DORADO REGIONAL PARK LAKES SEA

<u>Criterion</u>	<u>Status</u>	<u>Justification</u>
A) <u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	- <u>Lucky morning-glory</u>

DRAFT

Criterion		Status	Justification
B/ C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> - <u>Freshwater marsh habitat</u> - <u>Multi-lake freshwater wetland complex</u> - <u>California bulrush-cattail marsh</u> - <u>Least bittern</u> - <u>Clark's marsh wren</u> - <u>Northern yellow warbler,</u> - <u>Wetland-dependent nesting herons</u> - <u>Rediscovered lucky morning-glory population</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> - <u>Wintering waterfowl concentrations</u> - <u>Nesting waterbird rookeries</u> - <u>Breeding marsh birds</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Not met	
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Not Met	
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Coastal-inland wetland linkage along San Gabriel River</u> - <u>Freshwater wetland stepping-stone along lower San Gabriel River</u>

Griffith Park SEA

General location and landform

The Griffith Park SEA is located at the extreme eastern end of the Santa Monica Mountains. The SEA

DRAFT

is an extensive, relatively undisturbed island of natural vegetation in an urbanized, metropolitan area, connected to the rest of the Santa Monica Mountains via highway overpasses along State Route 101 through Cahuenga Pass. It supports the coastal sage scrub, chaparral, riparian, and southern oak woodland plant communities that are typical in the foothills and valleys of southern California. For nearly a century, Griffith Park has been an island of natural vegetation surrounded by urban and suburban development. The geographic location makes the area important for scientific study, for genetic interchange between otherwise isolated populations, and for recreation of urban residents.

The SEA is located partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Burbank and Hollywood.

Biological Resources

Vegetation and communities

The SEA encompasses most of Griffith Park and the open space surrounding the Hollywood Reservoir, bounded by State Route 101, State Route 134, and Interstate-5. Griffith Park is the easternmost extent of the Santa Monica Mountains, and a stepping stone to the Verdugo and San Gabriel Mountains, the latter only two miles distant. It is a very important natural area for animals and plants species that occur in both the Santa Monica and San Gabriel mountains. Because of its large acreage, Griffith Park maintains relatively large populations of biological communities given its isolation, even including top predators, such as bobcats and (occasionally) mountain lions. The life zones of Griffith Park are somewhat compressed, with foothill species such as Eastwood's manzanita and interior live oak occurring at a relatively low elevation. Many—probably dozens—of plant species found in Griffith Park are absent elsewhere in the eastern Santa Monica Mountains and on the floor of the Los Angeles Basin, only appearing in the western Santa Monicas and Verdugo Mountains.

Griffith Park has extensive areas of coastal chaparral and is an island of refuge for native animals in the Santa Monica Mountains. Its mosaic of habitats includes coastal sage scrub, riparian areas, and southern oak woodland. The mosaic of habitats is especially valuable to preserve. Griffith Park is in the City of Los Angeles and protected in this respect, but no approved and implemented management plan preserves its natural habitat in perpetuity.

The SEA includes patches of remnant coast live oak woodland (*Quercus agrifolia*), which has many oak trees, and in many places retains its natural understory and component species, including a large population of acorn woodpecker (*Melanerpes formicivorus*), and scarce amphibians such as arboreal salamander (*Aneides lugubris*). In addition, birds rely on large urban islands of habitat for areas to rest and feed along their migration. In the case of Griffith Park, this function is more significant than might be expected because it serves as a corridor for any gene flow and species movement that may take place between the Santa Monica and San Gabriel mountains via the Verdugo Mountains.

This habitat transitions as one moves uphill into chaparral, with patches of native grassland on clay soils (largely on the upper slopes, far from human intrusion). North-facing sides of rocky outcrops support "moss gardens"—seasonally wet vegetation that is characterized by multiple kinds of lichens, mosses, liverworts and other non-vascular plants along with lance-leaf live-forever (*Dudleya lanceolata*), and other flowering plants.

Multiple rare plants are known from the park and adjacent open space, including Plummer's mariposa-lily southern California black walnut, Hubby's phacelia, ocellated Humboldt lily, San Gabriel oak, and small-flowered morning-glory. Sensitive wildlife species include southern California rufous-crowned sparrow, southern California legless lizard, coastal whiptail, and coast horned lizard. For many of these

DRAFT

scrubland and chaparral species, populations in the park represent some of their only occurrences on the floor of the Los Angeles Basin.

Wildlife Movement

Griffith Park has long been isolated from the rest of the Santa Monica Mountain Range, the Los Angeles River and the Verdugo Mountains (a little less than two miles to the east) because of the freeways, concrete river projects and urbanization that surround the park. Although some species have disappeared, including the ringtail (*Bassariscus astutus*) and (likely) the western spotted skunk (*Spilogale gracilis*), the park supports sustaining populations of most other Santa Monica Mountains wildlife species, including mule deer, gray fox, bobcat, and at least occasionally, mountain lion. All of these species may potentially utilize road overpasses that cross Cahuenga Pass, providing access to the larger Santa Monica Mountains.

Although some stretches of the Los Angeles River may not provide suitable primary corridors, it is important to reinstate Griffith Park's connection to the Los Angeles River for the future of wildlife and plant connectivity. In the management draft for Griffith Park wildlife, the authors outline some of the important connections to maintain or enhance: bridges and underpasses over and under State Route 101 and culverts that feed into the Los Angeles River channel.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE GRIFFITH PARK SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	<u>Not Met</u>	<u>– No known core populations occur within this SEA.</u>

DRAFT

	Criterion	Status	Justification
B/C)	<p><u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u></p>	Met	<ul style="list-style-type: none"> – <u>Remnant southern California black walnut woodland</u> – <u>Ocellated Humboldt lily and Plummer’s mariposa lily occurrences</u> – <u>Coast horned lizard and coastal scrub-dependent fauna</u> – <u>Mosaic of coastal sage scrub, chaparral, riparian, and oak woodland</u> – <u>One of few large natural habitat blocks on Los Angeles Basin floor</u>
D)	<p><u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u></p>	Not Met	–
E)	<p><u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u></p>	Met	<ul style="list-style-type: none"> – <u>Easternmost extent of the Santa Monica Mountains</u> – <u>Foothill species occurring at unusually low elevations</u> – <u>Isolated urban habitat island supporting compressed life zones</u> – <u>Basin-floor occurrences of species otherwise absent locally</u>

DRAFT

Criterion		Status	Justification
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Large urban island of relatively undisturbed native vegetation</u> – <u>Intact mosaic of coastal sage scrub, chaparral, and oak woodland</u> – <u>Remnant coast live oak woodland with natural understory</u> – <u>Preserved native habitat within heavily urbanized Los Angeles Basin</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Stepping-stone linkage between Santa Monica, Verdugo, and San Gabriel Mountains</u> – <u>Movement via Cahuenga Pass overpasses (State Route 101)</u> – <u>Urban refuge facilitating gene flow between mountain ranges</u> – <u>Connectivity potential to Los Angeles River corridor</u>

Harbor Lake Regional Park SEA

General location and landform

The Harbor Lake Regional Park SEA is located within Kenneth Malloy Harbor Regional Park in the Harbor City community of the City of Los Angeles, which is approximately 15 miles south of downtown Los Angeles and just west of the Interstate-110. It is bordered to the north by the Pacific Coast Highway, but includes a small segment of drainage, the Wilmington Drain (“Bixby Slough”) north of Pacific Coast Highway. The lake is named “Machado Lake.” The SEA supports some of the last remaining freshwater wetlands that once covered the southern and western areas of the Los Angeles Basin. The SEA supports significant areas of aquatic and terrestrial plant communities, which provide habitat to a variety of birds and wildlife.

DRAFT

The site is located in the Torrance Quadrangle of the United States Geological Survey (USGS) 7.5 Minute Map Series (USGS, 1964).

Biological Resources

Vegetation and communities

The SEA boundaries encompass the lake areas that contribute to the biological function of the Harbor Lake ecosystem. It includes a sizeable natural lake with a freshwater marsh along its northern, eastern and southern shores and the Bixby Slough, habitats that once covered the South Bay area. This type of habitat has been filled, drained, and lost to development throughout most of southern California. Numerous special-status bird species occur here, such as nesting least bittern, yellow-breasted chat (*Icteria virens*), Clark's marsh wren and yellow warbler (*Setophaga petechia*). The slough is a freshwater marsh/willow-riparian environment that supports a surprising diversity of wildlife. The freshwater marsh is a good example of the freshwater marshes that used to occur along the fault lines of the Los Angeles Basin and at the various river mouths. However, it is highly disturbed, having been subjected to a massive restoration effort (2014) that re-established the wetland and riparian habitat, particularly along the lake's eastern margin.

Still, the SEA supports an astonishing diversity of birds (nearly 300 species as of March 2026; <https://ebird.org/hotspot/L1552663>) and is considered an Audubon Important Bird area, though most species are present only briefly, generally in spring and fall migration. It encompasses a substantial expanse of remaining permanent and seasonal wetland habitat that once covered the South Bay/Harbor area. Most habitat of this type in the County has been filled, drained, and lost to development. Most of the bird species found here are dependent in some way on the surface moisture and vegetation, and would not be able to survive without it. It is also a habitat that supports several species of reptiles and amphibians, notably black-bellied slender salamander (*Batrachoseps nigriventris*) and isolated populations of California kingsnake (*Lampropeltis californiae*) and side-blotched lizard (*Uta stansburiana*). Western toad may be extirpated, based on a lack of recent observations despite records of other amphibian taxa in the area, suggesting some level of survey effort.

Least Bell's vireos have bred since at least 2018 (1 – 2 pairs), and the population of southern tarplant, located in the overflow area south of Machado Lake may be among the largest in the world, with thousands of individuals in good years (and potentially supporting more, with restoration). While a prior version of this document reported that this SEA contains habitat that supports a population of vernal pool fairy shrimp (*Branchinecta lynchi*), this population may not be extant and requires verification.

Wildlife Movement

The SEA does not lie within any identified terrestrial movement routes for wildlife. However, it is an important regional stop on the Pacific Flyway in southern Los Angeles County, supporting dozens of species of waterbirds and landbirds during spring, fall and winter.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

DRAFT

CRITERIA ANALYSIS SUMMARY OF THE HARBOR LAKE REGIONAL PARK SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>Least Bell’s vireo breeding population</u> – <u>Coastal California gnatcatcher (occasional occurrences)</u> – <u>Southern tarplant (large regional population)</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>Remnant freshwater marsh and willow-riparian habitat, one of few remaining freshwater wetland complexes in Los Angeles Basin</u> – <u>Least bittern nesting habitat</u> – <u>Yellow-breasted chat and Clark’s marsh wren occurrences</u> – <u>Northern yellow warbler in marsh-riparian vegetation</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in County.</u>	Met	<ul style="list-style-type: none"> – <u>Freshwater marsh supporting breeding waterbirds</u> – <u>Pacific Flyway stopover for migratory birds</u> – <u>Seasonal wetland feeding habitat for aquatic and terrestrial species</u> – <u>Permanent surface water supporting regionally limited wildlife habitat</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Not Met	
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Not Met	

Criterion		Status	Justification
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Not Met	– <u>Regional stopover site along Pacific Flyway (avian movement only)</u>

Joshua Tree Woodlands SEA

General location and landform

The Joshua Tree Woodlands SEA is located in the western portion of the Antelope Valley from the Antelope Valley California Poppy Reserve State Natural Reserve area west, in an unincorporated area of the County. The SEA is located primarily on the western Antelope Valley floor between the Tehachapi Mountains and the western San Gabriel Mountains. The topography of the SEA is extremely flat with the land sloping less than 200 feet across approximately five miles. The location and orientation of the SEA represents a matrix of remnant stands of Joshua tree woodland and other desert habitats among a patchwork of disturbed areas.

In its prior configuration, the SEA consisted of eight separate (and separated) units, seven of which are in close proximity to each other, with most located between the Kern-Los Angeles County line to the north, and the California Aqueduct and Fairmont Butte to the south. However, this current analysis recommends connecting these units, and extending the boundaries of this SEA out to include a) regenerating Joshua tree woodland on former agricultural lands; and b) additional adjacent desert habitats, many of which either provide additional habitat for Joshua tree-associated wildlife, or that serve as buffer habitats between agricultural and residential areas and Joshua tree woodland.

The SEA connects to the existing San Andreas SEA, the Antelope Valley California Poppy Reserve State Natural Reserve, Arthur B. Ripley Desert Woodland State Park, and the County’s George F. Bones Desert Pines Wildlife Reserve. Because many of these areas were not contiguous with one another, but clearly support high-quality wildlife habitat, we have made an effort to better connect these with other SEAs. An additional disjunct unit of this SEA that had been identified along Interstate 5 at the far northwestern corner of the Liebre Mountains is now part of the larger Liebre/Sierra Pelona SEA, as it is ecologically and physically connected to that area.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Neenach School, Fairmont Butte, Black Mountain, and Lebec.

Several broad washes are found within this SEA, with rich alluvial soils that become waterlogged following rainstorms. Since 2000, many of the former agricultural fields have become fields of photovoltaic panels to generate renewable energy, a land use that largely precludes wildlife movement through these properties. While some vegetation may be retained under the panels, the habitat is largely transformed to an industrial use, which has permanently altered the ecology of the region. This has only increased the value of the remaining habitat on undeveloped parcels, both for the desert and grassland species that rely on them, as for the connectivity opportunities (see below).

The California Aqueduct flows through this SEA as an open (though cement-lined) channel, and is an

important resource for migratory and wintering waterfowl.

Biological Resources

Vegetation and communities

This SEA encompasses many of the remaining old-growth stands of Joshua trees (*Yucca brevifolia*) on the west side of the Antelope Valley, as well as important sand sheet and wash communities dominated by saltbushes (*Atriplex* spp.). Significant areas of native wildflowers are also encompassed by this SEA, in many cases forming the understory of the Joshua tree woodland. Joshua tree woodland is a complex biological community of the gradual slopes of the higher elevation Mojave Desert and once covered much of the Antelope Valley. As Joshua trees nearly exclusively within the Mojave Desert, these Los Angeles County populations are at the western limit of the species' range. Because Joshua trees occur in areas that are easily developed for residences and agriculture, their habitat has become very fragmented in the County.

Fragmentation is a concern because the Joshua trees depend on a small moth for reproduction. Only two species of moth can successfully pollinate western Joshua trees, and in Los Angeles County, there is only the western Joshua tree moth (*Tegeticula synthetica*). The moth may have limited dispersal abilities, and the Joshua trees cannot reproduce from seeds without pollination from this particular moth. Cross pollination is regarded as essential to a species' genetic diversity, which is essential to adaptation to environmental change.

This part of the Antelope Valley is very important as a migratory flyway for birds, from small passerines to the larger raptors, such as the state-threatened Swainson's hawk (*Buteo swainsoni*) and turkey vultures (*Cathartes aura*). The Swainson's hawk maintains its only nesting areas in the county here, selecting trees along windbreaks between current or former agricultural parcels, or around homes ("ranchettes").

The Joshua trees within the SEA exhibit a variety of growth forms, from tall, spreading trees to more clumped, almost shrubby plants connected by rhizomes (formerly considered a distinct variety, *Yucca brevifolia* var. *herbertii*). These clumps can be clones, with individuals all sharing the same genetic identity. The Joshua tree woodland with many shrub components of the biological community intact, including a floor covered by slender goldfields (*Lasthenia gracilis*) and a high diversity of other wildflowers in the spring.

The County's last remaining population of burrowing owls, now reduced to scattered individuals rather than colonies (which occurred historically), are found here. The Bell's sparrow is a very common resident within this SEA, with the localized Mojave Bell's sparrow (ssp. *canescens*) resident, along with many Mojave Desert species near the western edge of their range, such as ladder-backed woodpecker. For some threatened species, such as American badger, uncertainty about their specific range and population within the County makes any assessment of regional importance impossible (for now). Likewise, the taxonomy of certain rare species has not been fully worked out (e.g., which legless-lizards are found here) Other localized species here include ferruginous hawk, LeConte's thrasher, Scott's oriole and a variety of desert-associated reptiles, but their current distributions, in light of massive land use transformation for industrial-scale solar and other projects, are now poorly understood.

Wildlife Movement

Due to significant barriers, including Hwy. 138 and the California aqueduct, wildlife movement within

DRAFT

the SEA is likely limited to local movement. Biogeographical boundaries may also play a role, as the Liebre/Sierra Pelona are ecologically separate from the Tehachapi Mountains just to the north (i.e., many species reach their distributional limits in one range or the other). One exception would be the pronghorn (*Antilocapra americana*), a reintroduced population of which extends east from Tejon Ranch. For desert species, movement across the Antelope Valley floor is probably much facilitated by the large patches of Joshua tree habitat as island-like stepping stones. Birds, and possibly bats, and other aerial organisms that use the habitat linkage along the desert side of the San Gabriel Mountains probably use the woodland in the SEA for resting and feeding. The expanded boundaries we propose for this SEA improve connections to both the San Andreas SEA to the south, as well as the Antelope Valley SEA to the east. Given the rapid expansion of industrial solar facilities—as well as the looming threat of massive data centers on remaining flat, undeveloped land near transmission lines (which describes most of the current Antelope Valley)—decisions on land use are now even more urgent than at any time in the region’s history.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE JOSHUA TREE WOODLANDS SEA

Criterion		Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	<ul style="list-style-type: none"> – Joshua tree woodland – Swainson’s hawk (only nesting areas in Los Angeles County) – Burrowing owl (last remaining County population) – Tricolored blackbird (wintering use) – Alkali mariposa lily
B/C)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	<ul style="list-style-type: none"> – Westernmost extent of Joshua tree woodland in Los Angeles County – Saltbush scrub supporting LeConte’s thrasher – Grassland and fallow agricultural habitats supporting raptors and shorebirds – Mojave Desert species near western range limit (e.g., ladder-backed woodpecker) – Remnant Mojave Desert plant communities within Antelope Valley

DRAFT

Criterion		Status	Justification
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in County.</u>	Met	<ul style="list-style-type: none"> – <u>Swainson’s hawk nesting in windbreak trees</u> – <u>Migratory bird flyway across western Antelope Valley</u> – <u>Breeding habitat for desert-associated passerines</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Westernmost extent of Joshua tree woodland supports many desert species at the far western edge of their range, such as Scott’s oriole, ladder-backed woodpecker, and many others.</u> – <u>Distinct Joshua tree growth forms including clonal stands</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Remnant old-growth Joshua tree woodland</u> – <u>Intact Joshua tree understory with native wildflowers</u> – <u>Preserved sand sheet and desert wash communities</u> – <u>Representative Mojave Desert vegetation mosaic</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Stepping-stone patches of Joshua tree woodland across Antelope Valley</u> – <u>Linkage to San Andreas SEA and Antelope Valley SEA</u> – <u>Resting and feeding habitat for migratory birds along desert foothills</u> – <u>Connectivity among remnant desert habitats fragmented by development</u> – <u>Woodland patches facilitating movement of desert species</u>

Liebre-Sierra Pelona SEA

General Location and Landform

Located in the northwestern portion of the County, Liebre Mountain and the Sierra Pelona extend the Transverse Range west toward the Coast Ranges that begin in Ventura County.

DRAFT

Together with the San Gabriel Mountains to the east, most of the land here is owned by the federal government and managed by the U.S. Forest Service as four separate Districts comprising Angeles National Forest. The Forest boundary encompasses roughly 700,000 acres of Los Angeles County's 3 million total acres, stretching from the far northwestern corner near Gorman, southeast to Claremont at the San Bernardino County line (the San Gabriel Mountains extend several miles east into San Bernardino Co.). The Forest consists of two large, unconnected blocks of territory, bisected by S.R. 14 and the Santa Clara River. These mountains contain most of the County's undeveloped open space, and nearly all its large roadless areas. Private lands with clusters of houses are limited in the Liebre-Sierra Pelona, but the communities of Green Valley and Bouquet Canyon are exceptions.

Like the San Gabriel Mountains, the Liebre-Sierra Pelona supported a rich Indigenous history, with native peoples taking full advantage of the available game and abundant fruits such as acorns (*Quercus* spp.), chia (*Salvia columbariae*), and manzanita (*Arctostaphylos* spp.). Trading routes and footpaths for seasonal movements were extensive, connecting different portions of the mountains, and villages, with the lowlands to the north and south.

Biological Resources

Vegetation and communities

Like the San Gabriel Mountains to the east, the vegetation of the Liebre-Sierra Pelona is exceptionally diverse. The peaks are lower in this range (none exceeding 6,000', vs. several over 9,000' in the San Gabriel Mountains), with more oak woodland and less truly montane habitat. As in the San Gabriels, small sag ponds forming marshy meadows are located along the northern base of the range, but just one main watershed dominates (Santa Clara River). Many Californian tree and shrub species typical of the Sierra foothills reach the southern extent of their global range within the Sierra Pelona, including California buckeye (*Aesculus californica*), gray pine (*Pinus sabiniana*), blue oak (*Quercus douglasii*), and foothill gooseberry (*Ribes quercetorum*). The oak (*Quercus*) species diversity in particular is exceptionally high here, as the range represents a crossroads of Sierran, Coast Range, Mojave, and southern California woodland flora. Finally, interesting desert extensions of species are frequently encountered here, with populations of western Joshua tree (*Yucca brevifolia*) and Scott's oriole (*Icterus parisorum*) found amid oak woodland and alluvial scrub habitat toward the northwestern corner of the range.

While most of the Threatened and Endangered high montane species of the San Gabriel Mountains are absent in the Liebre-Sierra Pelona, this range supports significant populations of arroyo toad (*Anaxyrus californicus*), California red-legged frog (*Rana draytonii*), California condor (*Gymnogyps californianus*), California spotted owl (*Strix occidentalis occidentalis*), and golden eagle (*Aquila chrysaetos*). Unarmored threespine stickleback (*Gasterosteus aculeatus williamsoni*) historically occurred and have been reintroduced to San Francisquito Canyon. County- and regionally rare bird populations of the Liebre-Sierra Pelona include likely nesting belted kingfisher (*Megaceryle alcyon*) in the upper Piru Creek drainage near the Ventura Co. line.

While the San Gabriel Mountains support more State and Federally protected plant species, Ross pitcher sage (*Lepechinia rossii*) is apparently endemic to the Liebre-Sierra Pelona, which support many regionally-rare plants, and those with small global ranges, including Robbin's nemacladus (*Nemacladus secundiflorus* var. *robbinsii*), Tehachapi monardella (*Monardella linoides* ssp. *oblonga*), Johnson's monkeyflower (*Diplacus johnstonii*), Mount Pinos larkspur (*Delphinium parryi* ssp. *purpureum*), and Peirson's morning-glory (*Calystegia peirsonii*).

DRAFT

The nesting raptor community of the Liebre-Sierra Pelona appears to be highly diverse, with historical nesting sites for California condor (Red Mountain), and breeding-season records of some of the County’s rarest nesting raptors, including prairie falcon (*Falco mexicanus*), long-eared owl (*Asio otus*), and northern saw-whet owl (*Aegolius acadicus*).

Wildlife Movement

Like the San Gabriel Mountains to the east, the Liebre-Sierra Pelona is a largely undeveloped, largely roadless mountain range bordering a megacity to the south. It may likewise be considered a “source” habitat for wide-ranging species such as mountain lion (*Puma concolor*) and golden eagle (*Aquila chrysaetos*) in the County. The freeways that border and cross the range, including Interstate 5 and S.R. 14, are thus of major concern for wildlife movement, and planning is underway to provide and improve wildlife crossing opportunities for these species in the form of underpasses and wildlife bridges.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE LIEBRE-SIERRA PELONA SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> - Arroyo toad - California red-legged frog - California condor - California spotted owl - Golden eagle - Ross pitcher sage (endemic)
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> - Diverse oak woodland assemblages - Regionally rare montane plants - California condor breeding use - Diverse nesting raptor community - Desert-montane vegetation overlap - Joshua tree occurrences in oak woodland
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> - Aquatic breeding habitat for arroyo toad and red-legged frog - Headwater stream habitats for native fish - Nesting raptor habitat in montane terrain

DRAFT

Criterion		Status	Justification
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> - <u>Southern range limits for Sierran woodland species</u> - <u>Exceptional oak species diversity</u> - <u>Desert-montane floristic overlap</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> - <u>Extensive relatively undisturbed montane habitats</u> - <u>Large contiguous oak woodland and chaparral</u> - <u>Intact watershed-scale habitat</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Regional source habitat for wide-ranging species</u> - <u>Elevational movement corridors</u> - <u>Large contiguous undeveloped mountain habitat</u>

Lower Los Angeles River SEA

General Location and Landform

Extending approximately six linear miles through the City of Long Beach, from Pacific Coast Highway north to Artesia Boulevard, and encompassing approximately 423 acres, the highly modified ecosystem of the lower Los Angeles River provides important habitat for a wide range of birds, and especially migratory shorebirds. The recommended SEA consists of the channel itself, administered by the Los Angeles County Flood Control District (LACFD), and associated restored wetlands of the Dominguez Gap (52 acres, restored by the LACFD in 2008) and DeForest Park (34 acres, restored by the City of Long Beach in 2018).

The SEA is located within the Downey United States Geological Survey (USGS) 7.5' California Quadrangle.

Although not a formal SEA Selection Criterion, it is relevant that the Gabrieleño-Kizh/Tongva people regard the Los Angeles River—Paayme Paxaayt (“west river”) or Wenot (“the life-giver”)—as a centerpiece of trade, culture, and spirituality. Historically, at least 45 villages flourished along its length, including Ahwaanga near the river’s mouth. As such, formally recognizing the ecological significance of the lower Los Angeles River and associated wetlands through SEA designation would be consonant with and respectful of TEK.

Biological Resources

Vegetation and communities

The recommendation to designate this area mainly reflects its unique value as a crucial stopover site for large numbers of shorebirds during fall migration, as summarized by Cooper (2006, p. 1): “The concrete-lined channel of the lower Los Angeles River in Long Beach supports one of the largest concentrations of shorebirds in southern California during fall migration. Each fall, the black-necked stilt (*Himantopus mexicanus*), western sandpiper (*Calidris mauri*), and least sandpiper (*Calidris minutilla*) occur in the thousands of individuals per day, and the American avocet (*Recurvirostra americana*) and long-billed dowitcher (*Limnodromus scolopaceus*) occur in the hundreds. Numbers peak from mid-August through late September, and eight systematic surveys between 9 August and 27 September 2000 yielded in excess of 14,000 birds on four of these counts. Despite the channel’s proximity to the coast, the species composition resembles that at other inland sites in the Southwest more closely than natural coastal wetlands in the region. Because of the channel’s constricted concrete borders, even slight rainfall can raise the water level in the channel, reducing the suitability of the habitat for shorebirds. This effect likely accounts for the area’s relatively light use by shorebirds outside the late summer and fall dry season.”

The Dominguez Gap Wetlands occupy two areas between the 405 freeway and Del Amo Boulevard: a 37-acre basin on the eastern side of the river channel and an 18-acre basin on the west side of the channel. The DeForest Wetlands occupy a basin 2.4 miles long on the eastern side of the river channel, from Del Amo Boulevard to Osgood Street. A review of eBird records show multiple summer records of least bitterns (*Botaurus exilis*), Clark’s marsh wrens (*Cistothorus palustris clarkae*), and northern yellow warblers (*Setophaga aestiva*), suggesting localized nesting by these California Species of Special Concern in the Dominguez Gap Wetlands and possibly in the DeForest Wetlands. Other wetland-dependent nesting species include green heron (*Butorides virescens*) and tree swallow (*Tachycineta bicolor*).

The river channel itself does not support natural communities, but the Dominguez Gap Wetlands and DeForest Wetlands have been restored using a combination of riparian trees, including western sycamore (*Platanus racemosa*), Fremont cottonwood (*Populus fremontii*), and willows (*Salix* spp.), along with mulefat (*Baccharis salicifolia*) and emergent freshwater marsh vegetation, such as California bulrush (*Schoenoplectus californicus*) and broad-leaf cattail (*Typha latifolia*). Upland slopes of the basins have been restored with native coastal sage scrub species, such as California buckwheat (*Eriogonum fasciculatum*), coyote brush (*Baccharis pilularis*), and California sagebrush (*Artemisia californica*).

A small area of the lower channel through north Long Beach supports a rare seasonal alkali wetland community, with patches of southern tarplant (*Centromadia parryi* ssp. *australis*), along with wetsalts tiger beetles (*Cicindelidia hemorrhagica*) and other wetland indicators.

Wildlife Movement

This channel and that of the San Gabriel River, several miles to the east, are the only geographic features connecting the southern coast of Los Angeles County to SEAs and other natural areas located 15 or more miles inland.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each

DRAFT

critterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE LOWER LOS ANGELES RIVER SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	<u>Met</u>	- <u>Southern tarplant</u>
B/ C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	<u>Met</u>	- <u>Restored freshwater marsh</u> - <u>Riparian woodland (sycamore-cottonwood-willow)</u> - <u>Seasonal alkali wetland</u> - <u>Least bittern</u> - <u>Clark’s marsh wren</u> - <u>Northern yellow warbler</u> - <u>Wetland dependent nesting birds</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	<u>Met</u>	- <u>Large migratory shorebird concentrations</u> - <u>Fall migration stopover habitat</u> - <u>Concentrated feeding and resting habitat for shorebirds</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	<u>Not Met</u>	
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	<u>Not Met</u>	

Criterion	Status	Justification
G) <u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Los Angeles River coastal-inland linkage</u> - <u>Urban river movement corridor</u>

Madrona Marsh Preserve SEA

General Location and Landform

The Madrona Marsh Preserve SEA lies on the northeast corner of Madrona Avenue and Sepulveda Boulevard, within the City of Torrance. The SEA boundaries encompass the natural part of the Madrona Marsh Preserve, which is a remnant of one of the last natural vernal wetlands in the County. Vernal marshes fill in the rainy season (winter and spring in our area) and dry completely by the end of summer. They are heavily used by migrant birds in the spring, and in the fall if rains are early. They usually have other resident flora and fauna that are specially adapted to the seasonal cycle of wetting and gradual complete desiccation.

The SEA is located within the Torrance United States Geological Survey (USGS) 7.5' California Quadrangle.

The SEA supports one of few remaining freshwater wetlands that once covered much of the southern and western Los Angeles Basin area. In the lowland are vernal marshes and an alkaline margin, and the upland supports a coastal prairie ("sand prairie") ecosystem and vernal pools.

The freshwater plants and animals found here are completely surrounded by residential development and industrial facilities. Most habitat of this type in the County has been filled, drained, and lost to development. In some areas, man-made lakes and ponds have created small freshwater marshes along their edges, but this is minimal in comparison to the large expanses of freshwater marsh that were once found in the Los Angeles Basin.

Biological Resources

Vegetation and communities

Freshwater marsh habitat at Madrona Marsh supports a great diversity of wildlife. Most of the bird species are dependent in some way on the surface moisture and vegetation, and would not be able to survive without it. It is also a habitat that supports several species of amphibians, including western toad (now localized in the coastal lowlands of southern California).

DRAFT

Madrona Marsh contains vernal pools, which harbor one of the world’s largest populations of southern tarplant (*Centromadia parryi* ssp. *australis*). The vernal pool community here (plants, invertebrates, and amphibians) is similar to the one found at Chatsworth Reservoir, but otherwise, several species have their *only* Los Angeles Basin populations at Madrona Marsh, such as pygmy boisduvalia (*Epilobium campestre*), water plantain (*Alisma triviale*), dwarf wooly-heads (*Psilocarphus brevissimus* var. *brevissimus*), water starwort (*Callitriche marginata*), California waterwort (*Elatine californica*), versatile fairy shrimp (*Branchinecta lindahli*) and others.

Madrona Marsh is one of three remaining wetlands with freshwater marsh in the County. This type of habitat once covered the South Bay area, and supports several species of amphibians including frogs and toads that are becoming rare throughout southern California. In addition, it has some vernal pools, which are very rare in the County. The marsh is also an important area for migratory birds. Over 275 bird species have been recorded using Madrona Marsh as of March 2026 (eBird). Because remnant freshwater marshes like Madrona Marsh are the only habitat of this type in southern portion of the County, they serve as miniature wildlife refuges. Kenneth Malloy Harbor Regional Park (Harbor Lake Regional Park SEA) and Ballona Freshwater Marsh (Ballona Wetlands CRA) are the two other freshwater marshes in this area. Waterfowl, shorebirds, marsh birds, and others can be found on the marsh in numbers during the spring and fall migration.

Wildlife Movement

The vernal pools situated within this SEA serve as an isolated, high-quality resource providing habitat linkage for migratory waterfowl. The vernal pools teem with arthropod and amphibian activity, and provide essential feeding grounds for long-distance migrants, as well as for resident species of reptiles, birds and mammals. The ponds do not lie within any identified terrestrial movement routes for wildlife.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE MADRONA MARSH PRESERVE SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	<u>Met</u>	<u>– Southern tarplant (one of the world’s largest populations)</u>

DRAFT

Criterion		Status	Justification
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	<u>Met</u>	<ul style="list-style-type: none"> – <u>Remnant vernal marsh and seasonal freshwater wetland</u> – <u>Coastal prairie (sand prairie) with vernal pools</u> – <u>One of few remaining freshwater wetlands in Los Angeles Basin</u> – <u>Vernal pool flora with Basin-limited distributions</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	<u>Met</u>	<ul style="list-style-type: none"> – <u>Seasonal wetland breeding habitat for amphibians</u> – <u>Important spring and fall migratory bird stopover</u> – <u>Concentrated feeding habitat for waterfowl and shorebirds</u> – <u>Limited freshwater wetland habitat in southern Los Angeles County</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	<u>Not Met</u>	

Criterion		Status	Justification
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Remnant vernal marsh and freshwater wetland ecosystem</u> – <u>One of last intact seasonal wetland systems in South Bay</u> – <u>Preserved coastal lowland vernal pool community</u> – <u>Representative historic Los Angeles Basin freshwater marsh</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Not Met	

Pacoima Wash SEA

General location and landform

Lower Pacoima Wash extends for more than mile downstream of the USFS boundary as an un-channelized, boulder-filled coastal-slope alluvial wash, and is blocked by a large flood-control dam at Maclay St., just east of the 210 Freeway. One of the rarest habitats in the state, alluvial fan scrub is restricted to the remaining natural-floor drainages in the Los Angeles Basin and nearby lowlands, where the Basin meets the foothills of the Transverse and Peninsular ranges (e.g., San Gabriel River, Santa Ana River, San Jacinto River). The alluvial habitats of these sites stand in stark contrast with the surrounding upland habitats, which typically support chaparral and coastal sage scrub (or, more often, disturbed versions of these habitats, filled with non-native grasses and forbs) not associated with seasonal water flow.

The SEA is located within the San Fernando United States Geological Survey (USGS) 7.5' California Quadrangles.

While the habitat at Pacoima Wash is less extensive and more disturbed than that at Big Tujunga Wash, the San Gabriel River, and a handful of other drainages in the eastern part of the County, it nonetheless supports numerous rare species (including many of those also found at Big Tujunga Wash), and thus represents an integral component of the original vegetation of the Los Angeles Basin.

Located at the northeastern edge of the densely urban San Fernando Valley, Pacoima Wash is bordered by El Cariso Community Regional Park to the west, and by dense residential and industrial development to the south; however, the northern connection to Angeles National Forest is still intact, and numerous large corridors into the ANF to the east (Lopez Canyon area) are also still open

DRAFT

(including via “Lopez Canyon Sylmar Hiking Trail” and Lime Kiln Canyon (note: a different Limekiln Canyon is located on the southeastern slopes of the Santa Susana range, several miles to the west).

Biological Resources

Vegetation and communities

The alluvial fan scrub present at Pacoima Wash supports a highly diverse scrub community with California buckwheat (*Eriogonum fasciculatum*) particularly prevalent, and scattered laurel sumac (*Malosma laurina*). Notable plant species present include Tujunga bushmallow (*Malacothamnus davidsonii*), scalebroom (*Lepidospartum squamatum*, an indicator species for this habitat), and cane cholla (*Cylindropuntia bernardina*), all of which are highly restricted in the Los Angeles Basin. All (southern California) areas of alluvial fan scrub feature plants at their extreme physiological range, trying to survive in an extremely harsh environment (boulder-filled floodwaters, searing hot temperatures). Even riparian trees such as western sycamore adopt a distinctive growth form on these bajadas (alluvial fan slopes), stunted and often with multiple trunks.

Sensitive wildlife species known from the wash, or from nearby similar habitat, include San Diegan legless lizard (*Anniella stebbinsi*), Blainville’s horned lizard (*Phrynosoma blainvillii*), two-striped garter snake (*Thamnophis hammondi*) and coastal whiptail (*Aspidoscelis tigris stejnegeri*). Numerous rare species are known from upstream of USFS boundary, and may also occur in the lower wash, at least seasonally, including rare fishes, southwestern pond turtle (*Actinemys pallida*) and amphibians such as arroyo toad (*Anaxyrus californicus*). Surveys are urgently needed to document the presence (or absence) of these and other sensitive taxa.

Trask’s shoulderband is known from the area, and at least some of these records may pertain to the narrowly-endemic Pacoima shoulderband subspecies (*Helminthoglypta traskii pacoimensis*), which is known only from “Pacoima Canyon” and is believed to be “Critically Endangered”; the subspecies is known from specimens but has not been photographed in life.

Wildlife Movement

The natural habitats of Pacoima Wash essentially stop at the edge of urban development and are blocked from the rest of the San Fernando Valley by the massive 210 Freeway. However, a series of flood control channels convey water from the wash southwest to a large complex of settling ponds near Arleta, and then south into the Los Angeles River channel. Numerous pocket parks have been installed or are planned along Pacoima and Tujunga Wash, and the Los Angeles River, emphasizing its importance as an urban wildlife corridor. This includes not just large and mid-sized mammals, but also pollinators taking advantage of native plants (natural or planted), and smaller reptiles supported by strips of dirt (rather than concrete) along the channel right-of-way. These urban habitats are essentially “anchored” by the extensive alluvial scrub habitat of Pacoima Wash, which in turn is ecologically and literally connected to the extensive habitats of the San Gabriel Mountains SEA.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE PACOIMA WASH SEA

DRAFT

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	- <u>Tujunga bushmallow</u>
B/ C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	- <u>Alluvial fan scrub</u> - <u>Scalebroom</u> - <u>Cane cholla</u> - <u>San Diegan legless lizard</u> - <u>Blainville's horned lizard</u> - <u>Two-striped garter snake</u> - <u>Coastal whiptail</u> - <u>Trask's shoulderband snail</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Not Met	
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	- <u>Flood-scoured, boulder-dominated alluvial fan scrub</u> - <u>Scalebroom and Tujunga bushmallow adapted to disturbance regimes</u> - <u>Multi-trunk western sycamore growth forms on bajadas</u>
F)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	- <u>Relatively intact alluvial fan scrub</u> - <u>Remnant natural wash habitat in urban basin</u> - <u>Boulder-dominated natural drainage</u>

Criterion	Status	Justification
G) <u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Linkage to San Gabriel Mountains SEA and Angeles National Forest</u> - <u>Anchor habitat for urban wash corridors</u> - <u>Connectivity via Pacoima-Tujunga-Los Angeles River system</u>

Palos Verdes Peninsula and Coastline SEA

General location and landform

The Palos Verdes Peninsula and Coastline SEA is chiefly located offshore around the Palos Verdes Peninsula, but also includes terrestrial habitat in Portuguese Bend, the Navy Fuel Depot in San Pedro and several other disjunct drainages and ridgelines. All of the many offshore rocks within 12 nautical miles of the coast are part of the California Coastal National Monument that managed by the Bureau of Land Management in the U.S. Department of the Interior. Many endangered marine mammals use this habitat. The SEA includes the inshore part of the Point Vicente State Marine Conservation Area (SMCA) and the Abalone Cove SMCA.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Torrance, San Pedro, and Redondo Beach.

In geological history, the Peninsula was once an island before sediments filled the Los Angeles Basin, and it has many biological associations today that originated during the time it was insular. Several uncommon plants, for example, are otherwise only known from offshore islands in the County, and some of the bird subspecies on the Peninsula are also only known elsewhere on the offshore islands.

The terrestrial part of the SEA includes several non-contiguous ravines, canyons and streambeds that are adjacent to rural residential areas. The area known as Portuguese Bend is one of the largest inland areas within the SEA. The SEA also includes the inshore part of the Point Vicente State Marine Conservation Area (PVSMCA), which extends 3 nautical miles in a southwest arc originating along the shoreline from 0.4 miles north of Point Vicente to Long Point. The PVSMCA is abutted to the east by the Abalone Cove SMCA (ACSMCA), which extends 3 nautical miles to the south from Long Point to Portuguese Point. No take of marine organisms is allowed in the PVSMCA; limited take is allowed in the ACSMCA.

The SEA is formed by the Palos Verdes Range, and is characterized by marine terraces atop rough terrain that is created by the erosion of the upland areas into many canyons. The flatter areas on the marine terraces are the areas that have been developed. Unparalleled headlands, rocky shoreline, and the land-sea interface provide for a tremendous variety of biotic resources in this area. It is one of the most biologically diverse and productive regions in the County, and contains several biotic communities, including rocky intertidal, kelp bed, coastal strand, and coastal sage scrub. One small

DRAFT

sandy beach is periodically present on an ephemeral basis at Portuguese Bend. This 10-mile stretch of coastline, between Point Fermin and Bluff Cove, is the only sizeable rocky intertidal area in the County.

Through the late 1900s, the bluff tops and cliffs had been disturbed by pedestrian use, residential development, and agriculture, with only very small, isolated ravines remaining in a natural state. Through the early 2000s, however, ongoing preservation and restoration work throughout the peninsula has resulted in a network of preserves managed by the Palos Verdes Peninsula Land Conservancy. Likewise, the shoreline of the peninsula, which had suffered major biological impairment through the late 1900s, including over-collection by humans and intense pollution, is now in recovery as preserved areas and enforcement of regulations has started to change the local culture from one of collection and extraction, to one centered around passive use. The health of the marine environment, relatively poor due to human influences, such as outfalls with toxic contents, appears to be slowly recovering.

Biological Resources

Vegetation and communities

Most of the interior Peninsula canyons and slopes are critical habitat for the federally-threatened coastal California gnatcatcher (*Polioptila californica californica*). Many of the interior areas are also critical habitat for the federally-endangered Palos Verde blue butterfly (*Glaucopsyche lygdamus palosverdesensis*). The gnatcatcher favors the coastal sage scrub or sumac scrub (lemonadeberry *Rhus integrifolia*) vegetation; the butterfly favors flatter areas with grassland plants of the upper marine terraces, such as Santa Barbara milk-vetch (*Astragalus trichopodus* var. *lonchus*), which is a principal food plant for its larvae. The gnatcatcher Critical Habitat covers virtually all of the terrestrial part of the SEA and extends well beyond the SEA on the Peninsula. The butterfly's critical habitat is chiefly in flatter areas of the marine terraces, but can include broader areas of the canyons. In addition to these species, the SEA supports a population of the coastal cactus wren, and several rare plants, most associated with coastal bluffs. These include San Diego bursage (*Ambrosia pumilla*), aphanisma (*Aphanisma blitoides*), Coulter's saltbush (*Atriplex coulteri*), South Coast saltscale (*Atriplex pacifica*), and Catalina crossosoma (*Crossosoma californicum*)—all rare globally—and locally-rare taxa such as Hubby's phacelia (*Phacelia hubbyi*), Lewis's evening-primrose (*Camissoniopsis lewisii*), California box-thorn (*Lycium californicum*), and woolly seablite (*Suaeda taxifolia*). State-sensitive (terrestrial) wildlife include southern California legless lizard (*Anniella pulchra*) and southern California rufous-crowned sparrow (*Aimophila ruficeps canescens*).

Rocky intertidal zones support a great number of species, owing to the oxygen and food-rich environment offered by this habitat. The natural features include a variety of substrates, both stable and unstable, the aeration of water and humidification of the air through wave splash, and the upwelling of nutrient-rich waters along the southern California coast.

Subtidally, one can find representatives of every phylum of animals (major groupings, such as mollusks, echinoderms, nudibranchs, etc.), many of which are not represented on land. Rocky substrates are as diverse as the intertidal areas. Sandy subtidal areas also support a great variety of species, but most are cryptic and submerge into the depths of the sand because of the high energy and unstable nature of their environment.

Kelp beds dominated by giant kelp (*Macrocystis pyrifera*) are an extremely productive habitat that provides food, attachment sites, and shelter for a diverse community of invertebrates, fish, and sea

DRAFT

mammals. They are an important nursery habitat for juvenile fish. Kelp beds were originally common off the southern California coast wherever rocks were present at shallow depths. However, due to man-made and natural phenomena, this habitat was severely diminished in the region, but a kelp bed habitat restoration program begun in the area in the 1960s appears to have been successful. Kelp beds have been widely reestablished offshore of the Peninsula.

The coastal cliffs found in the area range in elevation from 100 to 300 feet and support coastal sage scrub and coastal strand bluff scrub. These and offshore rocks offer ideal roosting and feeding sites for numerous shorebirds, gulls, and other seabirds, including the California brown pelican (*Pelecanus occidentalis californicus*), Brandt's and pelagic cormorants, and various shorebirds. The American peregrine falcon (*Falco peregrinus anatum*) (delisted at the state and federal levels) breeds on sea-facing cliffs here, and some of the County's last remaining breeding American Kestrels (*Falco sparverius*) may still attempt to nest around the remaining vacant lots and grassy areas on the peninsula.

Wildlife Movement

Though isolated from terrestrial habitats elsewhere in the Los Angeles Basin, the Palos Verdes peninsula is a key stepping stone for migratory birds moving through the region. Local wildlife is also supported by the remaining natural canyons preserved on the peninsula. So, while it does not fall within any identified terrestrial movement routes for wildlife because it is isolated by the ocean on one side, the fact that it is separated by miles of developed land on the remaining terrestrial sides may actually increase its importance for connectivity.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE PALOS VERDES PENINSULA AND COASTLINE SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	<u>Met</u>	<ul style="list-style-type: none">- <u>Coastal California gnatcatcher</u>- <u>Palos Verdes blue butterfly</u>- <u>Coastal cactus wren</u>- <u>Monarch butterfly overwintering sites</u>- <u>Black abalone and white abalone (offshore habitat)</u>- <u>Multiple globally rare coastal bluff plant species</u>

DRAFT

Criterion		Status	Justification
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>Channel Islands–affiliated flora and fauna with only mainland LA County occurrences</u> – <u>Coastal bluff scrub and marine terrace grassland communities</u> – <u>Rocky intertidal and kelp bed habitats rare in Los Angeles County</u> – <u>Multiple state-rare coastal bluff plant assemblages</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Coastal cliffs providing roosting and nesting sites for seabirds</u> – <u>Rocky intertidal and kelp beds supporting marine breeding and feeding areas</u> – <u>Offshore waters supporting marine mammal movement and migration</u> – <u>Shorebird and seabird feeding habitat along coastal interface</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Former island landform with insular biogeographic affinities</u> – <u>Biota more similar to Channel Islands than mainland Los Angeles Basin</u> – <u>Isolated peninsula supporting distinct coastal populations</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Only extensive rocky intertidal shoreline in Los Angeles County</u> – <u>Intact coastal sage scrub and bluff scrub preserves</u> – <u>Restored kelp bed and nearshore marine communities</u> – <u>Preserved marine terrace and canyon habitats</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Stepping-stone for migratory birds along southern LA County coast</u> – <u>Coastal canyon habitats supporting localized wildlife movement</u>

Puente Hills SEA

General location and landform

The Puente Hills SEA is located in the Puente Hills in the southeastern portion of the County. The Puente Hills are an inland topographical feature that separates the San Gabriel Valley to the north and the coastal plain to the south. The hills are oriented east-west and stretch from the San Gabriel River on the west approximately to the San Bernardino-Los Angeles County line to the east, where they transition into the Chino Hills. The SEA includes portions of the Whittier Narrows Dam Recreation Area, including flood control basins and nearby spreading grounds for the San Gabriel River and Rio Hondo, as well as much of the undeveloped land throughout the Puente Hills. Nearly the entire SEA is designated as the Puente-Chino Hills State Important Bird Area (IBA) by Audubon California. The area hosts migrating and resident birds that use the extensive mosaic of lowland terrestrial habitats, and notable extensive areas of grassland and oak and walnut woodlands.

The SEA encompasses portions of the El Monte, Baldwin Park, Whittier, La Habra and Yorba Linda United States Geological Survey (USGS) 7.5' California Quadrangles. The majority of this SEA lies within unincorporated areas of the County.

In the westernmost part of the SEA, previously adopted boundaries encompass portions of the Montebello Hills that have subsequently been developed with homes since 2010, leaving the remaining open space narrow and fragmented. Nevertheless, these areas continue to be inhabited by the federally-threatened coastal California gnatcatcher (*Polioptila californica californica*) (formerly thought to be the largest population in the County). This area is part of Critical Habitat for the gnatcatcher, which extends to Montebello Boulevard on the north side and to the west side of San Gabriel Boulevard, while the SEA boundary stops at the transmission lines' southern edge. Some of the canyons in the area support oak woodland.

Biological Resources

Vegetation and communities

The Puente Hills SEA includes some of the most significant coastal-lowland riparian and scrub habitats left in the County. It includes a projection that extends from the Montebello Hills over San Gabriel Boulevard to the oak woodland bordering the Rio Hondo Channel among oil field structures. The projection continues upstream along the natural riparian course of the Rio Hondo to a point where the Rio Hondo is encased as a concrete flood control channel. This area of the Rio Hondo usually has surface water and is on the migration route for migrating and wintering waterfowl. The riparian area of the Rio Hondo immediately north of San Gabriel Boulevard has long been recognized as important to birds and has been called the Bosque del Rio Hondo since Spanish occupation of this area (since developed into public parkland). While the abundance of birdlife is frequently highlighted, the area also provides habitat for amphibians, reptiles, and mammals.

This SEA represents the only large complex of multiple, relatively undisturbed habitats in southeastern portion of the County, and some of the best-preserved examples of lowland (<1,000' elevation) southern California habitats. It is regionally important to many resident species, as well as migrating species, which would otherwise not be able to meet their habitat requirements. In particular, large mammal and overwintering birds of prey and migratory songbirds make use of this area. The Puente Hills are a well-known migration corridor for migratory songbirds during spring migration (April and May). On foggy May mornings, the hilltops can support hundreds of individual migrant songbirds, which forage actively in all available habitats.

DRAFT

The Puente Hills represent the clear northern edge of the Peninsular Ranges. For this reason, taxa such as red diamond rattlesnake, occur here at the edge of their range, where they co-occur with the more widespread Pacific rattlesnake. Several bird and wildlife species extend west through the Puente Hills into the Los Angeles Basin, where they are otherwise rare (e.g., western gray squirrel, long-tailed weasel, arboreal salamander, and grasshopper sparrow).

The SEA encompass natural areas of the basin and portions of the San Gabriel River and Rio Hondo and associated upland habitat, along with Legg Lake, which features relatively extensive freshwater marsh, riparian habitat, and regionally-significant rookeries of nesting large waterbirds. The riparian habitat within Whittier Narrows supports some of the largest breeding populations of riparian-obligate birds in the Los Angeles Basin, including least Bell's vireo and yellow-breasted chat.

A plan for the Whittier Narrows area, sometimes called the Emerald Necklace, proposes to eventually develop a string of parks along the San Gabriel River and Rio Hondo, stretching from the Whittier Narrows Dam County Recreation Area along the rivers to the Santa Fe Dam area, into the City of Irwindale. This would better connect wildlife from the Puente Hills, Chino Hills, Santa Ana Mountains and the other Peninsular Ranges of southern California with the San Gabriel Mountains, and the Transverse Ranges across the northern side of the San Gabriel Valley.

Much of the summit area in the Puente Hills is conserved and under restoration by the Puente Hills Habitat Preservation Authority (PHHPA), which has sponsored important studies on habitat and wildlife movement, restoration, and the wildlife movement tunnel under Harbor Boulevard at the eastern side of the SEA.

The SEA at the western end of the Puente Hills includes walnut woodland and riparian scrub within Sycamore Canyon, and mixed riparian forest of ash, sycamore, and oak within Turnbull Canyon. The ridges around Turnbull Canyon are mixed grassland and chaparral, with coastal sage scrub and a scattered population of the federally-threatened coastal California gnatcatcher. Mixed chaparral and cactus scrub on the Turnbull Canyon and Sycamore Canyon slopes support a local population of the sensitive coastal cactus wren (*Campylorhynchus brunneicapillus sandiegensis*). Other rare species include a diverse raptor community which includes the white-tailed kite (*Elanus leucurus*), nearly extirpated as a nester in Los Angeles County, and a handful of rare plants including intermediate mariposa lily (*Calochortus intermedius*).

The southern SEA boundary in this area skirts development in the community of City of La Habra Heights and includes the southern slope of Powder Canyon, which has a variety of natural habitats including riparian, walnut, and north-slope oak woodlands, California walnut woodland, rocky hillsides, chaparral, and some grassland. The walnut woodlands in this area are reported to be the best remaining stands south of Ventura County. Pathfinder Park, also within the SEA, has riparian and chaparral habitat. An area around the ridgeline development along Vantage Pointe Drive has been excluded from the SEA. The northern SEA boundary continues to the edge of development at the southern edge of the communities of Rowland Heights to include walnut woodland, oak woodland, grassland, and chaparral vegetation on the north-facing slopes and central ridge of the Puente Hills. This area includes tributaries in the Puente Hills for the Brea Canyon watershed.

Tonner Canyon is in the Santa Ana River watershed, and the ridge separating Brea and Tonner canyons is generally considered the divide between the Puente Hills and the Chino Hills, all of which have significant areas of remaining habitat. Though no public access is allowed, it may hold the most significant biodiversity of the entire SEA, with records nearby of southern California legless lizard, red-diamond rattlesnake, south coast gartersnake (Chino Hills State Park), and potentially breeding

DRAFT

northern harrier, turkey vulture, western meadowlark, and grasshopper sparrow.

The majority of the SEA lies within unincorporated County jurisdiction. Other local jurisdictions have also been included within the SEA in order to delineate the boundaries of functioning habitat units. These include the City of Diamond Bar, the City of Industry, the City of La Habra Heights, the City of Montebello, the City of Pico Rivera, the City of South El Monte, and the City of Whittier.

Wildlife Movement

Evidence of significant wildlife movement throughout the Puente Hills SEA has been documented in an ongoing carnivore study commissioned by the Santa Monica Mountains Conservancy as part of a multi-jurisdictional effort to establish a region wide wildlife movement linkage. Additional studies on wildlife movement through the area include several focusing on the Harbor Boulevard and Colima Road Underpasses. This SEA represents the Los Angeles County portion of a continuous series of natural open spaces within the Puente Hills and Chino Hills. Overall, this open space extends north and west from State Route 91 (SR-91) in Orange and Riverside Counties to the Whittier Narrows reach of the San Gabriel River. The open space physically links the Puente/Chino Hills with the Santa Ana Mountains and the San Gabriel Mountains, respectively. By virtue of these linkages and a complex of interconnected habitat units throughout the hills, the Puente/Chino Hills function as both an important wildlife linkage and resident habitat area for regional wildlife populations.

Maintenance of biological diversity and population viability is accorded throughout the SEA and the chance of local species extinctions due to isolation is minimized. This function is acutely important for wide-ranging species that meet their breeding and/or habitat requirements over broad areas.

Although several major arterial roads and highways cross the hills, continued use of undercrossings and surface crossings by wildlife has been documented. This movement is largely east-west trending between large habitat blocks located in the western, central and eastern portions of the SEA. Species documented as moving through the area include bobcat, gray fox (*Urocyon cinereoargenteus*), mule deer (*Odocoileus hemionus*), and the occasional mountain lion (*Puma concolor*).

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE PUENTE HILLS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none">- <u>Least Bell's vireo</u>- <u>Coastal California gnatcatcher</u>- <u>Coastal cactus wren</u>- <u>White-tailed kite</u>- <u>Intermediate mariposa lily</u>

DRAFT

Criterion		Status	Justification
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>California walnut woodland (regionally significant stands)</u> – <u>Oak riparian woodland along Rio Hondo and San Gabriel River</u> – <u>Mosaic of low-elevation coastal scrub, grassland, and woodland habitats</u> – <u>Habitat for multiple California Bird Species of Special Concern (e.g., yellow-breasted chat)</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Waterbird rookeries at Legg Lake and Whittier Narrows spreading grounds</u> – <u>Migratory songbird stopover along Puente Hills ridgelines</u> – <u>Riparian breeding habitat for riparian-obligate birds</u> – <u>Concentrated feeding habitat for raptors and overwintering birds</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Northern extent of Peninsular Range flora and fauna in Los Angeles County</u> – <u>Co-occurrence of Peninsular Range and Los Angeles Basin species</u> – <u>Range-edge occurrences of species such as red diamond rattlesnake</u>

DRAFT

Criterion		Status	Justification
E)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Intact California walnut woodland stands</u> – <u>Relatively undisturbed oak woodland and riparian habitats</u> – <u>Preserved low-elevation southern California habitat mosaic</u> – <u>Restored summit habitats managed for conservation</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Continuous linkage between Puente Hills and Chino Hills open space</u> – <u>Regional connection between Santa Ana Mountains and San Gabriel Mountains</u> – <u>East-west wildlife movement across interconnected habitat blocks</u> – <u>Documented movement of bobcat, mule deer, gray fox, and mountain lion</u> – <u>Functional linkage across Whittier Narrows and river corridors</u>

San Andreas SEA

General location and landform

The San Andreas SEA is located in the western portion of the Antelope Valley, largely on unincorporated lands. This large SEA roughly follows the San Andreas Fault, and includes many diverse habitats, as it represents an ecotone, or meeting place for the Central Valley, the Tehachapi Mountains (which include many elements of the Sierra Nevada); San Gabriel Mountains, and the Mojave Desert. These areas provide a great variety of habitats, as well as areas of water important for wildlife, plant dispersal and connectivity. The SEA includes a small portion of the south-facing Tehachapi foothills near Gorman, which are known for wildflower field displays in years of good rainfall. The SEA extends east and south across grasslands at the western tip of the Antelope Valley, and includes Quail Lake, a natural sag pond that was dammed to receive water from the West Branch of the California Aqueduct. From Quail Lake, the SEA extends southeast up the northern foothills of

DRAFT

Liebre Mountain, Sawmill Mountain, and Portal Ridge; large portions of Leona Valley; the northern base of Ritter Ridge, Fairmont and Antelope buttes; and portions of Anaverde Valley. It also includes Lake Palmdale and adjacent Una Lake, terminating near Barrel Springs. While the San Andreas Fault continues to the east, those areas are included as part of the Antelope Valley SEA. The Antelope Valley and adjacent desert slopes of the SEA are recognized by Audubon California as the Lancaster Globally Important Bird Area (IBA).

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Frazier Mountain, Lebec, La Liebre Ranch, Neenach School, Fairmont Butte, Little Buttes, Black Mountain, Liebre Mountain, Burnt Peak, Lake Hughes, Del Sur, Lancaster West, Sleepy Valley, Ritter Ridge, and Palmdale.

The majority of land within the SEA lies within unincorporated area of the County. Other jurisdictions include the Angeles National Forest, the City of Palmdale, and the City of Lancaster.

Biological Resources

Vegetation and communities

The transition of several habitat types including creosote bush scrub, Joshua tree/California juniper mixed woodland, and desert chaparral, makes the SEA valuable for educational and scientific reasons. The close proximity of the Fairmont and Antelope buttes to the San Gabriel Mountains renders them unique in their species composition and ecological relationships and, therefore is of scientific interest. This area also harbors the southern limit of the Sierra Nevada foothill woodland community, which includes blue oak, gray or foothill pine, and California buckeye, as well as rare relic stands of great basin sagebrush scrub.

The northern slopes of Ritter Ridge (southwest of Palmdale) support one of the most pristine mixed stands of Joshua tree and California juniper in Los Angeles County.

The location of the SEA at the confluence of five major geographical areas, the Mojave Desert, the Central Valley, the San Gabriel Mountains of the Transverse Ranges, the Coastal Ranges, and the Tehachapi Mountains has produced a community-rich area with desert, foothill, and montane environments. The SEA encompasses large, mostly undisturbed examples of all of these communities. Some of the westernmost clonal stands of western Joshua trees (*Yucca brevifolia*) are present within this SEA, with a small number of stands and individuals known west of Interstate-5. The SEA includes the northern slope of the Angeles National Forest with its diversity of chaparral, grasslands, and oak and conifer forests.

The entire area along the San Andreas Fault is rich in wetlands and bogs, but Leona Valley has these in abundance, even in many yards. The wetlands in the San Andreas Fault valley and Portal Ridge have supported nesting tricolored blackbird, now very rare as a breeder in southern California, with more individuals arriving in winter to utilize the pasturelands, stables and other agricultural features of the SEA.

The far western area of the SEA, along the Kern-Los Angeles County line, is coincident with designated Critical Habitat for the federally-endangered California condor (*Gymnogyps californicus*), a bird that nearly went extinct and has been kept extant in the wild by captive breeding, beginning in the 1980s. Many sensitive plant species are known from the SEA and have virtually no other habitat in the County, including Clokey's cryptantha, spreading navarretia, Horn's milk-vetch (possibly restricted to Lake Palmdale), and Mason's neststraw. Important populations of rare wildlife are also

DRAFT

found here, for pond turtle (apparent hybrids), golden and bald eagles, and California spotted owl (in montane areas).

Several protected areas are contained within this SEA, including the Antelope Valley California Poppy Reserve State Natural Reserve.

These desert buttes are concentrated wintering grounds for birds of prey, including ferruginous hawks, northern harriers, and a small population of resident burrowing owls, and include small alfalfa farms and pastures that support a plentiful food supply of rodents and rabbits. The most westerly buttes in the Mojave Desert are found here, and with their proximity to the San Gabriel and Liebre Mountains, support a distinctive transitional habitat that fits neatly into neither the desert nor the montane biotope. The open space associated with Fairmont and Antelope buttes provide vital habitat to many wide-ranging species, which forage in outlying habitat, but use the buttes for nesting, roosting, denning, and refuge.

A reintroduced population of pronghorn (*Antilocapra americana*) has been detected within this SEA. Ranch yards provide critical breeding habitat for the County's remaining nesting Swainson's hawks, which also forage within the SEA. Lakes and other wetland areas along the Fault and throughout the SEA provide breeding habitat for amphibians and feeding habitat for migrating birds that traverse the slopes adjacent to the Mojave Desert.

Wildlife Movement

The SEA includes several important linkages for wildlife movement. The foothills in the western-most part of the SEA are an important linkage between the San Gabriel Mountains, the Tehachapi Mountains, and the Coastal Ranges. This linkage to the Tehachapi Mountains is important because they connect to the southernmost extent of the Sierra Nevada Mountains. The Tehachapi Mountains represent the only mountain linkage from the Transverse Ranges and the Coast Ranges to the Sierra Nevada Range. This feature may be an important topographic reference for migrating birds, as well as providing high elevation foraging grounds along the migratory route. The several ranges that meet at the western end of the SEA, provide a valuable link for gene flow between divergent subspecies, varieties, and populations of many species. The SEA includes numerous drainages that extend onto the Antelope Valley floor towards resources, such as the Fairmont and Antelope buttes. These washes provide an important linkage for animals traveling between the Valley floor, the buttes and the western part of the San Gabriel Mountains. In addition, Anaverde Creek, Amargosa Creek, and Pine Canyon facilitate east-west wildlife movement through the mountains, Portal Ridge, and Ritter Ridge. Tributary drainages from the Santa Clara River, such as Elizabeth Lake Canyon and San Francisquito Canyon, connect coastal drainages and the coastal ecoregion to the Fault and interior watersheds. The frequency of valuable riparian communities along this travel route, which are located within an otherwise arid climate, further contributes to the SEA's importance for wildlife and habitat linkages in the region.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

DRAFT

CRITERIA ANALYSIS SUMMARY OF THE SAN ANDREAS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>California condor</u> – <u>Swainson’s hawk</u> – <u>Tricolored blackbird</u> – <u>Burrowing owl</u> – <u>California spotted owl</u> – <u>Golden eagle</u> – <u>Bald eagle</u> – <u>Western pond turtle</u> – <u>Western Joshua tree (range-limited populations)</u> – <u>Short-joint beavertail</u> – <u>Clokey’s cryptantha</u> – <u>Spreading navarretia</u> – <u>Horn’s milk-vetch</u> – <u>Mason’s neststraw</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>At least 40 special-status species (excluding Threatened and Endangered) representing a variety of habitat types.</u> – <u>Mixed Joshua tree– California juniper woodland</u> – <u>Great Basin sagebrush relic stands</u> – <u>Fault-associated wetlands and sag ponds</u> – <u>Transitional desert– montane habitat mosaic in western Antelope Valley</u>

DRAFT

Criterion		Status	Justification
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Rocky buttes supporting nesting, roosting, and refuge for raptors</u> – <u>Ranchlands providing breeding habitat for Swainson’s hawk</u> – <u>Lakes and wetlands supporting amphibian breeding and migratory birds</u> – <u>Wintering grounds for raptors in desert butte complexes</u> – <u>Foraging and movement habitat for pronghorn</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Westernmost stands of Joshua tree woodland</u> – <u>Southern limit of Sierra Nevada foothill woodland community</u> – <u>Convergence of multiple biogeographic regions</u> – <u>Co-occurrence of desert, montane, and foothill species assemblages</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Large intact desert, foothill, and montane habitat mosaic</u> – <u>Extensive undisturbed ranchlands and open space</u> – <u>Intact Joshua tree–juniper woodland stands</u> – <u>Preserved fault-associated wetlands</u>

DRAFT

Criterion		Status	Justification
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Linkage between San Gabriel Mountains, Tehachapi Mountains, and Coastal Ranges</u> – <u>Connection from Transverse Ranges to Sierra Nevada via Tehachapi Mountains</u> – <u>Wash and drainage corridors linking Antelope Valley floor to mountain habitats</u> – <u>East-west movement along Portal Ridge and Ritter Ridge</u> – <u>Riparian corridors connecting coastal and interior watersheds</u>

San Gabriel Canyon SEA

General location and landform

The San Gabriel Canyon SEA is centered on the southern foothills of the central San Gabriel Mountains, at the entrance to the San Gabriel River canyon. Three major tributaries emerge here, which flow from the mountains and interconnecting terrain: Santa Anita, Monrovia and Sawpit canyons, located above (or partially within) the cities of Sierra Madre, Arcadia, Monrovia, Duarte, Bradbury, Irwindale, and Azusa. A substantial part of the SEA along the San Gabriel River is in the Audubon California-designated Important Bird Area (IBA) of the Los Angeles Flood Control Basin IBA. The San Gabriel River has largely been dammed and channelized, but with infrequent clearing of the detention basins and wash areas, substantial parts of the San Gabriel River have reverted to riparian habitat or the even more rare alluvial fan habitat, both of which support many resident and migratory birds, as well as numerous species of reptiles, amphibians, mammals and invertebrates.

The SEA is found within the, Mount Wilson, Azusa, San Dimas, and Glendora U.S. Geological Survey (USGS) 7.5' California Quadrangles.

Biological Resources

Vegetation and communities

The wide range of elevation, topography, slope aspect, and geology represent a wide array of physical habitats within this SEA. Consequently, multiple major plant communities exist in close proximity, including riparian canyon habitat, a great diversity of shrublands (both burned and unburned), plus woodlands and forests. This has led to an extremely high species diversity, including some of the

DRAFT

highest bird species diversity in the County. Of particular note, this SEA contains the last remaining relatively well-developed lower montane riparian habitats in the eastern County and dammed drainages have created additional riparian and wetland formations along Sawpit and Santa Anita canyons. Clusters of sensitive plant occurrences and vegetation microhabitats are frequently encountered here.

The mouth of the San Gabriel Canyon supports numerous special-status species, including most of the range of the San Gabriel Mountains dudleya (*Dudleya densiflora*) and San Gabriel River dudleya (*Dudleya cymosa* ssp. *crebrifolia*), both in different parts of the granitic rubble and canyon walls in the main canyon and tributaries. This SEA also includes the primary habitat area for the San Gabriel bedstraw (*Galium grande*), which is another local endemic. The Los Angeles Flood Control Basin IBA covers all of the SEA in the San Gabriel River and downstream at the Santa Fe Dam Recreation Area.

The south banks and ridge of Clamshell Canyon is critical habitat for the federally-endangered Branton's milk-vetch (*Astragalus brauntonii*), a leguminous shrub that requires interbedded sandstone and carbonate substrate deposited near the coastline of former oceans. Very limited areas of this substrate occur at the boundary of the San Gabriel Mountains in this area (most of the rocks of the San Gabriel Mountains are igneous granites and metamorphic rocks). Santa Anita Canyon has some stands of Pacific madrone (*Arbutus menziesii*), mainly known elsewhere from the Pacific coast north of Santa Barbara to British Columbia. The Santa Anita Canyon stands comprise one of the few places madrone is found between Santa Barbara and Baja California.

The Santa Fe Flood Control Basin is one of the most unusual vegetation habitats in the County (alluvial fan scrub), with multiple special-status species, including coastal California Gnatcatcher, least Bell's vireo and mesa horkelia (*Horkelia cuneata* var. *puberula*).

Elsewhere, the SEA provides important habitat for some of the rarest species in the County, including such aquatic species as southwestern pond turtle (*Actinemys pallida*), Santa Ana speckled dace (*Rhinichthys gabrielino*), and Santa Ana sucker (*Pantosteus santaanae*).

Wildlife Movement

Wildlife movement within the SEA takes on two primary modes; first, due to the steep topography, it is logical to expect considerable movement of wildlife up and down the sizeable drainages, which course through this SEA to connect the forest interior with foothill areas. Consequently, this type of movement occurs on a seasonal and more frequent basis, particularly for large mobile mammals whose full range of habitat needs are typically met over broad areas, including American black bear, mountain lion (*Puma concolor*), mule deer (*Odocoileus hemionus*), gray fox (*Urocyon cinereoargenteus*) and other medium-sized mammals. The second major type of movement occurs across the flanks of the foothills and lower mountains, in an east-west direction. Particularly for migratory birds, a corridor linking lower elevation riparian and oak woodland habitats in the SEA is of high use and importance.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

DRAFT

CRITERIA ANALYSIS SUMMARY OF THE SAN GABRIEL CANYON SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> - <u>Coastal California gnatcatcher</u> - <u>Least Bell's vireo</u> - <u>Southwestern pond turtle</u> - <u>Santa Ana speckled dace</u> - <u>Santa Ana sucker</u> - <u>Bald eagle</u> - <u>Braunton's milk-vetch</u> - <u>San Gabriel bedstraw</u> - <u>San Gabriel Mountains dudleya</u> - <u>San Gabriel River dudleya</u> - <u>Mesa horkelia</u> - <u>Mountain lion (Southern California/Central Coast ESU)</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> - <u>More than two dozen special-status species (excluding Threatened and Endangered)</u> - <u>Alluvial fan scrub (rare in Los Angeles County)</u> - <u>Lower montane riparian habitat in eastern County</u> - <u>Oak riparian woodland and southern willow scrub</u> - <u>Coastal sage scrub and walnut woodland mosaic</u> - <u>High-diversity foothill canyon habitat complex</u>

DRAFT

Criterion		Status	Justification
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Waterbird rookeries in upper San Gabriel Canyon reservoirs</u> – <u>Nesting habitat for mainland bald eagle at San Gabriel Reservoirs</u> – <u>Waterbird rookery at Santa Fe Dam Recreation Area</u> – <u>Riparian breeding habitat for migratory birds</u> – <u>As of 2026, the County’s sole mainland nesting pair of bald eagle is also found at the San Gabriel Reservoirs.</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Endemic plant concentration associated with localized substrates</u> – <u>Co-occurrence of multiple narrowly endemic San Gabriel Mountains plants</u> – <u>Isolated population of Pacific madrone in southern California</u> – <u>Unique alluvial fan scrub vegetation in flood control basin</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Extensive relatively undisturbed foothill canyon habitats</u> – <u>Intact lower montane riparian and woodland communities</u> – <u>Preserved alluvial fan scrub habitat</u> – <u>Large continuous natural habitat at canyon mouth</u>

Criterion		Status	Justification
G)	Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.	Met	<ul style="list-style-type: none"> – North-south movement along major canyon drainages – East-west foothill movement corridor – Riparian linkage across San Gabriel River floodplain – Movement corridor for large mammals between mountains and foothills

San Gabriel Mountains SEA

General location and landform

The San Gabriel Mountains, together with the Liebre Mountain and the Sierra Pelona just to the northwest, are part of southern California’s Transverse Range, an east-west trending range of high, steep granitic mountains forming the northern edge of the Los Angeles Basin. Most of the land in these mountains is owned by the federal government and managed by the U.S. Forest Service as four separate Districts comprising Angeles National Forest. In addition, considerable areas of private lands are scattered throughout the forest, some including (rural) residential communities such as Green Valley and Kagel Canyon. The Forest boundary encompasses roughly 700,000 acres of Los Angeles County’s 3 million acres, stretching from the far northwestern corner near Gorman, southeast to Claremont at the San Bernardino County line (the San Gabriel Mountains extend several miles east into San Bernardino Co.). The Forest consists of two large, unconnected blocks of territory, split by S.R. 14 and the Santa Clara River. Despite covering just under 25% of the County’s total land area, ANF contains most of the County’s undeveloped open space, and nearly all its large roadless areas.

The San Gabriel Mountains contain some of the highest peaks in southern California, including Mount San Antonio (Mt. Baldy, aka *Joat/Yoát* “Snowy Mountain” to the Tongva, a local tribe), at 10,064’, Mt. Baden-Powell (9,339’), and a dozen other peaks over 8,000’.

A massive range containing and straddling multiple biomes (including the Mojave Desert to the north, and the coastal Los Angeles Basin to the south), the San Gabriel Mountains supported a rich Indigenous history, with native peoples taking full advantage of the available game and abundant fruits such as acorns (*Quercus* spp.), chia (*Salvia columbariae*), and manzanita (*Arctostaphylos* spp.). Trading routes and footpaths for seasonal movements were extensive, connecting different portions of the mountains, and villages, with the lowlands to the north and south.

Biological Resources

Vegetation and communities

It is difficult to summarize the vegetation of so diverse an area as the San Gabriel Mountains. Nearly all the habitats of mainland southern California may be found within Angeles National Forest, with the exception of desert-floor vegetation (e.g., creosote bush scrub, alkali sink scrub) and coastal scrub

DRAFT

vegetation at immediate coastal sites such as the Palos Verdes Peninsula. Microhabitats abound—small sag ponds forming marshy meadows are located along the northern base of the range (natural wetlands along the coastal slope have largely been replaced by urbanization). Several major watersheds originate in the San Gabriel Mountains, producing the major waterways of southern California (including Santa Clara River, Los Angeles River, San Gabriel River, and in the north, Big Rock and Little Rock creeks).

As of 2008—i.e., prior to long-term drought that began in c. 2010, and a series of devastating fires including the Station Fire (2009, 169,000+ acres) and the Bobcat Fire (2020, 115,000+ acres)—the vegetation present in the Angeles National Forest (including the bulk of the Liebre-Sierra Pelona and the San Gabriel Mountains portions) was estimated at more than 50% chaparral, with 23% conifer woodland and 9% “hardwoods” (largely oak woodland) (Schaaf et al. 2008). Since 2020, however, the amount of coniferous forest has plummeted, and oak woodland is similarly reduced, due to fire and drought. Today, the pockets of high-elevation coniferous forest and meadow habitat, rich in rare species, are truly imperiled, both by forces easily managed (e.g., human intrusion and trampling) and largely out of human control (long-term drought and climate change).

In a 1998 analysis (updated with newer taxonomy in 2005), Angeles National Forest was thought to support 23 “Forest Sensitive Plant Species”, most entirely restricted in the County to the San Gabriel Mountains, and dozens of sensitive wildlife species. The San Gabriel Mountains are known to support 24 endemic or near-endemic plant taxa, 15 of which are *only* found here⁵. Many of these are listed as Threatened or Endangered by state or federal agencies or California Native Plant Society, including Great’s aster (*Symphytotrichum greatae*), San Gabriel manzanita (*Arctostaphylos glandulosa* ssp. *gabrielensis*), Palmer’s mariposa lily (*Calochortus palmeri* var. *palmeri*), San Gabriel linanthus (*Linanthus concinnus*), wooly mountain-parsley (*Oreonana vestita*), Mount Gleason paintbrush (*Castilleja gleasoni*), Bear Valley woolypod (*Astragalus leucolobus*), San Antonio milkvetch (*Astragalus lentiginosus* var. *antonius*), Ewan’s woodbeauty (*Drymocallis cuneifolia* var. *ewanii*), southern alpine buckwheat (*Eriogonum kennedyi* var. *alpigenum*), Johnston’s buckwheat (*E. microtheca* var. *johnstonii*), San Gabriel Mountains monardella (*Monardella australis* ssp. *gabrielensis*), Brown’s Flat monardella (*M. breweri* ssp. *glandulifera*), Abram’s oxytheca (*Acanthoscyphus parishii* var. *abramsii*), Rock Creek broomrape (*Aphyllon validum* ssp. *validum*), lemon lily (*Lilium parryi*), gray-leaved violet (*Viola pinetorum* ssp. *grisea*), and Peirson’s lupine (*Lupinus peirsonii*). No fewer than 30 rare, but not Threatened/Endangered, plant species are known from the San Gabriel Mountains, many occurring nowhere else in the County, including Jepson’s bedstraw (*Galium jepsonii*), urn-flowered alumroot (*Heuchera caespitosa*), California adder’s tongue (*OphioGLOSSUM californicum*) and silky lupine (*Lupinus elatus*).

Examples of sensitive wildlife taxa of the San Gabriel Mountains not found anywhere else in the County include the amphibians mountain yellow-legged frog (*Rana muscosa*) and San Gabriel Mountains slender salamander (*Batrachoseps gabrieli*); birds such as nesting bald eagle (*Haliaeetus leucocephalus*), black swift (*Cypseloides niger*), purple martin (*Progne subis*); the rare fish Santa Ana speckled dace (*Rhinichthys gabrielino*) and Santa Ana sucker (*Catostomus santaanae*) are found in the Big Tujunga watershed and locally elsewhere; San Gabriel elfin (*Callophrys mossii hidakupa*; a

⁵ <https://tchester.org/sgm/plants/species/endemics.html>

DRAFT

butterfly) and Nelson's (or "desert") bighorn sheep (*Ovis canadensis nelsoni*) are restricted to rocky habitat in the eastern portion of the range. Several other sensitive species occur primarily in the San Gabriel Mountains in the County, with small, often relict populations in the lower Liebre-Sierra Pelona range (e.g., California spotted owl, *Strix occidentalis occidentalis*).

Dozens of wildlife species that aren't recognized as sensitive at the state level are restricted in the County to the San Gabriels, with many occurring in the County *only* in these mountains. These include birds such as common nighthawk (*Chordeiles minor*), flammulated owl (*Psilosops flammeolus*), gray flycatcher (*Empidonax wrightii*), hermit thrush (*Catharus guttatus*), Lincoln's sparrow (*Melospiza lincolni*), MacGillivray's warbler (*Geothlypis tolmei*), Nashville warbler (*Leiothlypis ruficapilla*), plumbeous vireo (*Vireo plumbeus*), Williamson's sapsucker (*Sphyrapicus thyroideus*) and Wilson's warbler (*Cardellina pusilla*).

Tragically, a localized subspecies of greenish blue, the San Gabriel Mountains blue butterfly (*Plebejus saepiolus aureolus*) was rendered extinct (last seen 1980s) after a water diversion project near Big Pine destroyed its wet meadow habitat. The nearest populations of greenish blue persist in the Piute, San Bernardino and San Jacinto mountains (iNaturalist), all of which are separate subspecies than that which formerly occupied the San Gabriel Mountains.

Wildlife Movement

As a massive, largely undeveloped, largely roadless mountain range bordering a megacity to the south, and surrounded by rapidly urbanizing land, the importance of the San Gabriel Mountains to wildlife connectivity throughout the County (and beyond) cannot be overstated. The San Gabriels may be considered a "source" habitat for wide-ranging species such as mountain lion (*Puma concolor*) and golden eagle (*Aquila chrysaetos*) in the County, and presumably "supply" many of the individuals dispersing to more urban open space areas such as the Verdugo Mountains and Santa Susana Mountains. Poorly known and rarely seen mammals such as southern California ringtail (*Bassariscus astutus octavus*) and western spotted skunk (*Spilogale gracilis*) likewise would be expected to maintain source populations not in isolated urban open spaces in the Los Angeles Basin, but in the rugged mountains to the north. The freeways that border and cross the San Gabriel Mountains are thus of major concern for wildlife movement, and planning is underway to provide and improve wildlife crossing opportunities for these species in the form of underpasses and wildlife bridges.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE SAN GABRIEL MOUNTAINS SEA

DRAFT

	Criterion	Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	<u>Met</u>	<ul style="list-style-type: none"> - <u>Mountain yellow-legged frog</u> - <u>California spotted owl</u> - <u>Bald eagle nesting sites</u> - <u>Golden eagle nesting sites</u> - <u>Santa Ana sucker</u> - <u>Santa Ana speckled dace</u> - <u>San Gabriel manzanita</u> - <u>San Gabriel linanthus</u> - <u>Mountain Gleason paintbrush</u>
B/ C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	<u>Met</u>	<ul style="list-style-type: none"> - <u>High-elevation conifer forest</u> - <u>Montane meadow and sag pond wetlands</u> - <u>Endemic San Gabriel Mountains flora (multiple taxa)</u> - <u>Localized montane bird assemblages</u> - <u>San Gabriel Mountains slender salamander</u> - <u>San Gabriel elfin butterfly</u> - <u>Nelson's bighorn sheep</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	<u>Met</u>	<ul style="list-style-type: none"> - <u>High-elevation meadow and sag pond breeding habitats</u> - <u>Headwater stream habitats for endemic fish and amphibians</u> - <u>Nesting raptors in montane habitats</u> - <u>Microhabitat-restricted endemic species</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	<u>Met</u>	<ul style="list-style-type: none"> - <u>High-elevation "sky island" habitats</u> - <u>Floristic similarity to northern montane regions</u> - <u>Large mountain stream systems lacking lowland analogues</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	<u>Met</u>	<ul style="list-style-type: none"> - <u>Extensive roadless montane habitat</u> - <u>Relatively undisturbed chaparral and forest communities</u> - <u>Intact elevational vegetation gradients</u>

Criterion	Status	Justification
G) <u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Regional source habitat for wide-ranging species</u> - <u>Elevational movement corridors</u> - <u>Linkage to surrounding foothill SEAs</u> - <u>Dispersal source for basin and transverse ranges</u>

San Rafael Hills SEA

General location and landform

The San Rafael Hills are a low-elevation transverse range located in central Los Angeles County, southeast of the Verdugo Mountains and west of the Arroyo Seco. The neighborhoods of Eagle Rock and Glassell Park lie to the south, and the urbanized San Fernando Valley to the west. Elevations range from approximately 720 to 1,889 feet above mean sea level. Two major canyons, Scholl and Sycamore, run east to west. Soils include multiple calcareous types. The San Rafael Hills retain substantial areas of native vegetation, steep rugged topography, incised canyons, and natural drainage systems. Although they are sometimes referred to as an island of open space surrounded by development, tenuous connections remain to the Altadena Foothills and Arroyos SEA (across the 210 freeway through Hahamonga Watershed Park), and to the Verdugo Hills SEA (across North Verdugo Road). The hills cover approximately 7,040 acres, 3,413 of which are recommended for designation as a new SEA. Most of the remaining acreage is low-density housing and other land uses suitable for designation as connectivity overlay.

The San Rafael Hills occur within portions of the United States Geological Survey (USGS) 7.5' quadrangles for Pasadena, Glendale, and Burbank.

Although not a formal SEA Selection Criterion, it is relevant that the San Rafael Hills are part of the ancestral homelands of the Gabrieleño-Kizh/Tongva people, with the historical village of Toviscangna located approximately five miles southeast of the hills near present-day Mission San Gabriel Arcangel. The San Rafael Hills were important as a natural resource area, providing materials that included willow for houses. As such, formally recognizing the ecological significance of the San Rafael Hills through SEA designation would be consonant with and respectful of TEK.

Biological Resources

Vegetation and communities

The San Rafael Hills support a mosaic of native plant communities characteristic of foothill environments in Los Angeles County, including expanses of chaparral, coastal sage scrub, coast live oak woodland, and localized riparian and oak-sycamore woodland along canyon bottoms and ephemeral drainages. North-facing slopes and shaded ravines tend to support denser chaparral and woodland assemblages, while south- and west-facing slopes are characterized by more open coastal sage scrub and drought-tolerant chaparral species.

DRAFT

Natural communities described for this area include various forms of chaparral, coastal sage scrub, oak woodland, walnut woodland, California sycamore woodland, and mixed riparian woodland (Soza et al. 2013). Sensitive and noteworthy plant species recorded the San Rafael Hills include one plant with a California Rare Plant Rank of 1B.2 (plants rare or endangered in California and elsewhere) and several watch-list plants of limited distribution: chaparral nolina (*Nolina cismontana*), Plummer's mariposa lily (*Calochortus plummerae*), fragrant pitcher-sage (*Lepechinia fragrans*), ocellated Humboldt lily (*Lilium humboldtii* ssp. *ocellatum*), southern California black walnut (*Juglans californica*), San Gabriel oak (*Quercus durata* var. *gabrielensis*) and Engelmann oak (*Q. engelmannii*). The hills are near the western edge of the range of Engelmann oak. A distinctive form of summer lupine (*Lupinus formosus*) is also known from here⁶.

Sensitive and noteworthy wildlife species known from the San Rafael Hills include one Candidate for State listing, Crotch's bumble bee (*Bombus crotchii*), and several California Species of Special Concern: coast range newt (*Taricha torosa*), Blainville's horned lizard (*Phrynosoma blainvillii*), coastal whiptail (*Aspidoscelis tigris stejnegeri*), two-striped garter snake (*Thamnophis hammondi*), and northern yellow warbler (*Setophaga aestiva*). Also present is Trask shoulderband snail (*Helminthoglypta traskii*), which has a NatureServe Rank of S2 (Imperiled, at high risk of extirpation in the state due to restricted range, few populations or occurrences, steep declines, severe threats, or other factors).

The San Rafael Hills represent the southernmost extent of the Transverse Range flora and fauna into the Los Angeles Basin; as such, they support species like Merriam's chipmunk, black bear, and others that do not (regularly) occur farther south. Mule deer also cannot penetrate the Basin's urban matrix farther south than the San Rafael Hills. For this reason, they are considered to be an important extension of such habitat, providing a stepping-stone from the Basin into the mountains.

Wildlife Movement

The San Rafael Hills are situated south of the San Gabriel Mountains, east of the Verdugo Mountains to the west, and the Arroyo Seco to the east, and thus may function as an important linkage for these expansive open spaces. Obstacles to terrestrial movement through the San Rafael Hills include State Route 2, which runs north/south through the eastern part of the range, and the 210 freeway, which separates the San Rafael Hills from the San Gabriel Mountains. They are also the first large open space block of habitat providing a source for dispersal lower into the Basin (e.g., via the Arroyo Seco), and vice-versa. Efforts to identify and protect the remaining connections across these major barriers are warranted.

⁶ The population of summer lupine occurring from near the base of the San Gabriel Mountains through Verdugo Hills and into the Griffith Park area of the eastern Santa Monica Mountains was originally described as *Lupinus pasadenensis*. This is a perennial herb with a very small range that was subsequently included within *L. formosus*, but which differs in having a much later flowering period than other populations of *L. formosus* in the region (blooms late winter, rather than mid-summer), and occurs not in grassland, but on gneiss, a form of a decomposed granite. Mature plants closely resemble the woody shrub-forming longleaf lupine in the area (*Lupinus longifolius*), which occurs in at least three populations in the County, one in the San Gabriel Mountains foothills, another in the Santa Monica Mountains, and a third in the low hills on the floor of the Los Angeles Basin (DSC, unpubl. data).

DRAFT

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE SAN RAFAEL HILLS SEA

Criterion	Status	Justification
A) <u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Not Met	
B/ C) <u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> - <u>Chaparral, coastal sage scrub, and oak woodland mosaic</u> - <u>Walnut woodland and oak-sycamore woodland</u> - <u>San Gabriel oak occurrences</u> - <u>Engelmann oak near western range limit</u> - <u>Chaparral nolina</u> - <u>Plummer’s mariposa lily</u> - <u>Fragrant pitcher-sage</u> - <u>Ocellated Humboldt lily</u> - <u>Southern California black walnut</u> - <u>Crotch’s bumble bee (Candidate)</u> - <u>Coast range newt</u> - <u>Blainville’s horned lizard</u> - <u>Coastal whiptail</u> - <u>Two-striped garter snake</u> - <u>Northern yellow warbler</u> - <u>Trask’s shoulderband snail (S2)</u>
D) <u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Not Met	

Criterion		Status	Justification
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> - <u>Southernmost extent of Transverse Range flora and fauna</u> - <u>Range-edge populations (Engelmann oak)</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> - <u>Relatively intact foothill habitat mosaic</u> - <u>Steep, rugged terrain limiting disturbance</u> - <u>Large contiguous native vegetation patches</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Linkage to Verdugo Mountains</u> - <u>Linkage to San Gabriel Mountains via Arroyo Seco</u> - <u>Stepping-stone habitat into Los Angeles Basin</u> - <u>Constrained concentrations across SR-2 and 1-210</u>

Santa Clara River SEA

General location and landform

The Santa Clara River SEA encompasses the entire Los Angeles County reach of the Santa Clara River, as well as its main tributaries, and associated upland habitats, extending from Acton Wash in the east, downstream/west to the Ventura County line. The orientation and extent of the SEA is dictated by the surface and subsurface hydrology of the Santa Clara River, including its headwaters, tributaries, and watershed basin. This large area lies within both incorporated (Santa Clarita) and unincorporated areas of the County. The SEA encompasses a wide variety of topographic features and habitat types, all of which contribute to its exceptionally high diversity, including that of rare flora and fauna. Nearly all of the SEA is designated by Audubon California as a Globally Important Bird Area (IBA); the Santa Clara River IBA extends beyond the SEA in both upstream and downstream directions (across Soledad Pass to the Barrel Springs area in the Antelope Valley and through Ventura County to the mouth of the River at the Pacific Ocean).

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Pacifico Mountain, Acton, Agua Dulce, Sunland, San Fernando, Mint Canyon, Oat Mountain, Newhall, and Val Verde.

Biological Resources

Vegetation and communities

The eastern portion of the SEA follows natural contours at the headwaters of the Santa Clara River watershed to incorporate much of upper watershed of Soledad Canyon, as well as the Kentucky Springs and the Aliso Canyon basins, and the downstream unnamed tributaries of the Santa Clara River to Arrastre Creek. This headwater areas lie within the Angeles National Forest (San Gabriel Mountains and Liebre/Sierra Pelona), which features semi-arid chaparral and desert scrub habitat. However, most drainages themselves support vegetation of semi-desert and interior riparian habitat, which ranges from Great Basin sagebrush in Kentucky Springs Wash and Big Springs to dense, mature, willow-cottonwood-sycamore woodlands along permanent streams in drainages such as Aliso Canyon. The SEA has a large population of Parish's Great Basin sagebrush (*Artemisia tridentata* ssp. *parishii*). Tributary drainages surrounding Santa Clarita support regionally distinct alluvial scrub formations, typified by the dominance of holly-leaf cherry (*Prunus ilicifolia* ssp. *ilicifolia*).

The surrounding uplands in the SEA support pinyon-juniper woodland at highest elevations, with chaparral communities of chamise, mountain mahogany, and manzanita, and buckwheat scrub. The alluvial plain formed along the Santa Clara River wash supports intact, high diversity xeric alluvial fan sage scrub reminiscent of Big Tujunga Wash and San Antonio Wash, but with a stronger Mojave Desert influence, especially as one moves east/upstream in the drainage. Alluvial terraces have been extensively cultivated for orchard crops and dryland agriculture, particularly toward Ventura County, and in more recent years, dense residential developments have heavily encroached on the habitats here.

The SEA still supports numerous threatened and endangered plant and wildlife species largely gone from elsewhere in the County. For example, populations of the federally-threatened red-legged frog (*Rana draytonii*) are known from San Francisquito Canyon and, at least formerly, from the Aliso Canyon watershed. From upstream of the Arrastre Creek confluence to downstream in the vicinity of the railroad stop of Lang (about 13 miles of river), the floodplain of the Santa Clara River has been designated Critical Habitat for the federally-endangered arroyo toad (*Anaxyrus californicus*); some of the confluence area of Mill Canyon is also critical habitat for the arroyo toad. Additional arroyo toad Critical Habitat has been identified in the Castaic Junction area; however, whether these areas are actually occupied by the species requires additional surveys, ideally after wet rainfall years. Part of the area of Critical Habitat for the toad is also proposed Critical Habitat for the state and federally-endangered unarmored threespine stickleback (*Gasterosteus aculeatus williamsoni*), a small fish with most of its global population in the County. Once widespread throughout the Los Angeles Basin, it is now restricted to the upper Santa Clara River.

The SEA includes the watershed of Bee Canyon, a tributary of the Santa Clara River. Bee Canyon supports an important population of the federally-endangered slender-horned spineflower (*Dodecahema leptoceras*) within its broad, floodplain area. Here in Bee Canyon, slopes of coastal sage scrub mixed with chaparral elements have supported breeding by the federally-threatened coastal California gnatcatcher (*Polioptila californica californica*). While portions of this SEA have coastal sage scrub and have been designated as Critical Habitat for the threatened coastal California gnatcatcher, the only consistent population of this species is located far to the northwest of the existing SEA, in hills west of Castaic Reservoir; in part for that reason, this area has been included in the main Santa Clara River SEA.

The habitat along the Santa Clara River supports what is likely the largest community of riparian-

DRAFT

obligate birds between Santa Ynez River in Santa Barbara County and the Prado Basin in Riverside County. In the Soledad Canyon stretch are breeding summer tanager (*Piranga rubra*) and other desert-associated species, as well as yellow-breasted chat (*Icteria virens*), and abundant northern yellow warblers (*Setophaga petechia*). Other notable wildlife species frequently found in the washes of this SEA include the San Diego black-tailed jackrabbit (essentially extirpated elsewhere on the coastal slope of the County), coast horned lizard, greater roadrunner, and lesser nighthawk. The area is notable for having a combination of species that are characteristic of the desert and characteristic of coastal influence.

Just west of the confluence with Arrastre Creek, the SEA includes slopes of Parker Mountain and the eastern watershed of Hughes Canyon around the basal contours of significant rock outcroppings above the river basin, and on the south side, around the Mill Canyon tributary basin. The rocky buttes on the north side of the river, while only a minor part of the watershed of the river, provide important nesting, roosting, and sheltering habitat values for bats, birds of prey, and other sensitive species foraging along the river corridor. Emerging from these buttes, Nellus and Bobcat drainages were identified by the South Coast Wildlands Project as important to connectivity across the Santa Clara River between the western and eastern highland areas of the San Gabriel Mountains.

In upstream areas of the river and its tributaries, flows become less seasonal, and riparian resources expand and diversify. San Francisquito Creek supports dense and mature southern riparian scrub and riparian woodland formations, along with small areas of freshwater marsh. The headwaters of San Francisquito Creek are on a low ridge that bounds the San Andreas Fault Zone, and this is an important connective element of the SEA, in that it completes the path from the Pacific Ocean through the mountains to the Mojave Desert (this entire area has been recommended for inclusion as an SEA, the Liebre/Sierra Pelona SEA). West of Sand Canyon, the river has been intermittently armored to allow for development within flood hazard zones. Some irregular extensions go north into tributaries that have remnant riparian habitat and outflows from runoff that flows into neighborhood storm drains. Despite the intermittent nature of water, the slightly elevated areas among braided channels support relatively intact stands of alluvial sage scrub, riparian woodland, and southern riparian scrub. It is worth noting that the dry zones of the river are essential to the continued genetic isolation and integrity of the unarmored three-spine stickleback population in the upper reaches of the Santa Clara River (and probably are important for many other species).

At the Mint Canyon crossing, just southwest of the community of Sleepy Valley, a lobe of the SEA extends along Mint Canyon to capture riparian woodlands of coast live oak, with a number of massive heritage trees (diameters greater than 36 inches). Residences are scattered and the natural communities of chaparral are intact on the canyon slopes. Elsewhere, the alluvial fans of Oak Springs Canyon and Sand Canyon are important recharge grounds for the river aquifer and provide important passage habitat for wildlife moving north out of the San Gabriel Mountains toward the river. Here, surface flows from both canyons enter the Santa Clara River basin through natural, unconfined channels. Existing rural residential developments are excluded from the SEA, but the remaining natural highland areas of the western banks of the Sand Canyon watershed are included.

The upper areas of the canyons draining the north slope of the San Gabriel Mountains within the SEA support massive oaks and big-cone Douglas-fir, and provide habitat for the California spotted owl (*Strix occidentalis*), as well as various species of local conservation interest in the County, including Mountain Quail (*Oreortyx pictus*) and Coast Mountain Kingsnake (*Lampropeltis multifasciata*, formerly San Bernardino mountain kingsnake—*L. zonata parvirubra*). Other uncommon raptors known to breed within the SEA (typically on rugged rock outcrops or along remote canyons) include golden eagle (*Aquila chrysaetos*), long-eared owl (*Asio otus*), and prairie falcon (*Falco mexicanus*).

DRAFT

Relatively extensive areas of willow-cottonwood forest and southern riparian scrub occur west of San Francisquito Creek and within the junction zone of Castaic Creek and the Santa Clara River. These river forests support numerous sensitive species and provide multi-layered riparian habitat for a wide diversity of wildlife species, particularly birds of prey and riparian-obligate songbirds, such as the federally-endangered least Bell's vireo (*Vireo bellii pusillus*). While the southwestern willow flycatcher (*Empidonax traillii extimus*) has been detected nesting (or at least summering) in the Ventura County portion of the Santa Clara River downstream of this SEA, it is not known to have done so in recent years in Los Angeles County. More biological surprises await discovery here as new areas are made available for biological exploration; for example, the rare Newhall sunflower (*Helianthus inexpectatus*) was recently discovered growing in single spring along the Santa Clara River, adjacent to what is now Newhall Ranch.

Upland areas of these lower portions of the SEA are arguably as important as the riparian habitat here, supporting numerous rare plants, including one of just two populations of the state-endangered San Fernando Valley Spineflower (*Chorizanthe parryi* var. *fernandina*), a diminutive, once-common buckwheat relative of flat mesa tops at the perimeter of the San Fernando Valley and adjacent hills. The plant became so rare that it was believed to be extinct until it was rediscovered during required surveys for development of what is now Newhall Ranch.

Finally, the northwestern portion of the SEA has recently (since 2000) been shown to be an important movement area for the endangered California Condor, with multiple birds often seen together in the hills along I-5 west of Castaic Reservoir. These birds rarely approach the developed areas of Castaic-Santa Clarita, indicating that they have a threshold for how much human activity they can tolerate.

Wildlife Movement

The Santa Clara River channel and its alluvial terraces and tributary creeks together form arguably the single most diverse, natural wildlife movement zone through the County. Historically (and prehistorically) the riparian corridor along the Santa Clara River has served as the primary east-west linkage between the Pacific coastline, coast ranges, interior ranges, high desert and southern Sierra Nevada (via the Tehachapi Range). Mobile species can enter the river basin anywhere along its length (at least outside of developed areas) and proceed in either direction without having to pass through narrow culverts or blind channels, with continuous vegetative cover and only short stretches of dry substrates. The overall drainage course provides a continuum of aquatic and terrestrial movement opportunities, shelter, forage, and resident habitat from the mouth of the river at Ventura County and the Pacific Ocean, to the Antelope Valley. The drainage course connects to both districts of the Angeles National Forest, and links together multiple large public resource preserves (Vasquez Rocks and Placerita County Natural Areas and the Angeles National Forest).

The tributary routes in and out of this SEA extend south into the Santa Susana Mountains, south and north into the San Gabriel Mountains, northward via Castaic, Bouquet and San Francisquito tributaries (over the coastal ranges and Liebre/Sierra Pelona into the San Joaquin Valley), west into the central coast ranges, or east through the Tehachapi Mountains, and into the southern Sierra Nevada. The Agua Dulce underpass of State Route 14 is an important crossing of the highway barrier for wildlife. From that point, north riparian areas exist where the creeks (Agua Dulce and Escondido) pass through Vasquez Rocks County Natural Area.

The eastern half of the Los Piñetos undercrossing of State Route 14 onto old oil development roads enables a major wildlife conduit between the Santa Susana Mountains, the San Gabriel Mountains, and the Santa Clara River. The adjacent part of the Santa Susana Mountains and Simi Hills SEA

DRAFT

includes the west half of the Los Piñetos undercrossing of State Route 14, connecting through the natural oak woodlands and drainages adjacent to the San Fernando Pass.

The Agua Dulce Canyon area was considered for inclusion as the SEA for its value as a wildlife corridor to provide connectivity across the Santa Clara River between the western and eastern highland areas of the San Gabriel Mountains and Liebre/Sierra Pelona. Natural areas of the Sand Canyon watershed, along with the major topography of ridgelines, earthquake escarpments, grasslands, and canyon habitat features and watersheds of Bear, Placerita, Whitney, and Elsmere canyons are the important features for wildlife linkage.

The present configuration of the tributary drainages has depended upon connectivity through the Santa Clarita Valley to the north, and for now these corridors remain tenuously intact and open. However, this connectivity is far from guaranteed long-term, placing at risk the countless wildlife species—mountain lion, gray fox, bobcat, as well as birds, reptiles, amphibians, and fishes—that depend on it.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE SANTA CLARA RIVER SEA

Criterion		Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	<ul style="list-style-type: none">– <u>Arroyo toad</u>– <u>Least Bell’s vireo</u>– <u>Southwestern pond turtle</u>– <u>Unarmored threespine stickleback</u>– <u>Slender-horned spineflower</u>– <u>Spreading navarretia</u>– <u>San Fernando Valley spineflower</u>– <u>Newhall sunflower</u>– <u>Western spadefoot</u>– <u>California condor</u>– <u>Golden eagle</u>– <u>Mountain lion (Southern California/Central Coast ESU)</u>

DRAFT

Criterion		Status	Justification
B/C)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	<ul style="list-style-type: none"> – <u>Estimated 42 special-status species (excluding Threatened and Endangered)</u> – <u>Xeric alluvial fan sage scrub with Mojave Desert influence</u> – <u>Great Basin sagebrush populations within river watershed</u> – <u>Willow-cottonwood riparian forest and southern riparian scrub</u> – <u>Pinyon-juniper woodland transition zones</u> – <u>Riverine and alluvial terrace habitat mosaic</u>
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	<ul style="list-style-type: none"> – <u>Breeding raptor habitat on rock outcrops</u> – <u>Vernal pools and seasonal wetlands</u> – <u>Riparian breeding habitat for migratory birds</u> – <u>Aquatic breeding habitat for amphibians and fish</u> – <u>Foraging habitat for wide-ranging mammals along river corridor</u>
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	<ul style="list-style-type: none"> – <u>Continuous drainage from desert to coast through mountain ranges</u> – <u>Co-occurrence of desert, coastal, and montane species within short distances</u> – <u>Strong desert-coastal ecological transition zone</u>

Criterion		Status	Justification
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Extensive intact alluvial fan sage scrub</u> – <u>High-quality riparian forest and woodland</u> – <u>Bigcone Douglas-fir-canyon oak forest</u> – <u>Remaining natural vernal pools in the County</u> – <u>Large biotically intact river corridor</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Major east-west wildlife movement corridor across the County</u> – <u>Continuous riparian linkage from Pacific coast to Antelope Valley</u> – <u>Tributary connections to San Gabriel Mountains, Santa Susana Mountains, and Liebre/Sierra Pelona</u> – <u>Movement corridors through Sand Canyon, Bouquet Canyon, and San Francisquito Creek</u> – <u>Linkages across State Route 14 via Agua Dulce and Los Piñetos crossings</u>

Santa Monica Mountains SEA

General location and landform

The Santa Monica Mountains SEA is located within the Santa Monica Mountains in a mostly unincorporated area of the County. Much of the area is in public ownership within the Santa Monica Mountains National Recreation Area, but many parcels, particularly smaller ones, are privately owned. Large California State Parks holdings, notably Malibu Creek State Park and Topanga State Park, are included in the SEA. The SEA includes nearly all the canyons and ridges from the Ventura-Los Angeles County line east through Topanga State Park to the Sullivan Canyon area, a large north-south-trending drainage that essentially divides the communities of Pacific Palisades Brentwood to the south, and Encino to the north.

This SEA recognizes the entire extent of a small, somewhat isolated mountain range straddling coastal and interior valley habitats, with a high diversity of topography and moisture regimes. Its natural

DRAFT

vegetation, which still dominates the SEA, is adapted to a Mediterranean climate, which itself is globally rare, existing elsewhere only along western portions of continents at 30 – 40° latitude.

Although the habitats may seem common within the Santa Monica Mountains, the limited global ranges of many of its constituent species, their special adaptations to climate, as well as the relatively intact character of the habitats and the diverse plant assemblage of the Santa Monica Mountains, make it unique. Development pressure within and bordering the SEA has been intense for many decades, so this needs to be carefully balanced to preserve these special resources as an intact ecosystem.

The southern boundary of the SEA reaches the northern boundary of the Malibu Coastline CRA, which extends offshore about a 0.8 mile to capture the kelp beds and rich natural marine communities of and off the coastline of the Santa Monica Mountains. The sandy beaches along this SEA are the least disturbed beaches of the County, some of them completely natural.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Thousand Oaks, Calabasas, Canoga Park, Triunfo Pass, Point Dume, Malibu, and Topanga.

Within the SEA boundary, small areas are not a part of the SEA due to dense development. For example, La Sierra, Malibu Lake, and most of the City of Malibu have been excluded. The majority of the SEA lies within the unincorporated area of the County. Other jurisdictions include the City of Malibu, the City of Los Angeles, City of Calabasas, the City of Agoura Hills, the City of Hidden Hills, and the City of Westlake Village.

Biological Resources

Vegetation and communities

Beginning at the intersection of the Ventura-Los Angeles County line and State Route 1, the southern area of the SEA includes all the terrestrial area of Leo Carrillo State Beach (a State Park with extensive grasslands, coastal chaparral, and an excellent riparian area of sycamore-oak forest along Arroyo Sequit). The Arroyo Sequit from the coast to the junction of West and East forks and about a 0.5 mile of the West Fork has been federally-designated as Critical Habitat for the southern steelhead (*Oncorhynchus mykiss irideus*), which is a salmonid that runs upstream for breeding, but spends most of its life in the ocean. Lower Malibu Creek is also considered Critical Habitat for the southern steelhead. Unfortunately, both runs have been absent for many years, leaving lower Topanga Canyon as the only extant run in the SEA (and in all of Los Angeles Co.); however, it is hoped that future restoration and restocking efforts could result in a return of these runs.

The SEA includes the riparian forest of Zuma Canyon as it emerges on the coast, with a small sub-watershed designated Critical Habitat for the federally-endangered Braunton's milkvetch (*Astragalus brauntonii*). The upstream areas of Santa Ynez, Pulga, and Temescal canyons that are uphill from the community of Palisades Highlands also include federally-designated critical habitat for the milkvetch. This rare plant, which thrives in the years immediately after fire and (under certain circumstances) soil disturbance, is narrowly endemic to four mountain ranges of southern California that have a particular marine-originated rock stratum of interbedded carbonates and sandstones.

While just outside the SEA, the mouth of Malibu Creek, Malibu Lagoon, was completely re-engineered as a fully-tidal lagoon in 2013 and replanted with native vegetation. It is designated critical habitat for the endangered tidewater goby (*Eucyclogobius newberryi*), a small fish that also persists at Topanga

DRAFT

Lagoon (which is part of the SEA). Malibu Lagoon is one of the only remaining coastal estuaries in the County, and the perennial stream that feeds it (greatly augmented by releases from a wastewater treatment upstream at Monte Nido) helps maintain the brackish conditions that are essential to a diverse and unique group of species of coastal estuaries.

In the far eastern portion of the SEA, Temescal, Sullivan and Rustic canyons are deep, mesic drainages that support some of the last populations of uncommon riparian-dependent plants and wildlife close to Los Angeles, including California newt (*Taricha torosa*), two-striped garter snake (*Thamnophis hammondi*) and white-veined monardella (*Monardella hypoleuca* ssp. *hypoleuca*). These species are also present in the cooler, more remote canyons of the central Santa Monicas, which have also supported recent efforts to reestablish the federally threatened California red-legged frog (*Rana draytonii*). Several special-status reptile species are still common within the SEA, including coast horned lizard.

An important feature of this area is a small tributary of Las Virgenes Creek (and ultimately Malibu Creek) named Liberty Canyon. The underpass of State Route 101 at Liberty Canyon Road along the drainage conveys relatively less vehicular traffic than other freeway crossings within several miles, and is one of the few active wildlife passage areas along the entire extent of State Route 101 through the Santa Monica Mountains, with undeveloped habitat on either side. The Wallis Annenberg Wildlife Crossing (scheduled to be completed in late 2026) was cited near here to further take advantage of this critical passage area, particularly for the mountain lion (*Puma concolor*), whose local population was recently designated as state Endangered.

Locally-important valley oak (*Quercus lobata*) savannah may be found on both sides of the freeway at Liberty Canyon. To the west, the unnamed ridge at the southern end of the City of Agoura Hills extends southwest towards Ladyface Mountain, a chaparral-dominated area with numerous canyons and ridges supporting federally-threatened Agoura Hills dudleya (*Dudleya cymosa* ssp. *agourensis*), and one of the nine critical habitat areas in the County of Los Angeles and in the SEA for the state- and federally-endangered Lyon's pentachaeta (*Pentachaeta lyonii*), as well as other uncommon and rare plants including a pale violet *Navarretia* of uncertain identity (form *pubescens*), but seemingly closely related to *N. mitracarpa* and *N. ojaiensis*.

The area also supports the southernmost known occurrences within the County of several species more indicative of arid or semi-arid interior basins, including California juniper (*Juniperus californica*), linear-leaved goldenbush (*Ericameria linearis*), Kern Brodiaea (*Brodiaea terrestris* ssp. *kernensis*), and butterfly mariposa lily (*Calochortus venustus*). The area near Las Virgenes Reservoir toward the Ventura County line supports another high concentration of locally-rare plants, and is also Critical Habitat for the Lyon's pentachaeta. Other important areas for rare plants include Malibu Creek State Park (particularly for rock outcrop species, like marcescent dudleya *Dudleya cymosa* ssp. *marcescens*) and Red Rock Canyon Park north of Topanga for species adapted to unusual soil types, including Santa Susana tarplant (*Deinandra minthornii*).

Excellent examples of state-rare vegetation communities are still found here, such as southern California coastal lagoon, southern California steelhead stream, southern sycamore alder riparian woodland, and valley needlegrass grassland. Relatively isolated canyons such as Zuma Canyon represent some of the last major drainages in the Los Angeles County portion of the Santa Monica Mountains with a natural year-round stream that supports a rich riparian community and remains in an undeveloped state. Cold Creek includes an excellent example of an undisturbed natural sandstone basin with springs and a perennial stream, and Tuna and Pena canyons are the last drainages in the central and eastern Santa Monica Mountains that have essentially no development between the

DRAFT

canyon mouth at the coast and upper areas of the watershed.

Wildlife Movement

The SEA contains major features that are important to wildlife movement, both within the Santa Monica Mountains, and as crossings north into larger areas of open space in the local mountains. A wide variety of wildlife use linkages throughout the SEA, including mountain lion, coyote, mule deer, bobcat, and essentially all the other medium-sized mammals of southern California. Aside from the Liberty Canyon area discussed above, on the eastern end of the mountains, Mulholland Drive bridge over the 405 Freeway and Bel Air Crest Rd. just to the south, are the only feasible ways wildlife transit between the Santa Monica Mountains sections west and east of Sepulveda Pass. The many natural drainages and ridgelines of the Santa Monica Mountains connect populations of wildlife within the range, where for now, development is sparse enough that barriers to movement are mainly in the form of roads, and increased vehicle traffic on these roads.

Malibu Creek State Park has been identified as the central core habitat area in the Santa Monica Mountains, serving as a connective hub between the Simi Hills to the north and the open space preserves of Topanga State Park to the east, and Muqu State Park to the west (which is near the Ventura-Los Angeles County line, but within Ventura County). These linkages still allow movement among large open space areas within the SEA as well as between areas outside the SEA, such as the Simi Hills and the western extent of the Santa Monica Mountains in Ventura County. The genetic flow through these areas is crucial in maintaining the diversity and viability of the species within the Santa Monica Mountains. Open space linkages between Kanan Road and Calabasas Parkway along State Route 101, as indicated by the National Park Service, are of particular importance for continued connectivity of wildlife populations, due to a lack of alternative routes and encroachment of development. Although there are significantly large open spaces within the SEA, maintaining habitat linkages between them is critical in providing for long-term sustainability.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE SANTA MONICA MOUNTAINS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> - <u>California red-legged frog</u> - <u>Mountain lion (Southern California/Central Coast ESU)</u> - <u>Braunton’s milk-vetch</u> - <u>Lyon’s pentachaeta</u> - <u>Santa Monica dudleya</u> - <u>Agoura Hills dudleya</u> - <u>Marcrescent dudleya</u> - <u>Malibu baccharis</u>

DRAFT

Criterion		Status	Justification
B/C)	<p><u>On a regional/Countywide basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u></p>	Met	<ul style="list-style-type: none"> – <u>Roughly two dozen special-status species (excluding Threatened and Endangered)</u> – <u>Rare riparian communities in coastal Mediterranean climate setting</u> – <u>Floristically diverse canyon and rock outcrop habitats</u> – <u>southern California coastal lagoon and steelhead stream communities</u> – <u>Valley needlegrass grassland and sycamore–alder riparian woodland</u>
D)	<p><u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u></p>	Met	<ul style="list-style-type: none"> – <u>Waterbird rookeries at Westlake Reservoir and Malibu Lagoon</u> – <u>Migratory bird stopover habitat at Malibu Lagoon and Malibu Creek</u> – <u>Narrow habitat linkages between large open space areas</u> – <u>Riparian breeding habitat for birds and amphibians</u> – <u>Malibu Lagoon and the upstream riparian woodland in Malibu Creek are important bird stopover habitats.</u>

DRAFT

Criterion		Status	Justification
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Mediterranean-climate mountain range with isolated coastal influence</u> – <u>Range limits for species such as California juniper and valley oak</u> – <u>Unusual floristic diversity associated with varied geology and microclimates</u> – <u>Isolated mountain range supporting coastal and interior species assemblages</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Undisturbed canyon-bottom riparian habitats (e.g., Zuma Canyon)</u> – <u>Intact chaparral and coastal sage scrub communities</u> – <u>Undeveloped perennial stream systems</u> – <u>Large contiguous natural habitat mosaic</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Linkages within Santa Monica Mountains along ridgelines and drainages</u> – <u>Liberty Canyon wildlife crossing and surrounding movement corridor</u> – <u>Connectivity between Santa Monica Mountains, Simi Hills, and Ventura County open space</u> – <u>Movement corridors across State Route 101 and Sepulveda Pass crossings</u>

Santa Susana Mountains and Simi Hills SEA

General location and landform

The Santa Susana Mountains and Simi Hills SEA is located northwest of the San Fernando Valley within unincorporated areas of the County and the City of Los Angeles, west of the communities of

DRAFT

Chatsworth and West Hills and north of Calabasas. It includes much of the Santa Susana Mountains to the north, the Santa Susana Pass, Chatsworth Reservoir, and the eastern portion of the Simi Hills to the south. The north face of the Santa Susana Mountains comprises the southwestern watershed of the Santa Clara River, and on the south face, the (Los Angeles County portion of the) Santa Susana Mountains are part of the direct coastal watershed of the Los Angeles River. The Simi Hills in Ventura County are part of the direct coastal drainage in their southernmost area (via Las Virgenes and Malibu canyons).

The SEA extends eastward to include all of the Santa Susana Pass area, much of which is preserved in the Santa Susana Pass State Historic Park. Just across the Ventura-Los Angeles County line near State Route 118 in Ventura County are Corriganville Park, a former and current natural location for movie production, and Rocky Peak Park. Corriganville is a regional park within the City of Simi Valley; Rocky Peak Park is administered by the Santa Monica Mountains Conservancy. Rocky Peak Park is the former Runkle Ranch and stretches from State Route 118 five miles northward to Las Lajas Canyon. The area is a vital wildlife habitat linkage between the Simi Hills and the Santa Susana Mountains. Spectacular sandstone boulders, outcroppings, oak savannahs, and perennial water sources provide diverse habitat for vertebrates and a number of rare plants. The Ventura-Los Angeles County line and the SEA boundary cross directly over Rocky Peak here. The majority of the land in the SEA is natural open space with very sparse disturbances in the form of ranches, oil wells, and narrow dirt roads.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Calabasas, Canoga Park, Oat Mountain, Santa Susana, Val Verde, and Newhall.

Biological Resources

Vegetation and communities

The southern side of Dayton Canyon, southwest of Chatsworth, is designated Critical Habitat for the Braunton's milk vetch (*Astragalus brauntonii*), which co-occurs with another rare plant, chaparral nolina (*Nolina cismontana*). The SEA also includes the unique habitats of the Chatsworth Nature Preserve (including Chatsworth Reservoir), which supports some of the large seasonal wetlands in the County, along with numerous rare plant and wildlife species with populations greater than anywhere else in the County, including western spadefoot (*Spea hammondi*) and southern tarplant (*Centromadia parryi* ssp. *australis*).

While portions of the SEA (from State Route 118 northward) have been designated Critical Habitat for the coastal California gnatcatcher (*Polioptila californica californica*), there are few confirmed records from here, with the bulk of the population lying to the southwest, within Ventura County.

North across the Interstate-5 is Newhall Pass, a rugged part of the Santa Susana Mountains near the junction of State Route 14, with substantial natural vegetation of oak woodlands, chaparral, and coastal sage scrub. On the west side of Sunshine Canyon Landfill, a broad lobe of the SEA extends along the ridgeline, which separates Sunshine and Bee Canyon to include Bee Canyon Park and Mission Point of O'Melveny Park. This is an area with walnut woodlands, oak woodlands, grasslands, and chaparral, including coastal sage scrub, which is the diverse and green vegetation typical of the Santa Susana Mountains. Several perennial stream areas occur within these canyons, and there are many natural springs and even small vernal pools on flatter portions.

Open space within the SEA supports a great variety of communities dominated by chaparral, oak

DRAFT

woodlands, coastal sage scrub, bigcone Douglas-fir-canyon oak woodland at the tops of deeper canyons, and native forb-rich grasslands, where they occur well below their normal elevational limit in southern California. Other oaks with groves in the Santa Susanas include coast live oak (*Quercus agrifolia*), valley oak (*Q. lobata*), canyon live oak (*Q. chrysolepis*), scrub oak (*Q. berberidifolia*), and interior live oak (*Q. wislizeni*).

This area supports localized but core populations of the Santa Susana tarplant (*Deinandra minthornii*), which is a tarplant like few others since it is perennial. Its distribution spreads through the Simi Hills and into the Santa Monica Mountains, but it is primarily at home among the sandstone boulders and terraces that prevail in the Santa Susana Mountains. The recently (1996)-described Malibu baccharis (*Baccharis malibuensis*) and Payne's bush lupine (*Lupinus paynei*) are two other narrowly-endemic plants with ranges centered on this SEA.

The numerous creeks and canyons support riparian scrub and woodland communities with oaks, sycamores, and willows. Not uncommon are cherry and California ash woodlands, which are dominated by holly-leaf cherry (*Prunus ilicifolia*). There are walnut woodlands of the California black walnut (*Juglans californica*) mixed with flowering ash (*Fraxinus dipetala*) and blue elderberry (*Sambucus mexicana*) and coast live oak. Flowering ash may be a tree up to 60 feet tall in the Santa Susana Mountains, whereas it usually is a low tree or even spindly shrub. The woodlands dominated by walnuts and flowering ash appear to be unique to the Santa Susana Mountains.

Wildlife Movement

The SEA includes several important linkages for wildlife movement. The Simi Hills and Santa Susana Mountains provide a vast open space corridor to foster wildlife movement between the Santa Monica Mountains to the south, San Gabriel Mountains to the east, and Los Padres National Forest to the north in the western San Gabriel Mountains of the Transverse Ranges. Dense, natural habitat associated with the majority of the SEA provides excellent opportunities for concealment and water sources, while the grasslands provide an abundance of prey. Examples of wildlife that use these linkages include mountain lion (*Puma concolor*), mule deer (*Odocoileus hemionus*), bobcat (*Lynx rufus*), and a number of medium-sized animals.

A very important wildlife passage between the Santa Susana Mountains and the Simi Hills is just west of the Ventura-Los Angeles County line connecting Corriganville Park and the Runkle Ranch. It is a broad tunnel under the freeway, which enabled the connection of the property that was divided by constructing State Route 118. This tunnel connects dirt roads and trails on either side, and is regularly used by mountain lions and other wildlife.

The connection to the Santa Clara River SEA is the Los Piñetos Road underpass of the State Route 14. Consistent wildlife movement has been recorded with motion-activated cameras there. The Weldon Canyon Road overpass of the Interstate-5 is another connection for the Newhall Wedge with the main part of the SEA. The Old Road underpass of the Interstate-5 is a broad connection. The Calgrove Avenue underpass is another broad connection, though busy with traffic.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

DRAFT

CRITERIA ANALYSIS SUMMARY OF THE SANTA SUSANA MOUNTAINS AND SIMI HILLS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>Western spadefoot</u> – <u>Mountain lion (Southern California/Central Coast ESU)</u> – <u>Braunton’s milk-vetch</u> – <u>Malibu baccharis</u> – <u>Santa Susana tarplant</u> – <u>Payne’s bush lupine</u> – <u>Chaparral nolina</u> – <u>Southern tarplant</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>Coastal lowland populations of greater roadrunner, hairy woodpecker, and coastal Bell’s sparrow</u> – <u>Canyon live oak ravine forest</u> – <u>Mainland cherry forest</u> – <u>Southern coast live oak riparian forest</u> – <u>Valley oak woodland</u> – <u>Walnut-flowering ash woodland unique to Santa Susana Mountains</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Vernal pool and seasonal wetland complex at Chatsworth Nature Preserve</u> – <u>Breeding habitat for western spadefoot and southern tarplant</u> – <u>Seasonal wetlands supporting rare wetland-dependent species</u> – <u>Concentrated amphibian breeding habitat</u>

DRAFT

Criterion		Status	Justification
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Low-elevation populations of bigcone Douglas-fir</u> – <u>Low-elevation canyon live oak communities</u> – <u>Unusual sandstone-associated endemic plant assemblages</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Large contiguous chaparral, oak woodland, and grassland communities</u> – <u>Undisturbed canyon riparian habitats</u> – <u>Intact walnut woodland and coastal sage scrub mosaic</u> – <u>Extensive natural open space with minimal disturbance</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Regional linkage between Santa Monica Mountains, San Gabriel Mountains, and Los Padres National Forest</u> – <u>Wildlife passage under State Route 118 connecting Simi Hills and Santa Susana Mountains</u> – <u>Connectivity to Santa Clara River SEA via Los Piñetos underpass</u> – <u>Movement corridors across Interstate 5 via Weldon Canyon and Calgrove crossings</u>

Sepulveda Basin SEA

General location and landform

The Sepulveda Basin comprises more than 2,000 acres of mostly undeveloped land including and alongside the Los Angeles River in the middle of the San Fernando Valley. It occupies the confluences of several tributaries of the Los Angeles River, Bull Creek, Encino Creek and Haskell Creek, and supports a mix of various bottomland vegetation communities largely lost on the floor of the Basin. It is isolated from the Santa Monica Mountains by a major freeway and residential development; however, Encino Creek provides at least hydrological connection between the Basin and that range.

DRAFT

A variety of ownership and management overlays cover Sepulveda Basin, including US Army Corps of Engineers (owns most of the land) and the city of Los Angeles (manages most of the land, with the exception of the “South Reserve” south of Burbank Blvd.).

The SEA is located within the Canoga Park and Van Nuys United States Geological Survey (USGS) 7.5' California Quadrangles.

Biological Resources

Vegetation and communities

The Sepulveda Basin, one of a handful of large remaining open space areas on the floor of the Los Angeles Basin, is a mixed-use recreational area with considerable natural habitats, including:

- A 1-mile natural-floor stretch of the Los Angeles River, with native willow-dominated riparian vegetation;
- “Bull Creek Restoration Area”, completed in 2009, located just north and west of Lake Balboa;
- Several marsh-lined ponds located in the designated “wildlife area”, as well as on the Woodley Lakes Golf Course;
- Multiple active colonial waterbird rookeries;
- Natural riparian scrub vegetation with scattered Fremont cottonwood and other trees in the “South Reserve” area south of Burbank Blvd.;
- A large constructed (cement-lined) reservoir (Lake Balboa) that supports waterbirds in winter.

Out of the 2,000+ acres within the Sepulveda Basin, 285 acres consisting of Bull Creek, the Los Angeles River, and Haskell Creek with its associated wildlife lake and adjacent undeveloped uplands, are recommended for SEA designation. While physically isolated from other open space areas of the County, its location along the Los Angeles River makes it a crucial component of future restoration efforts (The River Project 2022).

The Los Angeles River channel itself and associated habitats support Fremont cottonwood (*Populus fremontii*), and willows (*Salix* spp.), along with mulefat (*Baccharis salicifolia*) and emergent freshwater marsh vegetation, such as California bulrush (*Schoenoplectus californicus*) and broad-leaf cattail (*Typha latifolia*). This habitat supports numerous wildlife species found only in freshwater marsh, including rails (family Rallidae) and dragonflies/damselflies (order Odonata). Sanford’s arrowhead (*Sagittaria sanfordii*), a CRPR 1B.2 plant, was recently discovered along the Los Angeles River channel, and numerous wetland plant species occur within the basin (Cooper et al. 2022). Since formal botanical surveys have never been conducted here, more may await discovery.

Sensitive and noteworthy wildlife species confirmed as nesting at Sepulveda Basin include the Endangered least Bell’s vireo (*Vireo bellii pusillis*) and multiple California Species of Special Concern, including the least bittern (*Botarus exilis*), Clark’s marsh wren (*Cistothorus palustris clarkae*), yellow-breasted chat (*Icteria virens*) and northern yellow warbler (*Setophaga aestiva*).

A large rookery consisting mainly of double-crested cormorant (*Nannopterum auritum*) and great blue heron (*Ardea herodias*) is located on a small island within the Wildlife Area, and a smaller great blue heron rookery is located adjacent to the Los Angeles River. Additional rookeries of herons and egrets

DRAFT

have been found scattered throughout the Basin, appearing unpredictably from year to year.

The Los Angeles River includes an unchannelized portion, with natural sand/silt banks, just south of Woodley Golf Course. This fragment represents one of the few remaining large examples of a natural riparian zone in the coastal lowlands. Upland slopes of the Los Angeles River levee have been restored with native coastal sage scrub species, such as California buckwheat (*Eriogonum fasciculatum*), coyote brush (*Baccharis pilularis*), and California sagebrush (*Artemisia californica*). American kestrel, an open country species now rare year round, has bred in the Basin at least once in the past 10 years. In grassland areas, both burrowing owl (*Athene cunicularia*) and loggerhead shrike (*Lanius ludovicianus*) are regular in fall and winter, one of the few locales in urban Los Angeles where they still occur.

Wildlife Movement

The Los Angeles River channel represents a major urban wildlife corridor through the urban Los Angeles Basin.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE SEPULVEDA BASIN SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> - <u>Least Bell's vireo</u> - <u>Sanford's arrowhead</u>
B/ C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> - <u>Freshwater marsh habitat</u> - <u>Rare unchannelized river habitat</u> - <u>Least bittern</u> - <u>Clark's marsh wren</u> - <u>Yellow-breasted chat</u> - <u>Northern yellow warbler</u> - <u>Rails (Rallidae)</u> - <u>Odonates (dragonflies and damselflies)</u>
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> - <u>Colonial waterbird rookeries</u> - <u>Breeding marsh birds</u> - <u>Wintering waterbirds</u>

Criterion		Status	Justification
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Not Met	
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> - <u>Unchannelized Los Angeles River segment</u> - <u>Natural riparian zone in coastal lowlands</u>
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> - <u>Los Angeles River wildlife corridor</u> - <u>Linkage toward Santa Monica Mountains</u> - <u>Basin floor movement corridor</u>

Tujunga Valley and Hansen Dam SEA

General location and landform

The Tujunga Valley and Hansen Dam SEA is located on the northern edge of the San Fernando Valley at the base of the Verdugo Mountains and the foothills of the San Gabriel Mountains. The SEA consists of the Tujunga Valley and Wash, starting in the riparian areas of the Big Tujunga, which is a main tributary of the Los Angeles River, and including Hansen Dam Flood Control Basin/Hansen Dam Park. The SEA is largely within the City of Los Angeles. The area is an Audubon California Important Bird Area (IBA), which is part of the Los Angeles Flood Control Basin IBA. The area southwest of the dam is used as a spreading ground. This has created several freshwater marsh areas that are used by marsh birds, migratory waterfowl, and shorebirds.

The SEA is located within portions of the United States Geological Survey (USGS) 7.5' California Quadrangles: San Fernando, Sunland and Van Nuys.

Biological Resources

Vegetation and communities

The wash, golf course, and reserve area support an unusual remnant vegetation of alluvial fan scrub, with California juniper (*Juniperus californica*), patches of prickly-pear (*Opuntia littoralis*) and cane cholla cactus (*Cylindropuntia bernardina*), along with the usual riparian and scrub plants. Large natural

DRAFT

areas of riparian vegetation remain on the banks of the wash, as well as along associated tributaries (including Haines Canyon and Little Tujunga Creek), including what may be the largest extent of southern cottonwood-willow riparian forest left in the Los Angeles Basin.

The extensive alluvial fan scrub, which once covered the base of the foothills across the northern edge of the Los Angeles Basin, has been nearly extirpated by development and flood control projects. It is only represented in a few places in the County, including here, Pacoima Wash, the San Gabriel Canyon SEA, and San Antonio Wash. The alluvial fan habitat mixed with riparian forest continues across the Interstate-210 as the Big Tujunga Wash approaches and flows into Hansen Dam Park. The SEA has a broad undercrossing of the Interstate-210, which includes a braid of its tributary Haines Canyon Creek crossing under Foothill Boulevard next to Wentworth Street. The Wentworth Street undercrossing is important to wildlife, as it connects to somewhat dispersed development in the Shadow Hills neighborhood and from there to the natural areas of the Verdugo Mountains and the Verdugo Mountains SEA. Mountain lions are resident in the Verdugo Mountains, making Big Tujunga/Hansen Dam an important connection for that population as well as for other animals and plants of the Verdugo Mountains.

The Hansen Dam area is recognized for its great importance to migrating birds on the Pacific Flyway (nearly 300 species of birds have been recorded here), as well as the very rare habitat of alluvial fan scrub, which has uncommon resident birds. A coastal-slope population of the cactus wren (*Campylorhynchus brunneicapillus*) is resident here, but is nearing extirpation. Recent records of coastal California gnatcatcher suggest incipient occupancy by this federally threatened species. Coastal-lowland populations of birds such as greater roadrunner, hairy woodpecker, lesser nighthawk, and possibly breeding Swainson's thrush are also notable, as they have little habitat left away from the deserts and mountains. Other potentially occurring rare wildlife here include the County's last coastal slope population of California glossy snake (*Arizona elegans occidentalis*) and arroyo toad (*Anaxyrus californicus*), both of which require broad, sandy alluvial washes now rare in the region.

The wash flows year round most years, and the area from Hansen Lake (in the Hansen Dam Park) upstream to beyond the SEA is Critical Habitat for the federally-threatened Santa Ana sucker (*Pantosteus santaanae*). Two other fish Species of Special Concern are found here, speckled dace (*Rhinichthys gabrielino*) and arroyo chub (*Gila orcuttii*), which together with rainbow trout, comprise most of the original fish fauna of the Los Angeles River watershed (the unarmored threespine stickleback, *Gasterosteus aculeatus williamsoni*, is extirpated). Other riparian/aquatic species found here include a large population of breeding least Bell's vireo (*Vireo bellii pusillus*) and the southwestern pond turtle (*Actinemys pallida*) in deeper pools such as at Haines Canyon.

East and adjacent to the large Angeles National golf course is a state reserve area for the state and federally-endangered slender-horned spineflower (*Dodecahema leptoceras*), a rare buckwheat relative restricted to alluvial washes west of the deserts, and Davidson's bush-mallow (*Malacothamnus davidsonii*), a flowering shrub also known as Tujunga bush-mallow owing to its highly restricted range.

Wildlife Movement

Although wildlife movement is hampered by development surrounding the SEA to the south, animals are still able to move through the adjacent hills and valleys to the north into the San Gabriel Mountains, and (tenuously) into the Verdugo Mountains to the south. Due to its large size and topographic complexity, many linkages are certain to occur within the SEA at various bottlenecks. These linkages allow movement between large open space areas within the SEA. Although there are significantly large open spaces within the SEA, contiguous habitat linkages between them are critical in reducing

DRAFT

bottlenecks and providing for long-term sustainability. A wide variety of wildlife use linkages throughout the SEA, including mountain lion (*Puma concolor*), coyote, mule deer, bobcat (*Lynx rufus*), and a number of medium-sized mammals. The Wentworth Street underpass of Interstate-210 is a conduit for wildlife through the dispersed residential neighborhoods of the Shadow Hills. The Shadow Hills are the westernmost extent of the Verdugo Mountains, which provide a large area of natural habitat for wildlife.

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE TUJUNGA VALLEY AND HANSEN DAM SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>Least Bell’s vireo</u> – <u>Santa Ana sucker</u> – <u>Santa Ana speckled dace</u> – <u>Mountain lion (Southern California/Central Coast ESU)</u> – <u>Southwestern pond turtle</u> – <u>Slender-horned spineflower</u> – <u>Davidson’s bush-mallow</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>Remnant alluvial fan scrub at base of San Gabriel foothills</u> – <u>One of the largest remaining southern cottonwood-willow riparian forests in Los Angeles Basin</u> – <u>Extensive riparian woodland and marsh habitats within urbanized landscape</u> – <u>Rare sandy alluvial wash habitat supporting specialized species</u> – <u>Freshwater marsh habitat created by spreading grounds</u>

DRAFT

Criterion		Status	Justification
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Important migratory bird stopover along Pacific Flyway (≈300 species recorded)</u> – <u>Breeding habitat for least Bell’s vireo</u> – <u>Aquatic habitat supporting native fish assemblage of Los Angeles River watershed</u> – <u>Amphibian and reptile breeding and foraging habitat in alluvial wash systems</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Large intact alluvial wash habitat rare in Los Angeles Basin</u> – <u>Coastal-slope alluvial fan scrub remnants at urban edge</u> – <u>Unusual persistence of desert-affiliated species in coastal basin setting</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Relatively intact Big Tujunga Wash alluvial fan system</u> – <u>Undisturbed riparian forest along perennial wash segments</u> – <u>Large natural flood control basin supporting native vegetation</u> – <u>Remnant alluvial fan communities representative of historical landscape</u>

Criterion		Status	Justification
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none"> – <u>Movement corridor between San Gabriel Mountains and Verdugo Mountains</u> – <u>Wildlife passage under Interstate 210 at Wentworth Street</u> – <u>Linkages along Big Tujunga Wash and tributaries</u> – <u>Connectivity for mountain lion and other wide-ranging mammals</u>

Verdugo Mountains SEA

General location and landform

The Verdugo Mountains SEA encompasses the Verdugo Mountains south of Interstate-210 and east of the Interstate-5, as well as a portion of these mountains north of Interstate-210, north into the Shadow Hills neighborhood south of Big Tujunga Wash. The Verdugo Mountains are a wilderness island in the middle of the urbanized metropolitan area of the County, surrounded by the cities of Los Angeles, Burbank and Glendale. The Verdugo Mountains, and surrounding neighborhoods like Sunland and Shadow Hills, have retained a rural atmosphere despite their proximity to urban Los Angeles.

The freeway frontage includes relatively large areas of native vegetation on either side and a handful of connections across Interstate-210, all of which are threatened with development or otherwise impacted. The dispersed equestrian ranchettes in the Shadow Hills neighborhood feature important fragments of native vegetation. A paved road (La Tuna Canyon Rd.) runs alongside the streamcourse in La Tuna Canyon, and here the SEA includes about two miles of native, natural vegetation bordered on the eastern, undeveloped section of the road.

The Verdugo Mountains have been impacted by dirt roads, one paved road (Tuna Canyon), as well as fuel breaks, transmission lines and isolated buildings for houses, radio towers, and water tanks. Because of their extent, however, the Verdugo Mountains are still considered largely natural and little impacted—a prime example of the chaparral and coastal sage chaparral scrub once prevalent in the County coastal areas.

The SEA is located at least partially in each of the following United States Geological Survey (USGS) 7.5' California Quadrangles: Burbank, Sunland, and Pasadena.

It is wholly within incorporated boundaries (cities of Los Angeles, Glendale, and Burbank), but much is preserved in conservation easements under the guidance of the Santa Monica Mountains Conservancy.

Biological Resources

Vegetation and communities

The Verdugo Mountains currently encompass wilderness area, which ranges through various chaparral, coastal sage chaparral scrub, southern willow scrub, coast live oak woodland and forest ecosystems, and many riparian areas with seasonal waterfalls. It is one of the few remaining natural regions in the urban Los Angeles area that supports abundant native wildlife and habitats, and also contains several rare and sensitive plant and animal species. Many species found in the San Gabriel foothills reach the edge of their ranges into the Los Angeles Basin here, rather than crossing the lowlands. In this way, the Verdugo Mountains represent a crossroads between the Transverse and Peninsular Ranges, and the Santa Monica Mountains to the west, supporting rare lowland populations of mountain lion, Merriam's chipmunk (*Neotamias merriami*) and western gray squirrel (*Sciurus griseus*).

The SEA may be thought of as a habitat island, providing what remains of a link between plant and animal populations found in the Santa Monica and San Gabriel mountains. Genetic interchange, by way of this linkage is important in perpetuating the genetic variability in isolated populations, and the maintenance of healthy ecosystems. Chaparral and coastal sage chaparral scrub cover the hillsides of the mountains, with lush riparian vegetation, including California bay (*Umbellularia californica*), western sycamore (*Platanus racemosa*), coast live oak (*Quercus agrifolia*), ferns, and ocellated Humboldt lily (*Lilium humboldtii* ssp. *ocellatum*), which are found in most of the stream drainages. The localized range of the localized *pasadenensis* form of summer lupine (*Lupinus formosus*) is centered here, reaching locally into the San Gabriel foothills and the eastern edge of Griffith Park. These plant communities provide habitat that is essential to the diverse and abundant fauna that are found in the area. The mountains are also home to a widely disjunct population of mission manzanita (*Xylococcus bicolor*). Other rare plants present are similar to those found in nearby Griffith Park, including Hubby's phacelia (*Phacelia hubbyi*), Plummer's mariposa-lily (*Calochortus plummerae*), and San Gabriel oak (*Quercus durata* var. *gabrielensis*). However, Braunton's milkvetch (*Astragalus brauntonii*), Brewer's Calandrinia (*Calandrinia breweri*), and chaparral rein orchid (*Piperia cooperi*), appear to be absent from the park, indicating the role of the Verdugos in preserving their populations.

While incompletely documented, the reptile community also appears to be largely intact, with coastal whiptail (*Aspidoscelis tigris stejnegeri*), coast horned lizard (*Phrynosoma blainvillii*), and coast patch-nosed snake (*Salvadora hexalepis virgulata*) present and regularly observed.

Wildlife Movement

Although wildlife movement is hampered by adjacent rural development in proximity to the SEA, animals are still able to move through the Verdugo Mountains in many areas. Due to its large size and topographic complexity, many linkages occur within the SEA at various bottlenecks. These linkages allow movement between large open space areas within the SEA, as well as between areas outside the SEA toward the Angeles National Forest. The genetic flow through these areas is crucial in maintaining the diversity and viability of the species within the Verdugo Mountains. Some areas of probable, possible, and perhaps future connection have been indicated in the General Boundary and Resources section. Although there are significantly large open spaces within the SEA, contiguous habitat linkages between them are critical in reducing bottlenecks and providing for long-term sustainability. A wide variety of wildlife use linkages throughout the SEA, including mountain lion (*Puma concolor*), coyote (*Canis latrans*), mule deer (*Odocoileus hemionus*), and bobcat (*Lynx rufus*).

DRAFT

Designation Criteria

The SEA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS SUMMARY OF THE VERDUGO MOUNTAINS SEA

Criterion		Status	Justification
A)	<u>The habitat of core populations of endangered or threatened plant or animal species.</u>	Met	<ul style="list-style-type: none"> – <u>Mountain lion (Southern California/Central Coast ESU)</u> – <u>Braunton’s milk-vetch</u> – <u>Mission manzanita (disjunct population)</u>
B/C)	<u>On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.</u>	Met	<ul style="list-style-type: none"> – <u>Large intact habitat island of natural vegetation within highly urbanized Los Angeles Basin</u> – <u>Coastal whiptail, coast horned lizard, and coast patch-nosed snake populations in urban mountain setting</u> – <u>Hubby’s phacelia and Plummer’s mariposa-lily occurrences</u> – <u>Chaparral-coastal sage scrub-riparian mosaic uncommon in urban Los Angeles</u> – <u>Range-edge populations of San Gabriel foothill species</u>

DRAFT

Criterion		Status	Justification
D)	<u>Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.</u>	Not Met	<ul style="list-style-type: none"> – <u>Habitat island supporting resident wildlife populations isolated from surrounding urban areas</u> – <u>Riparian canyon habitat providing breeding and foraging areas</u> – <u>Movement stepping-stone between surrounding mountain ranges for dispersal</u> – <u>Refuge habitat for reptiles and small mammals within urban matrix</u>
E)	<u>Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.</u>	Met	<ul style="list-style-type: none"> – <u>Crossroads between Transverse Ranges, Peninsular Ranges, and Santa Monica Mountains</u> – <u>Species occurring at Los Angeles Basin range limits</u> – <u>Disjunct population of mission manzanita</u> – <u>Isolated urban mountain range supporting intact native fauna</u>
F)	<u>Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.</u>	Met	<ul style="list-style-type: none"> – <u>Large contiguous chaparral and coastal sage scrub communities</u> – <u>Intact canyon riparian habitats with minimal disturbance</u> – <u>Representative remnant of coastal basin mountain vegetation</u> – <u>Mostly natural vegetation despite surrounding urban development</u>

DRAFT

Criterion		Status	Justification
G)	<u>Areas that provide functionally important habitat connectivity between or within designated SEAs, including relatively intact or lightly disturbed habitat that forms functional linkages or corridors connecting portions of the same SEA or linking one SEA to another.</u>	Met	<ul style="list-style-type: none">– <u>Linkage between San Gabriel Mountains and Santa Monica Mountains (fragmented but functional for some taxa)</u>– <u>Movement corridor through Shadow Hills to Big Tujunga Wash</u>– <u>Internal ridge and canyon linkages within Verdugo Mountains</u>– <u>Genetic interchange stepping-stone among regional mountain ranges</u>

SEA Description Sources

~~Altadena Foothill and Arroyo SEA Sources~~

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The Altadena SEA Update Study*, Los Angeles County, California.~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Los Angeles County, Department of Parks and Recreation. Date unknown. *Bonelli Regional Park Bird List*, San Dimas, California.~~

~~Quinn, R.P. Date unknown. *The Status of Walnut Forests and Woodlands (Juglans californica) in Southern California*, Department of Biological Sciences, California State Polytechnic University, Pomona, California.~~

~~Sawyer, Keeler Wolf and Evens. 2009. *A Manual of California Vegetation*, Second Edition Sacramento: California Native Plant Society Press.~~

~~Toyon Design, Landscape Architecture. August 5, 2004. *Biological Survey and Habitat Needs Analysis for the Altadena Foothills, Los Angeles County, CA*. Prepared for the Altadena Foothills Conservancy~~

~~Antelope Valley SEA Sources~~

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

DRAFT

~~CalMat Properties Co.1988.A Plan for Reclamation:Big Rock Creek, Attachment:Technical Supplement, SEA No. 48, Los Angeles County, California.~~

~~County of Los Angeles, Department of Regional Planning.2006.The Antelope Valley SEA Description, Los Angeles County, California.~~

~~County of Los Angeles, Department of Regional Planning.1994.Mescal Creek Surface Mining and Reclamation Plan, Preliminary Draft Environmental Impact Report, Los Angeles County, California.~~

~~Edwards Air Force Base.1999.Checklist of Birds.Two page leaflet.~~

~~Envicom Corporation.1999.Biological Constraints Analysis for 4 Aces FilmLocations Property, SEA Nos. 48 & 53, Los Angeles County, California.~~

~~Fishman, R.1988.Biological Survey and Impact Assessment for Tentative Parcel No. 19426, 150th St. East and Avenue O, Lake Los Angeles, SEA No. 53 (revised), Los Angeles County, California.~~

~~Fishman, R.1988.Biological Survey and Impact Assessment for Tentative Parcel No. 19426, 150th St. East and Avenue O, Lake Los Angeles, SEA No. 53, Los Angeles County, California.~~

~~Frank Hovore & Associates.2000.County of Los Angeles Biological Constraints Analysis, Birsner Property, Crystalaire, P.M. 25189.~~

~~Frank Hovore & Associates.2000.Antelope Valley Courthouse Site, Biological Impacts Assessment.~~

~~Frank Hovore & Associates.1996.Palmdale Water District, 1996 Water System Master Plan, Biological Constraints Assessment.~~

~~Furgo-McClelland (West), Inc.1992.Draft Biota Report for Surface Mining Permit 90296, SEA No. 55, Mescal Creek (revised), Ventura, California.~~

~~Furgo-McClelland (West), Inc.1992.Draft Biota Report for Surface Mining Permit 90296, SEA No. 55, Mescal Creek, Ventura, California.~~

~~Hanes, T. and J. Weintraub.1989.Biota and Project Plans for Tentative Tract No. 40248, Puzzle/Jesus Creek Area, SEA No. 55, Pearblossom, California.~~

~~Hanes, T. and J. Weintraub.1988.Biological Survey of a 380 Acre SEA, Puzzle/Jesus Creek Area, SEA No. 55, Pearblossom, California.Department of Biology, Cal State Univ. Fullerton.~~

~~Holland RF. Preliminary descriptions of the terrestrial natural communities of California. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Impact Sciences, Inc.1992.Biota Report for Tentative Tract Map 50804, SEA Nos. 48 and 53, Los Angeles County, California.~~

~~Impact Sciences, Inc.1991.Biological Constraints Analysis for Lake Los Angeles Site, SEA Nos. 51-54, Los Angeles County, California~~

~~Mattoni, R.Butterflies of Greater Los Angeles.Wall poster, 2 sides.~~

~~Michael Brandman Associates.1989.Biological Resources Assessment, Assessor's Parcel No. 3029-~~

DRAFT

~~12-08, SEA No. 53, Los Angeles County, California.~~

~~Michael Brandman Associates. 1989. *Results of Survey for the Mohave Ground Squirrel (Spermophilus mohavensis) on Tentative Parcel No. 19426*, Los Angeles County, California.~~

~~Rado, T. October 1991. *Biological Constraints Analysis for Tentative Tract No. 48746, SEA No. 52*, Los Angeles County, California.~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press~~

~~Stephenson, J.R. and G.M. Calcarone, 1999. *Southern California Mountains and Foothills Assessment: Habitat and Species Conservation Issues*. Gen. Tech. Rep. GTR-PSW-172. Albany, CA. Pacific SW Res. Sta., Forest Serv., U.S. Department Agric., 402 pp.~~

~~VTN West, Inc. 1988. *Biota Report for Tentative Tract 45238, Parcel Map 19161, SEA No. 49, Case No. 87-511*, Los Angeles County, California.~~

~~Yorke, Callyn D. 1991. *Biological Resources Report, SEA No. 55, Parcel Map No. 13584-80 Ac.*, Los Angeles County, California.~~

~~Yorke, Callyn D., Ph.D. 1990. *Biological Resources Report on APN 3064-16-10, 22, Double G Ranch, SEA No. 55*, Los Angeles County, California.~~

Cruzan Mesa Vernal Pools SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded. Second Edition*. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~Boyd, S. 1999. *Vascular Flora of the Liebre Mountains, Western Transverse Ranges, California*. *Aliso* 18(2): 93-139~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The Cruzan Mesa Vernal Pools SEA Update Study*, Los Angeles County, California.~~

~~Fish and Wildlife Services. 2000. *Federal Register*. "Endangered and Threatened Wildlife and Plants; Proposed Determination of Critical Habitat for the Coastal California Gnatcatcher; Proposed Rule". Fish and Wildlife Services; 32 pp.~~

~~Frank Hovore & Associates. *Golden Valley Ranch Biological Resources Report*. 1999.~~

DRAFT

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~PCR Services Corporation. 2000. Biological Resources Assessment of the Proposed Cruzan Mesa Vernal Pools Significant Ecological Area. 19 pp.~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition*. Sacramento: California Native Plant Society Press.~~

~~Stephenson, J.R. and G.M. Calcarone. 1999. *Southern California Mountains and Foothills Assessment: Habitat and Species Conservation Issues*. General Technical Report GTR-PSW-172. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture; 402 pp.~~

~~U.S. Fish and Wildlife Service. 1997. *Vernal Pools of Southern California Draft Recovery Plan*. U.S. Fish and Wildlife Service, Portland, Oregon. 113+pp.~~

~~Zedler, P.H. 1987. *The Ecology of Southern California Vernal Pools: A Community Profile*. U.S. Fish and Wildlife Service Biological Report 85(7.11). 136 pp.~~

East San Gabriel Valley SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The East San Gabriel Valley SEA Description*, Los Angeles County, California.~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Los Angeles County, Department of Parks and Recreation. Date unknown. *Bonelli Regional Park Bird List*, San Dimas, California.~~

~~Michael Brandman Associates. 1995. *Draft Environmental Impact Report, Frank G. Bonelli Regional County Park Master Plan, State Clearinghouse No. 94061006*, Los Angeles, California.~~

~~Quinn, R.P. Date unknown. *The Status of Walnut Forests and Woodlands (Juglans californica) in Southern California*, Department of Biological Sciences, California State Polytechnic University, Pomona, California.~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition*. Sacramento: California Native Plant Society Press.~~

DRAFT

Griffith Park SEA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals* (898 taxa). 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 2006. *The Griffith Park SEA Description*, Los Angeles County, California.

City of Los Angeles, Department of Recreation and Parks. 2005. *Griffith Park Master Plan Draft/Redraft of Chapter Four, The Plant and Animal World*.

Cooper, Daniel S. and Paul Mathewson. 2008. *Griffith Park Wildlife Management Plan—Draft*. 88 pp.

http://www.laparks.org/dos/parks/griffithpk/wildlife/GP_WMP_Draft4.pdf.

Dale, N. 1986. *Flowering Plants: The Griffith Park, Coastal & Chaparral Regions of Southern California*.

Griffith Park Master Plan Working Group. December 2005. *Griffith Park Master Plan, Draft [Redraft of Chapter Four]*, City of Los Angeles, California.

Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition*. Sacramento: California Native Plant Society Press.

U.S. Department of the Interior, Fish and Wildlife Service. 1998. *Draft Recovery Plan for Six Plants from the Mountains Surrounding the Los Angeles Basin*, Region One, Portland, Oregon.

Harbor Lake Regional Park SEA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

Boyd, S. 1999. *Vascular Flora of the Liebre Mountains, Western Transverse Ranges, California*. *Aliso* 18(2):93-139

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals* (898 taxa). 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

DRAFT

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~GDM Environmental. December 2006. *Proposition O, Final Project Concept Report, Machado Lake Ecosystem Rehabilitation Project, Section 2.2.3*. City of Los Angeles, California.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The Harbor Regional Park SEA Description*, Los Angeles County, California.~~

~~Fish and Wildlife Services. 2000. *Federal Register*. "Endangered and Threatened Wildlife and Plants; Proposed Determination of Critical Habitat for the Coastal California Gnatcatcher; Proposed Rule". Fish and Wildlife Services; 32 pp.~~

~~Frank Hovore & Associates. *Golden Valley Ranch Biological Resources Report*. 1999.~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Sawyer, Keeler Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~Stephenson, J.R. and G.M. Calcarone. 1999. *Southern California Mountains and Foothills Assessment: Habitat and Species Conservation Issues*. General Technical Report GTR-PSW-172. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture; 402 pp.~~

~~U.S. Fish and Wildlife Service. 1997. *Vernal Pools of Southern California Draft Recovery Plan*. U.S. Fish and Wildlife Service, Portland, Oregon. 113+pp.~~

~~UtopianNature.com. 2005. Heindel, Mitch. <http://www.utopianature.com/kmhrp/home.html>~~

~~Last Accessed August 16, 2011.~~

~~Zedler, P.H. 1987. *The Ecology of Southern California Vernal Pools: A Community Profile*. U.S. Fish and Wildlife Service Biological Report 85(7.11). 136 pp.~~

Joshua Tree Woodlands SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The Joshua Tree Woodlands SEA*~~

DRAFT

~~Description, Los Angeles County, California.~~

~~EIP Associates.1990.Biota Report, SEATAC Project 88537, the Hughes Ranch, Quartz Hill, Los Angeles County, California.~~

~~England & Nelson Environmental Consultants.1976.Land Capability/Suitability Study, Los Angeles County General Plan Revision Program, Significant Ecological Areas Report, Riverside, California.~~

~~ESA (Environmental Science Associates).1999.Biological Constraints of Los Angeles Prayer Mountain, SEATAC Analysis Report, Los Angeles County, California.~~

~~Goode, S.1989.SEATAC Biota Report on #88502, 47811 Three Points Road, Los Angeles County, California.~~

~~Goode, S.1989.Revised SEATAC Biota Report on #88502, 47811 Three Points Road, Los Angeles County, California.~~

~~Goode, S.1989.Revised Impacts and Mitigations, SEATAC Biota Report on #88502, 47811 Three Points Road, Los Angeles County, California.~~

~~Holland RF.Preliminary descriptions of the terrestrial natural communities of California.1986 and 1992 Update.California Department of Fish and Wildlife unpublished report.~~

~~Independent Environmental Consultants.1994.Biological Constraints Analysis, the Amargosa Creek Improvement Project, Identification and Analysis of Impacts to SEA No.56, Los Angeles County, California.~~

~~Sierra Delta Corporation.1988.Biota Report for Zond Systems, Inc., Los Angeles Wind Electric Generating Station, Gorman, California, Conditional Use Permit 86453, Las Vegas, Nevada.~~

~~Sawyer, Keeler-Wolf and Evens.2009.A Manual of California Vegetation, Second Edition Sacramento:California Native Plant Society Press.~~

Madrona Marsh Preserve SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012)The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded.Second Edition.Jepson Herbarium, Berkeley, CA.1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database.January 2011.Special Animals (898 taxa).60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database.April 2011.Special Vascular Plants, Bryophytes, and Lichens List.Quarterly publication.71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database.January 2011.State and Federally Listed Endangered and Threatened Animals of California.13 pp.~~

~~City of Torrance, 2010, General Plan Update Draft EIR, 5.3.1 Environmental Setting.~~

~~County of Los Angeles, Department of Regional Planning.2006.The Madrona Marsh SEA Description, Los Angeles County, California.~~

DRAFT

~~Fish and Wildlife Services.2000.Federal Register."Endangered and Threatened Wildlife and Plants; Proposed Determination of Critical Habitat for the Coastal California Gnatcatcher; Proposed Rule".Fish and Wildlife Services; 32 pp.~~

~~Holland RF.Preliminary descriptions of the terrestrial natural communities of California.1986 and 1992 Update.California Department of Fish and Wildlife unpublished report.~~

~~Sawyer, Keeler-Wolf and Evens.2009.A Manual of California Vegetation, Second Edition Sacramento:California Native Plant Society Press.~~

~~U.S. Fish and Wildlife Service.1997.Vernal Pools of Southern California Draft Recovery Plan.U.S. Fish and Wildlife Service, Portland, Oregon.113+pp.~~

~~Zedler, P.H.1987.The Ecology of Southern California Vernal Pools:A Community Profile.U.S. Fish and Wildlife Service Biological Report 85(7.11).136 pp.~~

Palos Verdes Peninsula and Coastline SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012)The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded.Second Edition.Jepson Herbarium, Berkeley, CA.1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database.April 2011.Special Vascular Plants, Bryophytes, and Lichens List.Quarterly publication.71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database.January 2011.State and Federally Listed Endangered and Threatened Animals of California.13 pp.~~

~~County of Los Angeles, Department of Regional Planning.2006.The Palos Verdes SEA Description, Los Angeles County, California.~~

~~Holland RF.Preliminary descriptions of the terrestrial natural communities of California.1986 and 1992 Update.California Department of Fish and Wildlife unpublished report.~~

~~Sawyer, Keeler-Wolf and Evens.2009.A Manual of California Vegetation, Second Edition Sacramento:California Native Plant Society Press.~~

~~U.S. Department of Commerce.National Oceanic and Atmospheric Administration.National Marine Sanctuary Program.2008.Channel Islands National Marine Sanctuary Management Plan/Final Environmental Impact Statemant.Silver Spring, MD~~

~~URS.2004.Environmental Impact Report Draft, Rancho Palos Verdes Natural Communities Conservation Planning Subarea Plan (SCH# 2003071008).Prepared for the City of Rancho Palos Verdes~~

Puente Hills SEA Sources

~~Bahrami, M.M., A.M.Dove, E.M.Neaves and B.A.Roberts.1997.Puente Hills Corridor:Greenspace Connectivity for Wildlife and People, Department of Landscape Architecture, California State Polytechnic University, Pomona, California.~~

DRAFT

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The Puente Hills SEA Description*, Los Angeles County, California.~~

~~Haas, C. and K. Crooks. 1999. *Carnivore Abundance and Distribution Throughout the Puente/ Chino Hills*, The Mountains and Recreation Conservation Authority, Malibu, California~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Michael Brandman Associates. 1993. *Responses to SEATAC Comments and Recommendations on Vista de Lomas Project, Powder Canyon SEA No. 17*, Los Angeles, California.~~

~~Michael Brandman Associates. 1992. *Supplemental Biota Report for the Vista de Lomas Hacienda Heights Project Powder Canyon SEA No. 17, Tentative Tract 51153*, Los Angeles County, California.~~

~~Michael Brandman Associates. 1992. *Biota Report for the Vista de Lomas Hacienda Heights Projects, Tentative Tract 51153*, Los Angeles County, California.~~

~~Michael Brandman Associates. 1991. *Phase I Study for the Tonner Canyon/Chino Hills SEA No. 15*. Los Angeles County, California.~~

~~Michael Brandman Associates. 1991. *Biological Constraints Analysis for the Apple Creek Estates Project*, Los Angeles County, California.~~

~~Michael Brandman Associates. 1989. *Supplement to Biological Resources Assessment, Brea Canyon Tentative Tract No. 44478*, Los Angeles County, California.~~

~~Michael Brandman Associates. 1989. *Supplement to Biological Resources Assessment, Brea Canyon Tentative Tract No. 44478*, Los Angeles County, California.~~

~~Michael Brandman Associates. 1989. *Biological Resources Assessment, Brea Canyon Tentative Tract No. 44478*, Los Angeles County, California.~~

~~Michael Brandman Associates. 1989. *Biological Resources Assessment, A and S Development Site*, Los Angeles, California.~~

~~Michael L. Ahlering & Associates, Inc. 1988. *Biota Supplement, Tentative Tract 46485*, Diamond Bar, California.~~

DRAFT

~~Planning Center, The. 1990. *Biological Resources Assessment, Southern California Golf Association Member's Club at Firestone, Tentative Tract No. 48380, SEA No. 15 Tonner Canyon, Los Angeles, California.*~~

~~Planning Center, The. 1989. *Biological Resources Assessment, Southern California Golf Association Member's Club at Firestone, Tentative Tract No. 48380, SEA No. 15 Tonner Canyon, Los Angeles, California.*~~

~~Puente Hills Habitat Preservation Authority.~~

~~<http://www.habitatauthority.org/publications.shtml>~~

~~Quinn, Ronald D. 1998. *Status of Black Walnuts (Juglans californica) in Southern Puente Hills Site, Department of Biological Sciences, California State Polytechnic University, Pomona, California.*~~

~~Quinn, Ronald D. 1990. *Addendum to Supplement Biota Report, Tract No. 48632, SEA 44, Hacienda Heights, CA. Los Angeles, California. Department of Biological Sciences, Cal State Polytechnic University, Pomona.*~~

~~Quinn, Ronald D. 1990. *Biota Report, Tract No. 48632, SEA 44, Hacienda Heights, CA. Los Angeles County, California. Department of Biological Sciences, Cal State Polytechnic University, Pomona*~~

~~Quinn, Ronald D. 1990. *Supplement to Biota Report, Tract No. 48632, SEA 44, Hacienda Heights, CA. Los Angeles County, California. Department of Biological Sciences, Cal State Polytechnic University, Pomona.*~~

~~Quinn, Ronald D. Date unknown. *The Status of Walnut Forests and Woodlands (Juglans californica) in Southern California, Department of Biological Sciences, California State Polytechnic University, Pomona, California.*~~

~~Sawyer, Keeler Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~SWCA Environmental Consultants. 2012. *Evaluation of functional connectivity for medium to large-bodied carnivores and mule deer across Colima Road in the Puente Hills Preserve, City of Whittier, California. Prepared for City of Whittier & Puente Hills Habitat Preservation Authority, Whittier, CA. 31 pp.*~~

~~<http://www.habitatauthority.org/pdf/Colima%20Road%20Wildlife%20Connectivity%20Assessment%201-5-12.pdf>~~

Rio Hondo College Wildlife Sanctuary SEA Sources

~~Bahrami, M.M., A.M. Dove, E.M. Neaves and B.A. Roberts. 1997. *Rio Hondo Wildlife Sanctuary Corridor: Greenspace Connectivity for Wildlife and People*, Department of Landscape Architecture, California State Polytechnic University, Pomona, California.~~

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals*~~

DRAFT

~~(898 taxa).60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database.April 2011.*Special Vascular Plants, Bryophytes, and Lichens List*.Quarterly publication.71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database.January 2011.*State and Federally Listed Endangered and Threatened Animals of California*.13 pp.~~

~~County of Los Angeles, Department of Regional Planning.2006.*The Rio Hondo Wildlife Sanctuary SEA Update Study*,Los Angeles County, California.~~

~~Haas, C.and K.Crooks.1999.*Carnivore Abundance and Distribution Throughout the Puente/ Chino Hills*, The Mountains and Recreation Conservation Authority, Malibu, California~~

~~Holland RF.*Preliminary descriptions of the terrestrial natural communities of California*.1986 and 1992 Update.California Department of Fish and Wildlife unpublished report.~~

~~Michael Brandman Associates.1993.*Responses to SEATAC Comments and Recommendations on Vista de Lomas Project, Powder Canyon SEA No.17*,Los Angeles, California.~~

~~Michael Brandman Associates.1992.*Supplemental Biota Report for the Vista de Lomas Hacienda Heights Project Powder Canyon SEA No.17, Tentative Tract 51153*,Los Angeles County, California.~~

~~Michael Brandman Associates.1992.*Biota Report for the Vista de Lomas Hacienda Heights Projects, Tentative Tract 51153*,Los Angeles County, California.~~

~~Michael Brandman Associates.1991.*Phase I Study for the Tonner Canyon/Chino Hills SEA No.15*.Los Angeles County, California.~~

~~Michael Brandman Associates.1991.*Biological Constraints Analysis for the Apple Creek Estates Project*,Los Angeles County, California.~~

~~Michael Brandman Associates.1989.*Supplement to Biological Resources Assessment, Brea Canyon Tentative Tract No.44478*,Los Angeles County, California.~~

~~Michael Brandman Associates.1989.*Supplement to Biological Resources Assessment, Brea Canyon Tentative Tract No.44478*,Los Angeles County, California.~~

~~Michael Brandman Associates.1989.*Biological Resources Assessment, Brea Canyon Tentative Tract No.44478*,Los Angeles County, California.~~

~~Michael Brandman Associates.1989.*Biological Resources Assessment, A and S Development Site*,Los Angeles, California.~~

~~Michael L.Ahlering & Associates, Inc.1988.*Biota Supplement, Tentative Tract 46485*,Diamond Bar, California.~~

~~Planning Center, The.1990.*Biological Resources Assessment,Southern California Golf Association Member's Club at Firestone, Tentative Tract No.48380, SEA No.15 Tonner Canyon*,Los Angeles, California.~~

~~Planning Center, The.1989.*Biological Resources Assessment,Southern California Golf Association*~~

DRAFT

~~Member's Club at Firestone, Tentative Tract No.48380, SEA No.15 Tonner Canyon, Los Angeles, California.~~

~~Quinn, Ronald.D.1998.*Status of Black Walnuts (Juglans californica) in Southern Rio Hondo Wildlife Sanctuary Site*, Department of Biological Sciences, California State Polytechnic University, Pomona, California.~~

~~Quinn, Ronald D.1990.*Addendum to Supplement Biota Report, Tract No.48632, SEA 44*,Hacienda Heights, CA.Los Angeles, California.Department of Biological Sciences, Cal State Polytechnic University, Pomona.~~

~~Quinn, Ronald D.1990.*Biota Report, Tract No.48632, SEA 44*,Hacienda Heights, CA.Los Angeles County, California.Department of Biological Sciences, Cal State Polytechnic University, Pomona~~

~~Quinn, Ronald D.1990.*Supplement to Biota Report, Tract No.48632, SEA 44*,Hacienda Heights, CA.Los Angeles County, California.Department of Biological Sciences, Cal State Polytechnic University, Pomona.~~

~~Quinn, Ronald.D.Date unknown.*The Status of Walnut Forests and Woodlands (Juglans californica) in Southern California*, Department of Biological Sciences, California State Polytechnic University, Pomona, California.~~

~~Sawyer, Keeler-Wolf and Evens.2009.*A Manual of California Vegetation, Second Edition* Sacramento:California Native Plant Society Press.~~

~~URS.2006.*Draft Master Environmental Impact Report, Rio Hondo College Master Plan, Section 3.3.Biological Resources*, Los Angeles, CA.~~

San Andreas SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti(editors, 2012)*The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*.Second Edition.Jepson Herbarium, Berkeley, CA.1600 pp., illus.~~

~~Boyd, S.1999.*Vascular Flora of the Liebre Mountains, Wester Transverse Ranges, California*.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*.Quarterly publication. 71 pp.~~

~~County of Los Angeles, Department of Regional Planning.2006.*The San Andreas SEA Description*, Los Angeles County, California.~~

~~EIP Associates.1990.*Biota Report, SEATAC Project 88537, the Hughes Ranch, Quartz Hill, Los Angeles County, California*.~~

~~England & Nelson Environmental Consultants.1976.*Land Capability/Suitability Study, Los Angeles County General Plan Revision Program, Significant Ecological Areas Report*, Riverside, California.~~

~~Environmental Audit, Inc.November 1989.*Biota Report for Barrel Springs Ranch 63 Acre Tract*, Palmdale, California.~~

DRAFT

~~ESA (Environmental Science Associates).1999.*Biological Constraints of Los Angeles Prayer Mountain, SEATAC Analysis Report*, Los Angeles County, California.~~

~~Goode, S.1989.*SEATAC Biota Report on #88502, 47811 Three Points Road*, Los Angeles County, California.~~

~~Goode, S.1989.*Revised SEATAC Biota Report on #88502, 47811 Three Points Road*, Los Angeles County, California.~~

~~Goode, S.1989.*Revised Impacts and Mitigations, SEATAC Biota Report on #88502, 47811 Three Points Road*, Los Angeles County, California.~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Independent Environmental Consultants.1994.*Biological Constraints Analysis, the Amargosa Creek Improvement Project, Identification and Analysis of Impacts to SEA No. 56*, Los Angeles County, California.~~

~~Sierra Delta Corporation.1988.*Biota Report for Zond Systems, Inc., Los Angeles Wind Electric Generating Station, Gorman, California, Conditional Use Permit 86453*, Las Vegas, Nevada.~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~Vollmar Consulting, Berkeley, CA. 2004. *Centennial Specific Plan Project Site 2003/2004 Botanical Survey Report. Landscape Setting; Special Status Plant Surveys; Perennial Bunchgrass Community Study. 108 pp. 13 fig, 11 tables, 2 graphs, 3 appendices.*~~

San Dimas Canyon and San Antonio Wash SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012)*The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~Baskin, J.N. and T.R. Haglund, T.R. 1995. *Distribution and Abundance of Native Fishes, Southwestern Pond Turtles and Two-Striped Garter Snakes Below Morris Dam and in Brown's Gulch, San Gabriel River Canyon*. San Marino Environmental Associates, San Marino, California. 45 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The San Dimas-San Antonio Wash*~~

DRAFT

~~SEA Description, Los Angeles County, California.~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Planning Consultants Research. 1995. *Biological Assessment for the Big Dalton Dam Modification Project*, Santa Monica, California.~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~Stephenson, J.R. and G.M. Calcarone. 1999. *Southern California Mountains and Foothills Assessment: Habitat and Species Conservation Issues*. General Technical Report GTR-PSW-172. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture; 402 pp.~~

San Gabriel Canyon SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~Baskin, J.N. and T.R. Haglund. 1995. *Distribution and Abundance of Native Fishes, Southwestern Pond Turtles and Two-Striped Garter Snakes Below Morris Dam and in Brown's Gulch, San Gabriel River Canyon*. San Marino Environmental Associates, San Marino, California.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning. 2006. *San Gabriel Canyon SEA Description*, Los Angeles County, California.~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Planning Consultants Research. 1995. *Biological Assessment for the Big Dalton Dam Modification Project*, Santa Monica, California.~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~Stephenson, J.R. and G.M. Calcarone. 1999. *Southern California Mountains and Foothills Assessment: Habitat and Species Conservation Issues*. General Technical Report GTR-PSW-172. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture.~~

DRAFT

Santa Clara River SEA Sources

~~Aspen Environmental Group.1995.Draft EIS Pacific Pipeline Project:Biological Assessment, Vol. II.Prepared for the California Public Utilities Commission.~~

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012)The Jepson Manual:Vascular Plants of California, Thoroughly Revised and Expanded.Second Edition.Jepson Herbarium, Berkeley, CA.1600 pp., illus.~~

~~Bowland, J.L.1989.Biota Report, PM 20108; PM 20148, Project Numbers 88371 and 88416, SEA No. 23, Acton, California.~~

~~Boyd, S.1999.Vascular flora of the Liebre Mountains, western transverse ranges, California. Publ. #5, Rancho Santa Ana Botanic Garden, Claremont, CA, in Aliso, 18(2):93-139~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. Special Animals (898 taxa). 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. Special Vascular Plants, Bryophytes, and Lichens List.Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. State and Federally Listed Endangered and Threatened Animals of California. 13 pp.~~

~~California Native Plant Society.1994.Inventory of Rare and Endangered Vascular Plants of California.~~

~~Chambers Group.1994.EIR:Soledad Canyon Sand and Gravel Mining.~~

~~CNDDB (California Natural Diversity Database).2000.Multiple Records Search for the Val Verde, Newhall, Mint Canyon, Acton, and Aqua Dulce Quadrangles.~~

~~Goatsworth, S.1990.Biota Report for Tentative Parcel Map 21462, SEA No. 61, Los Angeles County, California.~~

~~Goatsworth, S.1989, resubmitted 1990.Biota Report for Tentative Parcel Map 20461, SEA No. 61, Los Angeles County, California.~~

~~Goatsworth, S.1989, resubmitted 1990.Biota Report for Tentative Parcel Map 20726, SEA No. 61, Los Angeles County, California.~~

~~County of Los Angeles, Department of Regional Planning.2006.The Santa Clara River SEA Update Study, Los Angeles County, California.~~

~~Dames and Moore & Sikand Engineering Associates.1991.Addendum to the SEATAC Report for a Portion of SEA No. 19, Tentative Map 44831, C.U.P. 86-491, Los Angeles County, California.~~

~~EAI (Environmental Audit Inc.).1990.Biota Report for Raphael 9.3 Acre Parcel, Los Angeles County, California.~~

~~England & Nelson Environmental Consultants.1976.Land Capability/Suitability Study, Los Angeles County General Plan Revision Program, Significant Ecological Areas Report, Riverside, California.~~

DRAFT

~~Fishman, R.H., Charlton, D. and R. Stafford.1990, revised August 1990.*Biota Survey and Impact Assessment for Parcel Map #20348, Angeles Forest Highway, Los Angeles County, California.*~~

~~Fishman, R.H., Charlton, D. and M. Hagan.1989, revised July 1989.*Biota Survey and Impact Assessment for Parcel Map #19258, Angeles Forest Highway, Los Angeles County, California.*~~

~~Frank Hovore & Associates.1999.*Biological Resources Report, Placerita Chaparral / Mitchell Development.*~~

~~Frank Hovore & Associates.1998.*Biological Resources Report, Golden Valley Ranch.*~~

~~Frank Hovore & Associates.1997.*Biological Constraints Assessment, Oak Springs Golf Course, TT 52004.*~~

~~Frank Hovore & Associates.1996.*Biotic Assessment and Report of Sensitive Resource Surveys, L.A. Department of Water and Power, Castaic Power Plant and Vicinity, L. A. County, California.*~~

~~Frank Hovore and Associates.1990.*SEATAC Biota Report, SEA No. 61, Kentucky Springs, Tentative Parcel Map No. 22107, Acton, California.*~~

~~Frank Hovore and Associates.1990.*SEATAC Biota Report, Supplemental Materials to Tierra Madre Consultants (TMC) Report on Tentative Parcel Maps 19628, 19899, 20056, 20057, San Francisquito Canyon, SEA No. 19.*~~

~~Frank Hovore and Associates.1989.*SEATAC Biotic Report, SEA No. 23 the Santa Clara River Buffer Zone, Tentative Tract 21273, Project No. 89315, Santa Clarita, California.*~~

~~Furgo McClelland (West), Inc.1992.*Biota Report for the Agua Dulce Quarry (Project No. 91307).*~~

~~Guthrie, D.1995.*Bird Surveys Along the Santa Clara River and its Tributaries Near Valencia, California.*~~

~~Guthrie, D.1993.*Bird Surveys Along the Santa Clara River and its Tributaries Near Valencia, California.*~~

~~Guthrie, D.1990.*Bird Surveys Along the Santa Clara River and its Tributaries Near Valencia, California.*~~

~~Guthrie, D.1988.*Bird Surveys Along the Santa Clara River and its Tributaries Near Valencia, California.*~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California.* 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Hovore, Frank T.*Checklist of the birds of Robinson Ranch golf course development.*2000.2 page leaflet.~~

~~ISI (Impact Sciences, Inc.).1995.*Biological Constraints Analysis, Tentative Parcel Map 19091 (West San Francisquito Creek), Los Angeles County, California.*~~

~~ISI (Impact Sciences, Inc.).1995.*Biota Report for Creekside Village Project, Tentative Tract Map 44831, Los Angeles County, California.*~~

DRAFT

~~ISI (Impact Sciences, Inc.).1995a.*Biological Constraints Analysis:Ranch Road South Project, Valencia, California, Tentative Tract Map 51931.*~~

~~ISI (Impact Sciences, Inc.).1995b.*Biota Report:West Creek Project.*~~

~~ISI (Impact Sciences, Inc.).1995c.*Biota Report:West Creek Project. Supplemental Information Submitted to SEATAC at Their Request on December 16, 1995.*Prepared for Los Angeles County Department of Regional Planning, Los Angeles, California.~~

~~ISI (Impact Sciences, Inc.) and RECON.1995.*Biota Report, Newhall Ranch Specific Plan, Tentative Tract Map 44831, Santa Clara River Valley, California.*~~

~~ISI (Impact Sciences, Inc.) and RECON.1994.*Biological Constraints Analysis:Newhall Ranch, California, Case No. 94087.*~~

~~Independent Environmental Consultants.1992.*Biological Constraints Analysis of Tentative Minor Land Division Map No. 23217, in Buffer Zone of SEA No. 23 Soledad Canyon, Santa Clara River, Los Angeles County, CA.*~~

~~Independent Environmental Consultants.1992.*SEATAC Report for Tentative Minor Land Division Map. No. 91253, Los Angeles, California.*~~

~~Independent Environmental Consultants.1992.*SEATAC Report for Tentative Minor Land Division Map. No. 91253 [Response to comments], Los Angeles, California.*~~

~~Independent Environmental Consultants.1990.*SEATAC Report for a Portion of SEA No. 19. Tentative Tract Map 44831. August 21.*Prepared for Valencia Company.~~

~~Independent Environmental Consultants and Sikand Engineering Associates.1988, revised 1989.*SEATAC Report for a Portion of SEA No. 23, Arbor Park, California.*~~

~~John M. Tettmer & Associates, Ltd.1994.*Response to U.S. Army Corps of Engineers Scoping Notice Comments for the U.S. Army Corps of Engineers, General Section 404 Permit, Santa Clara River from Castaic Creek to One Half Mile Above the Los Angeles Aqueduct and Portions of San Francisquito Creek and the Santa Clara River, South Fork, Los Angeles County, California.*~~

~~John M. Tettmer & Associates, Ltd.1993.*Biological Assessment:General Permit for the Santa Clara River from Castaic Creek to One Half Mile Above the Los Angeles Aqueduct, Portions of San Francisquito Creek, and the Santa Clara River, South Fork, Los Angeles County, California.*~~

~~John M. Tettmer & Associates, Ltd.1993, addendum submitted October 1993.*U.S. Army Corps of Engineers, General Section 404 Permit, Santa Clara River from Castaic Creek to One Half Mile Above the Los Angeles Aqueduct and Portions of San Francisquito Creek and the Santa Clara River, South Fork, Los Angeles County, California.*~~

~~Los Angeles County Museum of Natural History.1991.*Los Angeles County Breeding Bird Atlas, Santa Clarita Valley blocks.*California.~~

~~Mane'e Planning, Economic and Environmental Services.1989.*Ecological Impact Evaluation for Surface Mining Permit 86-357, Curtis Sand and Gravel (revised), Yucaipa, California.*~~

DRAFT

- ~~Marsh, K.G. 1990. *Biota Report, Los Angeles County Project No. 89189, for Surface Mining Within SEA No. 23 (Santa Clara River)*, Los Angeles County, California.~~
- ~~MBA (Michael Brandman & Associates). 1991. *Phase I Report: San Francisquito Canyon SEA No. 19*, Los Angeles County, California.~~
- ~~MBA (Michael Brandman & Associates). 1991. *Phase I Study for the Kentucky Springs SEA No. 61*, Los Angeles County, California.~~
- ~~MBA (Michael Brandman & Associates). 1992. *Biological Constraints for Clougherty Ranch Project*.~~
- ~~MBA (Michael Brandman & Associates). 1993 (revisions and supplemental info.). *Biota Report for the Tesoro del Valle Project*.~~
- ~~Mattoni, R. *Butterflies of greater Los Angeles*. wall poster, 2 sides. Rachel Tierney Consulting. 1996. *Supplemental Biota Report. Soledad Rock Quarry*.~~
- ~~Sawyer, Keeler Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press~~
- ~~Schoenherr, A. 1976. *The Herpetofauna of the San Gabriel Mountains, Los Angeles County, California, including distribution and biogeography*. Spec. Publ. Southwestern Herpetologists Society.~~
- ~~SEATAC *Biota Report, West Creek (VTTM 52455) and East Creek (VTTMs 44831, 52667) San Francisquito Canyon, Santa Clarita, Los Angeles County, CA*. 1998~~
- ~~SEATAC *Biota Report, North Valencia Annexation 2 Project (VTTM's 44831, 52667) San Francisquito Canyon, Santa Clarita, Los Angeles County, CA*. 1998~~
- ~~Stephenson, J.R. and G.M. Calcarone, 1999. *Southern California mountains and Foothills Assessment: Habitat and Species Conservation Issues*. Gen. Tech. Rep. GTR-PSW-172. Albany, CA. Pacific SW Res. Sta., Forest Serv., U.S. Department Agric., 402 pp.~~
- ~~Taylor & Company. 1995. *Biota Report: Los Angeles County Project No. 94 129, Soledad Rock Quarry, Surface Mining Permit, Conditional Use Permit*, Los Angeles County, California.~~
- ~~Tierra Madre Consultants. 1990. *Biota Report on Tentative Parcel Maps 19628, 19899, 20056, and 20057*.~~
- ~~Warren B. Houghton. 1986. *Biota Report on Parcel Map 17684, SEA No. 19*, Los Angeles County, California~~
- ~~Yorke, C. 1990. *Addendum to Biological Resources Report, Project No. 90113 / PM 21856, Kentucky Springs*, Los Angeles County, California.~~
- ~~Yorke, C. 1990. *SEATAC Biota Report, APN 3056-12-31, Kentucky Springs, SEA No. 61, Project No. 90113*, Los Angeles County, California.~~
- ~~Yorke, C. 1989. *SEATAC Biota Report, APN 3209-14-21, Santa Clara River Buffer Zone SEA 23*.~~
- ~~Yorke, C. 1989. *Biota Report for Frank Collins, C/O Archer Real Estate, SEA No. 23, Canyon Country, California*.~~

DRAFT

Santa Felicia SEA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 1993. *Additional Information and Analysis Regarding the Environmental Impact Report for the Sunshine Canyon Landfill Extension*, Los Angeles County, California.

County of Los Angeles, Department of Regional Planning. 2006. *The Santa Felicia SEA Description*, Los Angeles County, California.

Guthrie, Daniel A. and B. A. Prigge. 1989 (Jan. 31, 1990). *Emendations I & II of Biological Survey of Case 88551, Housing Development for Tract No. 47329, Unincorporated Area of Chatsworth*, Los Angeles County, California.

Guthrie, Daniel A. and B. A. Prigge. 1989 (revised Dec. 26, 1991). *Biological Survey of Case 88551, Housing Development for Tract No. 47329, Unincorporated Area of Chatsworth*, Los Angeles County, California.

Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.

Los Angeles Department of Regional Planning, 2010 Revised Draft Santa Clarita Valley Area Plan.

Michael Brandman Associates. 1995. *Biological Constraints Analysis for Tentative Parcel Map No. 23793*, Los Angeles County, California.

Michael Brandman Associates. 1989. *Supplemental to Biological Resources Assessment of Indian Wells Estates, Tentative Tract 44327*, Los Angeles County, California.

Michael Brandman Associates. 1988. *Biological Resources Assessment, Woolsey Canyon Project Site, Tentative Tract 45756*, Los Angeles County, California.

Penrod, Kristeen, Clint R. Cabañero, Paul Beier, Wayne Spencer, Claudia Luke, and Esther Rubin. 2005. *A linkage design for the Sierra Madre Castaic Connection*. South Coast Missing Linkages Project. www.scwildlands.org. x, 98 pp., Appendices A, B.

Regional Environmental Consultants (RECON) and Impact Sciences. 1995. *Biota Report, Newhall Ranch Specific Plan, Volume I of II, Santa Clara River Valley*, California, Tentative Tract Map 44831.

DRAFT

~~Regional Environmental Consultants (RECON) and Impact Sciences.1995.*Biota Report, Newhall Ranch Specific Plan, Volume II of II, Santa Clara River Valley*, California, Tentative Tract Map 44831.~~

~~Sawyer, Keeler Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition*. Sacramento: California Native Plant Society Press~~

~~Ultrasystems Engineers & Constructors, Inc.Environmental Services.1991.*Seatac Report.Tentative Tract No. 43896*, Los Angeles County, California.Dale Poe Development Corporation.~~

~~Ultrasystems Engineers & Constructors, Inc.Environmental Services.1991.*Seatac Report. Supplemental Information.Tentative Tract No. 43896*.Los Angeles County, California.Dale Poe Development Corporation.~~

~~Ultrasystems Engineers & Constructors, Inc.Environmental Services.1991.*Seatac Report. Supplemental Report No. 2.Tentative Tract No. 43896*.Los Angeles County, California.Dale Poe Development Corporation.~~

~~Ultrasystems Engineers & Constructors, Inc.Environmental Services.1990.*Biotic Report for Sunshine Mountain Park, Travel Trailer Park, Santa Clarita Valley*, California.~~

Santa Monica Mountains SEA Sources

~~Atlantis Scientific.1990.*Biological Impact Report, Project No. 90088/PM - 21998/CUP*, Beverly Hills, California.~~

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012)*The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*.Second Edition.Jepson Herbarium, Berkeley, CA.1600 pp., illus.~~

~~Biological Assessment Services.1992.*Biological Resources Assessment Vesting Tentative Tract No. 47927, SEA No. 12, Palo Comado Canyon*, Los Angeles County, California.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*.Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~C.W. Cook Co.Inc.1990.*SEATAC Information Packet for SEA No. 4, Project No. 86089 (Biology Reports)*, Los Angeles, California.~~

~~Coatsworth, S. and T. Laughlin.1989.*Biota Report fo Tentative Tract No. 48238, SEA No. 5*, Los Angeles County, California.~~

~~County of Los Angeles, Department of Regional Planning.2006.*The Santa Monica Mountains SEA Description*, Los Angeles County, California.~~

~~County of Los Angeles, Department of Regional Planning.1999.*The Santa Monica Mountains North Area Plan (Draft)*, Los Angeles County, California.~~

DRAFT

~~County of Los Angeles, Department of Regional Planning. 1996. *Environmental Impact Report Vol. 1 (Screencheck Draft)*, Project No. 89-251, SEA No. 12, Los Angeles County, California.~~

~~Dale, N. 1986. *Flowering Plants: The Santa Monica Mountains, Coastal & Chaparral Regions of Southern California*.~~

~~David Carroll & Associates. 1992. *Biological Constraints Analysis Report*, Project No. 91-359, SEA No. 3-Buffer, Latigo Canyon Road, Los Angeles County, California.~~

~~David Carroll & Associates. 1991. *Biota Report for Subdivision Parcel Map No. 89494 Latigo Canyon*, Los Angeles County, California.~~

~~David J. Tanner & Associates. 1988. *Biological Resources Assessment and Tree Inventory and Evaluation for Malibu Terrace, Tract No. 45342 and No. 45343*, SEA No. 12, Los Angeles County, California.~~

~~Edelman, Paul. 1990 (revised 2/91). *Critical Wildlife Corridor/Habitat Linkage Areas Between the Santa Susana Mountains, The Simi Hills and The Santa Monica Mountains*, Los Angeles County, California.~~

~~Envicom Corporation. 1999. *Biota Report (Resubmittal No. 2)*, Beautiful Homes Site, Los Angeles County, California.~~

~~Envicom Corporation. 1992. *Biological Constraints Analysis, Tentative Parcel Map 22352*, SEA No. 3, Los Angeles County, California.~~

~~Envicom Corporation. 1992. *Soka University Expansion Project*, Los Angeles County, California.~~

~~Envicom Corporation. 1990. *Biota Report for Lots 159 and 160 of Tentative Tract No. 32268*, Calabasas, California.~~

~~Envicom Corporation. 1990. *Biota Report for Old West Valley Driving Range (2nd Submittal)*, Calabasas, California.~~

~~Envicom Corporation. 1990. *Biota Report for Old West Valley Driving Range*, Calabasas, California.~~

~~Envicom Corporation. 1990. *Supplemental Oak Tree Report for Old West Valley Driving Range*, SEA No. 12, Calabasas, California.~~

~~Envicom Corporation. 1989. *Oak Tree Report for Lots 159 and 160 of Tentative Tract No. 32268 for Bulk Grading Disposal for Adjacent Tract No. 37824*, Calabasas, California.~~

~~Envicom Corporation. 1989. *Biota Report, SEA No. 3-Buffer, Mulholland Highway Residential Development (4th Submittal)*, Los Angeles County, California.~~

~~Envicom Corporation. 1989. *Biota Report (resubmittal) for Conditional Use Permit Case No. 87361*, SEA No. 5, Los Angeles County, California.~~

~~Envicom Corporation. 1989. *Biota Report for Conditional Use Permit Case No. 87361*, SEA No. 5, Los Angeles County, California.~~

~~Environmental Audit, Inc. 1990. *Biota Report for Pepperdine University LRDP Developments, Including CUP Application No. 90-218*, Malibu, California.~~

DRAFT

- ~~Environmental Audit, Inc.1990.*Biota Report for Raphael 9.3 Acre Parcel (Tentative Parcel Map No. 20549) at 31905 Pacific Coast Highway, Malibu, California.*~~
- ~~Hartzell, A.K. and H. Hewitt.1992.Fish and Wildlife's Control Over Land Use.*California Planner* 4(8): 1-9.~~
- ~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California.* 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~
- ~~Impact Sciences, Inc.1988.*Biota Report for Calabasas Park West Project, SEA No. 12, Los Angeles County, California.*~~
- ~~Impact Sciences, Inc.1988.*Biota Report for Calabasas Road Expansion, SEA No. 12, (Amended June 1988), Los Angeles County, California.*~~
- ~~Independent Environmental Consultants.1993.*SEATAC Report for Tentative Minor Land Division Map No. 22648, Case No. 91253, SEA No. 3 Buffer, 32356 Mulholland Highway, in Unincorporated Territory of the Santa Monica Mountains, Los Angeles County, California.*~~
- ~~L. Newman Design Group, Inc.1995.*Oak Tree Report, Vesting Tentative Tract No. 47927, SEA No. 12, Los Angeles County, California.*~~
- ~~Lee Newman & Associates, Inc.1988.*Tree Report: The Salvation Army Camp (Mt. Crags Camp and Camp Gilmore), Los Angeles County, California.*~~
- ~~Lieberstein, T.A.1989.*Reserve Design in the Santa Monica Mountains.*Thesis for M.A. degree, Cal State Univ. Northridge.~~
- ~~Land and Water Company, The.1992.*Biological Constraints Analysis Supplemental Report, Potential Impacts of Development and Mitigation, Project No. 91-359, Los Angeles County, California.*~~
- ~~McAuley, M.1996 (2nd ed.).*Wildflowers of the Santa Monica Mountains.*Canyon Publishing:Canoga Park, CA, USA.~~
- ~~McClelland Consultants (West), Inc.1991.*Biota Report for Tentative Tract Map No. 21221 (3rd submittal), Los Angeles County California.*~~
- ~~Michael Brandman Associates.1991.*Phase I Study for the Cold Creek SEA No. 9, Los Angeles County, California.*~~
- ~~Michael Brandman Associates.1991.*Phase I Study for the Las Virgenes SEA No. 6, Los Angeles County, California.*~~
- ~~Michael Brandman Associates.1991.*Phase I Study for the Tuna Canyon SEA No. 10, Los Angeles County, California.*~~
- ~~Michael Brandman Associates.1991.*Biological Resources Assessment for the Liberty Canyon Project (Resubmittal), Project No. 90152, SEA No. 12, Los Angeles County, California.*~~
- ~~Michael Brandman Associates.1991.*Biological Resources Assessment for the Liberty Canyon Project, Project No. 90152, SEA No. 12, Los Angeles County, California.*~~

DRAFT

~~Michael Brandman Associates. 1990. *Biological Resources Assessment for the Cordillera Reclaimed Water Tank and Ancillary Facilities, Las Virgenes Municipal Water District, SEA No. 12, Los Angeles County, California.*~~

~~Michael Brandman Associates. 1989. *Biological Resources Assessment for the _____ Seminole Water Tank No. 2, Los Virgenes Municipal Water District, SEA No. 3 Buffer, Los Angeles County, California.*~~

~~Michael Brandman Associates. 1989. *Biological Resources Assessment for Malibu Terrace Project Site, Project No. 87527, SEA No. 12, Los Angeles County, California.*~~

~~Nelson, Erik. 2000. "Cameras Capture Critter Commuters", *Daily News*, Monday May 22, 2000.~~

~~P&D Environmental Services. 1995. *Biological Resources Assessment for Tentative Tract No. 47927, SEA No. 12, Palo Comado Canyon, Los Angeles County, California.*~~

~~P&D Environmental Services. 1995. *Biological Resources Assessment Vesting Tentative Tract No. 47927, SEA No. 12, Palo Comado Canyon, Los Angeles County, California.*~~

~~Planning Center, The. 1991. *Biological Resources Assessment for Tentative Tract No. 21887, SEA No. 3, Los Angeles County, California.*~~

~~Planning Center, The. 1990. *Biological Resources Assessment for Tentative Tract No. 47927, SEA No. 12 Palo Comado Canyon, Los Angeles County, California.*~~

~~Raven, Peter H., Henry J. Thompson, and Barry A. Prigge. 1986. *Flora of the Santa Monica Mountains, California, 2nd edition. Southern California Botanists Special Publication No. 2.*~~

~~Rindlaub, K. 1989. *Biological Resources Evaluation, Julian Property, Loma Metisse and Rotunde Mesa Road, Santa Monica Mountains, SEA No. 5, Los Angeles County, California.*~~

~~Rindlaub, K. 1989. *Biological Resources Evaluation, Julian Property, Loma Metisse and Rotunde Mesa Road, Santa Monica Mountains, Los Angeles County, California.*~~

~~Ryan Ecological Consulting. 2010. *The Western Snowy Plover in Los Angeles County, California. Report to California Department of Fish and Wildlife. 136 pp.*~~

~~<http://losangelesaudubon.org/images/stories/pdf/westernsnowyploverin%20losangelescounty.pdf>~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~U.S. Department of the Interior, Fish and Wildlife Service. 1998. *Draft Recovery Plan for Six Plants from the Mountains Surrounding the Los Angeles Basin, Region One, Portland, Oregon.*~~

~~U.S. Department of the Interior, National Park Service. 1998. *Santa Monica Mountains National Recreation Area Land Protection Plan, Los Angeles County, California.*~~

~~USFWS. 2000. *Draft Recovery Plan for the red-legged Frog (Rana aurora draytonii)*. U.S. Fish and Wildlife Service, Portland, Oregon. 258 pp.~~

~~Vincent N. Scheidt, M.A., Biological Consultant. 1991. *Biota Report for the Vierich Lot Split Project,*~~

DRAFT

~~Parcel Map No. 21737; File No. 90259, Los Angeles County, California.~~

~~Yorke, Callyn D., Ph.D.1993.*Response to SEATAC Comments of February 1, 1993, 50 Acres, 2700 Corral Canyon Road, Malibu, Project No. 90-504/TR 50290*, Los Angeles County, California.~~

~~Yorke, Callyn D., Ph.D.1992.*Biological Resources Report on 50 Acres, 2700 Corral Canyon Road, Malibu, Project No. 90504*, Los Angeles County, California.~~

~~Yorke, Callyn D., Ph.D.1992.*Biological Resources Report on 10.8 Acres, SEA No. 5 Buffer, Malibu (Project No. 88217)*, Los Angeles County, California.~~

~~Yorke, Callyn D., Ph.D.1992.*Biological Resources Report on 50 Acres, 2700 Corral Canyon Road, Project No. 90504*, Los Angeles County, California.~~

~~Yorke, Callyn D., Ph.D.1991.*Biological Resources Report on 10.8 Acres, SEA No. 5 Buffer, Malibu (Project No. 88217)*, Los Angeles, California.~~

~~Yorke, Callyn D., Ph.D.1991.*Biological Resources Report on 50 Acres, 2700 Corral Canyon Road, Project No. 90504*, Los Angeles County, California.~~

Santa Susana Mountains and Simi Hills SEA Sources

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012)*The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~County of Los Angeles, Department of Regional Planning.1993.*Additional Information and Analysis Regarding the Environmental Impact Report for the Sunshine Canyon Landfill Extension*, Los Angeles County, California.~~

~~County of Los Angeles, Department of Regional Planning.2006.*The Santa Susana Mountains-Simi Hills SEA Description*, Los Angeles County, California.~~

~~Envicom Corporation.1992 (revised Feb. 4, 1993).*A Consideration of Wildlife Movements in the Santa Susana Mountains*, Los Angeles County, California.~~

~~Guthrie, Daniel A. and B. A. Prigge.1989 (Jan. 31, 1990).*Emendations I & II of Biological Survey of Case 88551, Housing Development for Tract No. 47329, Unincorporated Area of Chatsworth*, Los Angeles County, California.~~

~~Guthrie, Daniel A. and B. A. Prigge.1989 (revised Dec. 26, 1991).*Biological Survey of Case 88551, Housing Development for Tract No. 47329, Unincorporated Area of Chatsworth*, Los Angeles County, California.~~

DRAFT

~~Guthrie, Daniel A. and B. A. Prigge. 1988. *Biological Survey of Case 88312, Christara Motion Picture Studio Site, 23747, The Old Road, Unincorporated Area of Newhall, Los Angeles County, California*~~

~~Henrickson, James, PhD. 1996. *Biological Constraints Analysis Zone Change—Conditional Use Permit 96-113, 22925 Coltrane Avenue, Within SEA No. 20, Santa Susana Mountains, Los Angeles County, California.*~~

~~Henrickson, James, PhD., Independent Environmental Consultants. 1992. *Biological Constraints Analysis of Tentative Minor Land Division Map No. 23217, in Buffer Zone of SEA No. 23, Soledad Canyon, Santa Clara River, Los Angeles County, California.*~~

~~Holland RF. *Preliminary descriptions of the terrestrial natural communities of California.* 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.~~

~~Impact Sciences, Inc. 1996 (Supplemental Info. Submitted Nov. 2, 1999). *Biota Report, Dahl Property, SEA No. 21, Tentative Parcel Map 23793, Chatsworth, California.*~~

~~Impact Sciences, Inc. and RECON. 1994. *Biological Constraints Analysis, Newhall Ranch, California, Case No. 94087.*~~

~~Michael Brandman Associates. 1995. *Biological Constraints Analysis for Tentative Parcel Map No. 23793, Los Angeles County, California.*~~

~~Michael Brandman Associates. 1989. *Supplemental to Biological Resources Assessment of Indian Wells Estates, Tentative Tract 44327, Los Angeles County, California.*~~

~~Michael Brandman Associates. 1988. *Biological Resources Assessment, Woolsey Canyon Project Site, Tentative Tract 45756, Los Angeles County, California.*~~

~~Regional Environmental Consultants (RECON) and Impact Sciences. 1995. *Biota Report, Newhall Ranch Specific Plan, Volume I of II, Santa Clara River Valley, California, Tentative Tract Map 44831.*~~

~~Regional Environmental Consultants (RECON) and Impact Sciences. 1995. *Biota Report, Newhall Ranch Specific Plan, Volume II of II, Santa Clara River Valley, California, Tentative Tract Map 44831.*~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press~~

~~Ultrasystems Engineers & Constructors, Inc. Environmental Services. 1991. *Seatac Report. Tentative Tract No. 43896, Los Angeles County, California. Dale Poe Development Corporation.*~~

~~Ultrasystems Engineers & Constructors, Inc. Environmental Services. 1991. *Seatac Report. Supplemental Information. Tentative Tract No. 43896. Los Angeles County, California. Dale Poe Development Corporation.*~~

~~Ultrasystems Engineers & Constructors, Inc. Environmental Services. 1991. *Seatac Report. Supplemental Report No. 2. Tentative Tract No. 43896. Los Angeles County, California. Dale Poe Development Corporation.*~~

~~Ultrasystems Engineers & Constructors, Inc. Environmental Services. 1990. *Biotic Report for Sunshine Mountain Park, Travel Trailer Park, Santa Clarita Valley, California.*~~

Tujunga Valley and Hansen Dam SEA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 2006. *The Tujunga Valley/ Hansen Dam SEA Description*, Los Angeles County, California.

Hartzell, A.K. and H. Hewitt. 1992. Fish and Wildlife's Control Over Land Use. *California Planner* 4(8):1-9.

Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.

Sawyer, Keeler Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.

URS Environmental Services. 2006. *Big Tujunga Dam Seismic Upgrade, Initial Study and Mitigated Negative Declaration*, Los Angeles County

U.S. Department of the Interior, Fish and Wildlife Service. 1998. *Draft Recovery Plan for Six Plants from the Mountains Surrounding the Los Angeles Basin*, Region One, Portland, Oregon.

U.S. Department of the Interior, National Park Service. 1998. *Santa Monica Mountains National Recreation Area Land Protection Plan*, Los Angeles County, California.

Valley Oaks Savannah SEA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 2006. *The Santa Monica Mountains SEA*

DRAFT

~~Update Study, Los Angeles County, California.~~

~~County of Los Angeles, Department of Regional Planning. 1996. *Environmental Impact Report Vol. 1 (Screencheck Draft)*, Project No. 89-251, SEA No. 12, Los Angeles County, California.~~

~~Dale, N. 1986. *Flowering Plants: The Santa Monica Mountains, Coastal & Chaparral Regions of Southern California*.~~

~~Edelman, Paul. 1990 (revised 2/91). *Critical Wildlife Corridor/Habitat Linkage Areas Between the Santa Susana Mountains, The Simi Hills and The Santa Monica Mountains*, Los Angeles County, California.~~

~~Los Angeles Department of Regional Planning, 2010 Revised Draft Santa Clarita Valley Area Plan.~~

~~McAuley, M. 1996 (2nd ed.). *Wildflowers of the Santa Monica Mountains*. Canyon Publishing: Canoga Park, CA, USA.~~

~~Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~U.S. Department of the Interior, National Park Service. 1998. *Santa Monica Mountains National Recreation Area Land Protection Plan*, Los Angeles County, California.~~

Verdugo Mountains SEA Sources

~~Atlantis Scientific. 1990. *Biological Impact Report, Project No. 90088/PM - 21998/CUP*, Beverly Hills, California.~~

~~Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.~~

~~Biological Assessment Services. 1992. *Biological Resources Assessment Vesting Tentative Tract No. 47927, SEA No. 12, Palo Comado Canyon*, Los Angeles County, California.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals* (898 taxa). 60 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.~~

~~California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.~~

~~C.W. Cook Co. Inc. 1990. *SEATAC Information Packet for SEA No. 4, Project No. 86089* (Biology Reports), Los Angeles, California.~~

~~Christopher A. Joseph and Associates. 2004. *FINAL ENVIRONMENTAL IMPACT REPORT FOR THE CANYON HILLS PROJECT*, Prepared for the City of Los Angeles~~

~~County of Los Angeles, Department of Regional Planning. 2006. *The Verdugo Mountains SEA Description*, Los Angeles County, California.~~

DRAFT

~~Dale, N. 1986. Flowering Plants: The Verdugo Mountains, Coastal & Chaparral Regions of Southern California.~~

~~David Carroll & Associates. 1992. *Biological Constraints Analysis Report, Project No. 91-359, SEA No. 3 Buffer, Latigo Canyon Road*, Los Angeles County, California.~~

~~David Carroll & Associates. 1991. *Biota Report for Subdivision Parcel Map No. 89494 Latigo Canyon*, Los Angeles County, California.~~

~~David J. Tanner & Associates. 1988. *Biological Resources Assessment and Tree Inventory and Evaluation for Malibu Terrace, Tract No. 45342 and No. 45343, SEA No. 12*, Los Angeles County, California.~~

~~Edelman, Paul. 1990 (revised 2/91). *Critical Wildlife Corridor/Habitat Linkage Areas Between the Santa Susana Mountains, The Simi Hills and The Verdugo Mountains*, Los Angeles County, California.~~

~~Sawyer, Keeler Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.~~

~~U.S. Department of the Interior, Fish and Wildlife Service. 1998. *Draft Recovery Plan for Six Plants from the Mountains Surrounding the Los Angeles Basin*, Region One, Portland, Oregon.~~

Existing SEAs descriptions have been updated as necessary with new sources from the 2025 SEA Study. To view the sources cited for the SEAs adopted in 2015, visit: <https://planning.lacounty.gov/planning-library/historical-planning-documents/>

Species & Natural Communities (see also “Regional and Local Studies”, below)

Allen, L.W., Carmona-Freeman, M., Cooper, D.S., Feenstra, J., Garrett, K.L., George, G., Loquvam, M., Osgood, E., Ryan, T., San Miguel, M., & Vigallon, S. (2009). Los Angeles County's Sensitive Bird Species. *Western Tanager* 75(3).

Allen, L.W., Garrett, K.L., & Wimer, M.C. (2016). *Los Angeles County Breeding Bird Atlas*. Los Angeles Audubon Society, Los Angeles.

Bell, E., Dagit, R. & Ligon, F. (2011). Colonization and persistence of a southern California steelhead (*Oncorhynchus mykiss*) population. *Bulletin of the Southern California Academy of Sciences*, 110, 1 – 16.

Benson, J.F., Mahoney, P.J., Vickers, T.W., Sikich, J.A., Beier, P., Riley, S.P., Ernest, H.B., & Boyce, W.M. (2019). Extinction vortex dynamics of top predators isolated by urbanization. *Ecological Applications*, 29(3), e01868. <https://doi.org/10.1002/eap.1868>

Benson, J.F., Abernathy, H.N., Sikich, J.A., & Riley, S.P.D. (2021). Mountain lions reduce movement, increase efficiency during the Covid-19 shutdown. *Ecological Solutions and Evidence*, 2, e12093. <https://doi.org/10.1002/2688-8319.12093>

Blood, B.R. (2021). An annotated checklist of the terrestrial mammals of Los Angeles County Part I: *Didelphimorphia, Lagomorpha, and Rodentia*. *Bulletin of the Southern California Academy of Sciences*, 120, 99 – 127.

Blood, B.R. (2022). *An annotated checklist of the terrestrial mammals of Los Angeles County Part II:*

DRAFT

- Eulipotyphla, Carnivora, Artiodactyla, and Chiroptera. Bulletin of the Southern California Academy of Sciences*, 121, 41 – 87.
- Bloom, P.H., Barton, R.G., & Kuehn, M.J. (2023). Swainson's hawk nesting population in the Antelope Valley of the western Mojave Desert, California. *Western Birds*, 54, 32 – 43. DOI 10.21199/WB54.1.3. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/487/459>
- Bolger, D.T. (2007). Spatial and temporal variation in the Argentine ant edge effect: Implications for the mechanism of edge limitations. *Biological Conservation*, 136, 295 – 305. <https://doi.org/10.1016/j.biocon.2006.12.002>
- Boyd, S. & Mistretta, O. (2006). *Lepechinia rossii (Lamiaceae)*, a narrow endemic from the western transverse ranges of southern California. *Madroño*, 53(1), 77-84. [https://doi.org/10.3120/0024-9637\(2006\)53\[77:LRLANE\]2.0.CO;2](https://doi.org/10.3120/0024-9637(2006)53[77:LRLANE]2.0.CO;2)
- Brattstrom, B.R. (2013). Distribution of the coast horned lizard, *Phrynosoma coronatum [blainvillii]*, in southern California. *Bulletin of the Southern California Academy of Sciences*, 112, 206 – 216.
- California Department of Fish and Wildlife (CDFW). (2025a). *California Natural Communities*. List dated February 27, 2025. Vegetation Classification and Mapping Program, California Department of Fish and Wildlife. Sacramento, CA.
- CDFW. (2025b). *California Sensitive Natural Communities*. List dated February 27, 2025. Vegetation Classification and Mapping Program, California Department of Fish and Wildlife. Sacramento, CA.
- California Native Plant Society, Rare Plant Program. (2025). *Rare Plant Inventory* (online edition, v9.5.1). Website <https://www.rareplants.cnps.org>
- California Natural Diversity Data Base (CNDDDB). (2025a). *Special Vascular Plants, Bryophytes, and Lichens List*. List dated July 2025. California Department of Fish and Wildlife. Sacramento, CA.
- CNDDDB. (2025b). *Special Animals List*. List dated July 2025. California Department of Fish and Wildlife. Sacramento, CA.
- CNDDDB. (2025c). *State and Federally Listed Endangered and Threatened Plants of California*. List dated July 2025. California Department of Fish and Wildlife. Sacramento, CA.
- CNDDDB. (2025d). *State and Federally Listed Endangered and Threatened Animals of California*. List dated July 2025. California Department of Fish and Wildlife. Sacramento, CA.
- CNDDDB. (2025e). *CNDDDB Quickview tool in BIOS 6*. Bios6 version 6.25.0416 Built on ArcGIS Maps SDK for JavaScript 4.32.
- Charlton, D., & Rundel, P. (2017). The vegetation and flora of Edwards Air Force Base, western Mojave Desert, California. *Aliso*, 35(2), 51 – 68. <https://doi.org/10.5642/aliso.20173502.02>
- Community Conservancy International/Natural History Museum of Los Angeles County Foundation, 2001. *The Biota of the Baldwin Hills: An Ecological Assessment*. K.C. Molina, ed., February

DRAFT

2001.

- Cooper, D.S. (2000). Breeding landbirds of a highly threatened open space: The Puente-Chino Hills, California. *Western Birds*, 31, 213-234. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/1491/1428>
- Cooper, D.S. (2006a). Shorebird use of the lower Los Angeles River channel: A novel wetland habitat. *Western Birds*, 37, 1 – 7. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/1377/1319>
- Cooper, D.S. (2006b). Annotated checklist of extirpated, reestablished, and newly-colonized avian taxa of the Ballona Valley, Los Angeles County, California. *Bulletin of the Southern California Academy of Sciences*, 105(3). <http://scholar.oxy.edu/scas/vol105/iss3/1>
- Cooper, D.S. (2009). *Ecological assessment of open space remnants in northeastern Los Angeles Mountains Recreation and Conservation Authority, Los Angeles, California. November 15, 2008, updated April 20, 2009.*
- Cooper, D.S., Hamilton, R.A. & Lucas, S. (2012). A population census of the Cactus wren in coastal Los Angeles County. *Western Birds*, 43, 151 – 163. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/1687/1599>
- Cooper, D.S., Mongolo, J., & Dellith, C. (2017). Status of the California gnatcatcher at the northern edge of its range. *Western Birds*, 48, 124 – 140. DOI: 10.21199/WB48.2.3. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/1143/1097>
- Cooper, D.S. & Muchlinski, A.E. (2015). Recent decline of lowland populations of the western gray squirrel in the Los Angeles area of southern California. *Bulletin of the Southern California Academy of Sciences*, 114(1), 42-53. DOI: 10.3160/0038-3872-114.1.42.
- Couffer, M.C. (2019). The current northernmost breeding pair of the California Gnatcatcher. *Western Birds*, 50, 188 – 193. DOI 10.21199/WB50.3.8. [https://archive.westernfieldornithologists.org/archive/V50/50\(3\)-p188-p193.pdf](https://archive.westernfieldornithologists.org/archive/V50/50(3)-p188-p193.pdf)
- Ernest, H.B., Vickers, T.W., Morrison, S.A., Buchalski, M.R., & Boyce, W.M. (2014). Fractured genetic connectivity threatens a southern California puma (*Puma concolor*) population. *PLoS One*, 9(10), e107985. <https://doi.org/10.1371/journal.pone.0107985>
- Fraga, N. (2015). *Diversity, endemism and conservation of California monkeyflowers (Phrymaceae): a case study in Erythranthe section Paradantha*. PhD Dissertation. Claremont Graduate University.
- Grinnell, J. (1909). *A bibliography of California ornithology*. Pacific Coast Avifauna No 5. Cooper Ornithological Club of California. Santa Clara, CA.
- Hamilton, R.A. & Cooper, D.S. (2010). *Conservation & Management Plan for Marina del Rey, Los Angeles Co., California, September 16, 2010*. Prepared for: County of Los Angeles, Dept. of Beaches and Harbors, Marina del Rey, CA. https://file.lacounty.gov/SDSInter/dbh/docs/150561_FinalMDRConservationMgmtPlan_9-16-10.pdf
- Hamilton Biological, Inc. (2024). *Regional Colonial Waterbird Survey Report, Los Angeles County*.

DRAFT

- California. Prepared for Los Angeles County Department of Beaches and Harbors. November 15, 2024. https://file.lacounty.gov/SDSInter/dbh/docs/1170634_HamiltonBio_FinalLACountyWaterbirdSurveyRpt_11-15-24_lowres_1.pdf
- Hanes, T. L.; Friesen, R. D.; K. Keane. 1989. Alluvial Scrub Vegetation in Coastal Southern California. In: Abell, Dana L., Technical Coordinator. 1989. Proceedings of the California Riparian Systems Conference: protection, management, and restoration for the 1990s; 1988 September 22-24; Davis, CA. Gen. Tech. Rep. PSW-GTR-110. Berkeley, CA: Pacific Southwest Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture; p. 187-193.
- Holl, S.A. & Bleich, V.C. (2010). *Biennial Symposium of the Northern Wild Sheep and Goat Council*, 17, 139 – 157.
- Longcore, T. Ed. (2016). *Urban Biodiversity Assessment: Baldwin Hills Biota Update*. Los Angeles: University of Southern California for Baldwin Hills Conservancy (Proposition 84) and Baldwin Hills Regional Conservation Authority (Proposition A). <https://baldwinhillsnature.bhc.ca.gov/wp-content/uploads/2016/12/baldwinhillsbiotaupdate.pdf>
- Los Angeles County Sensitive Bird Species Working Group. (2009). Los Angeles County's sensitive bird species. *Western Tanager* 75(3), 1 – 8. Los Angeles Audubon Society.
- Morse, K. (2022). *Systematics and Conservation of the Genus Malacothamnus (Malvaceae)*. PhD Dissertation. Claremont Graduate University.
- PCR Project Team. (2006). *Sensitive Species Occurring or Potentially Occurring within the Los Angeles SEA. Biological Resource Assessment Los Angeles County SEA*. Prepared by "PCR Project Team", January 2006.
- Provance, M.C., & Sanders, A.C. (2013). Lucky Morning Glory, *Calystegia felix (Convolvulaceae)*: A new species from southern California, with notes on the historical ecology of the Chino ciénega belt. *PhytoKeys* 32 (December 2013), 1 – 26. <https://doi.org/10.3897/phytokeys.32.6020>.
- Remington, S., & Cooper, D.S. (2015). Bat survey of Griffith Park, Los Angeles, California. *Southwestern Naturalist* 59(4), 471-477. DOI: 10.1894/SGM-32.1
- Simpson, M.G., Simpson, L.M., & André, J.M. (2024). *Cryptantha acrimuricata (Boraginaceae)*, a distinctive new taxon of series *Muricatae*." *PhytoKeys* 250 (December 2024), 193 – 213. <https://doi.org/10.3897/phytokeys.250.138635>.
- Sharifi, M.R., Brostoff, W.N., & Rundel, P.R. (2017). Community structure and demography in a saline pan-dune mosaic in the western Mojave Desert. *Aliso* 35(2), 79 – 86. <https://doi.org/10.5642/aliso.20173502.04>.
- Read, E.A. (2015). A flora of the Ballona Wetlands and environs. *Bulletin of the Southern California Academy of Sciences*, 114, 149 – 163.
- Remington, S. (2016). Bat surveys of the Baldwin Hills, Los Angeles County, California, 2014 – 2015. (pp. 72 – 101) In T. Longcore, Ed. (2016). *Urban Biodiversity Assessment: Baldwin Hills Biota*

DRAFT

- Update. Los Angeles: University of Southern California for Baldwin Hills Conservancy (Proposition 84) and Baldwin Hills Regional Conservation Authority (Proposition A).
- Ryan, T.P., Vigallon, S., Griswold, R., Egger, C., Sheakley, S., Nguyen, L., & Schallmann, R. (2023). Western snowy plover in coastal Los Angeles and Orange Counties, California. *Western Birds*, 54, 44 – 64. DOI 10.21199/WB54.1.4. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/488/460>
- Shuford, W.D., Molina, K.C., Kelly, J.P., Condeso, T.E., Cooper, D.S., & Jongsomjit, D. (2020). Breeding status of the double-crested cormorant in the interior of California, 2009-2012. *Western Birds*, 51, 270-292. DOI: 10.21199/WB51.4.1 <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/773/733>
- Shuford, W.D., Kelly, J.P., Condeso, T.E., Cooper, D.S., Molina, K.C., & Jongsomjit, D. (2020). Distribution and abundance of colonial-nesting herons, egrets, and night-herons in California, 2009 – 2012. *Western Birds*, 51, 190-220. DOI: 10.21199/WB51.3.2 <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/768/728>
- Ryan, T.P., Vigallon, S., Cooper, D.S., Dellith, C., Johnston, K., & Nguyen, L. (2019). Return of beach-nesting Snowy Plovers to Los Angeles County following 68-year absence. *Western Birds*, 50, 16-25. DOI: 10.21199/WB50.1.2 <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/820/781>
- Smith, James P. Jr. (2019). California floras, manuals, and checklists: A bibliography. *Botanical Studies* 70. https://digitalcommons.humboldt.edu/botany_jps/70
- Thorne, J. H., Viers, J.H., Price, J., & Stoms, D. M. (2009). Spatial Patterns of Endemic Plants in California. *Natural Areas Journal* 29(4), 344 – 66. <https://doi.org/10.3375/043.029.0402>.
- Vaughan, T.A. (1954). *Mammals of the San Gabriel Mountains of California*. University of Kansas Publications, Museum of Natural History. Volume: 7(9). <https://www.gutenberg.org/cache/epub/34848/pg34848.txt>
- Vickers, T.W., Sanchez, J.N., Johnson, C.K., Morrison, S.A., Botta, R., Smith, T., Cohen, B.S., Huber, P.R., Ernest, H.B., & Boyce, W.M. (2015). Survival and mortality of pumas (*Puma concolor*) in a fragmented, urbanizing landscape. *PLoS One*, 10(7), e0131490. <https://doi.org/10.1371/journal.pone.0131490>
- White, M.D., Stallcup, J.A., Spencer, W.D., Strittholt, J.R., & Heilman, G.E. (2003). *Conservation significance of Tejon Ranch, a biogeographic crossroads*. Prepared by the Conservation Biology Institute with South Coast Wildlands Project and California Wilderness Coalition for Environment Now, Santa Monica, CA.
- SEA Background and Historical Context**
- County of Los Angeles. (1980). *Technical Supplement E, Significant Ecological Areas/Habitat Management Areas in Los Angeles County*. Pp. E-1 to E-43 In: County of Los Angeles General Plan Technical Supplement. November 1980.
- County of Los Angeles. (2022c). *Significant Ecological Program, Significant Ecological Areas (SEA) and Coastal Resource Areas*. Map. January 2022.

DRAFT

England & Nelson Environmental Consultants. (1976). *Los Angeles County Significant Ecological Area Study, 1976*. Prepared for Los Angeles County Department of Regional Planning and Environmental Systems Research Institute. 50 pp. + Appendices (A-E).

PCR. (2000). *Los Angeles County Significant Ecological Area Update Study 2000. Background Report*. Prepared for Los Angeles County Department of Regional Planning by PCR Services Corporation, Frank Hovore & Associates, and FORMA Systems. Nov. 2000. 37 pp. + Appendices (A-E).

LA County General Plan and Related Information

City of Los Angeles. (2001). Conservation Element of the City of Los Angeles General Plan. Adopted by the City Council September 26, 2001.

County of Los Angeles, Department of Parks and Recreation. (2022). *Los Angeles Countywide Comprehensive Parks and Recreation Needs Assessment Plus (PNA+)*. Prepared by MIG. Adopted by the Los Angeles County Board of Supervisors on Dec. 6, 2022.

County of Los Angeles. (2022a). Appendix E: Conservation and Natural Resources Element Resources. (pp. 42-237). In *Los Angeles County 2035 General Plan*. County of Los Angeles. <https://planning.lacounty.gov/long-range-planning/general-plan/appendices/>.

County of Los Angeles. (2022b). Chapter 9: Conservation and Natural Resources Element. (pp. 11-48). In *Los Angeles County 2035 General Plan*. County of Los Angeles. <https://planning.lacounty.gov/long-range-planning/general-plan/general-plan-elements/>

SCAG (Southern CA Association of Governments). (2014). SCAG Conservation Framework and Assessment. Presentation prepared by Northwest Habitat Institute. Dec. 16, 2014. <https://scag.ca.gov/sites/default/files/2024-05/oscwq111914nhi.pdf>

Climate Change and Adaptation

Allen, E.B., Temple, P.J., Bytnerowicz, A., Arbaugh, M.J., Sirulnik, A.G., & Rao, L.E. (2007). Patterns of understory diversity in mixed coniferous forests of southern California impacted by air pollution. *The Scientific World Journal*, 7, 247 – 63. <https://doi.org/10.1100/tsw.2007.72>.

California Climate Insurance Working Group. (2021, January). *Protecting communities, preserving nature, and building resilient economies: California climate insurance report*. California Department of Insurance. <https://www.insurance.ca.gov/01-consumers/180-climate-change/>. Accessed July 28, 2025.

Gardali, T., Seavy, N.E., DiGaudio, R.T., & Comrack, L.A. (2012). A climate change vulnerability assessment of California's at-risk birds. *PloS One* 7(3): e29507. DOI 10.1371/journal.pone.0029507. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0029507>

Hennessy, S., Jennings, M., Molinari, N., Magee, C., Pairis, A., & Safford, H. (2024). *Climate-adapted conservation strategy for southern California montane forests*. Zenodo. <https://doi.org/10.5281/ZENODO.13129913> (N.d.).

Iknayan, K.J., & Beissinger, S.R. (2018). Collapse of a desert bird community over the past century

DRAFT

- driven by climate change. *Proceedings of the National Academy of Sciences* 115(34), 8597 – 602. <https://doi.org/10.1073/pnas.1805123115>.
- Jenning, M., et al. (2019). *Climate Resilient Connectivity for the South Coast Ecoregion of California. Final Report*. Prepared for California Department of Fish and Wildlife. State Wildlife Grant #F16AF00551; Project #G1698064). San Diego State University.
- Jordan, S.E. (2023). *Microclimatic Refugia of Chaparral Relicts in Mountains of the Mojave Desert*. A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in Biology. California State University, Northridge.
- Jordan, S.E. & Rachman, R. (2024). Response of an avifaunal community to the La Tuna Fire in the Verdugo Mountains of southern California. *Western Birds*, 55:116 – 129. DOI 10.21199/WB55.2.3. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/937/892>
- Lauer, A., English, D., & Richardson, M.H. (2023). How ‘green energy’ is threatening biodiversity, human health, and environmental justice: An example from the Mojave Desert, California. *Sustainable Environment* 9(1), 2192087. <https://doi.org/10.1080/27658511.2023.2192087>.
- Los Angeles County Chief Sustainability Office. (2021). *Los Angeles County Climate Vulnerability Assessment*.
- Rojas, I.M., Jennings, M., Conlisk, E., Syphard, A.D., Mikesell, J., Kinoshita, A.M., West, K., Stow, D., Storey, E., De Guzman, M.E., Foote, D., Warneke, A., Pairis, A., Ryan, S., Flint, L.E., Flint, A.L. & Lewison, R. (2022). A landscape-scale framework to identify refugia from multiple stressors. *Conservation Biology* 36, e13834. <https://doi.org/10.1111/cobi.13834>
- Thorne, J.H., Boynton, R.M., Holquin, A.J., Stewart, J.A.E., & Bjorkman, J. (2016). *A climate change vulnerability assessment of California’s terrestrial vegetation*. California Department of Fish and Wildlife, Sacramento, CA.
- Unitt, P., & Hargrove, L. (2018). *Southward and downslope extensions of breeding ranges of birds in southern California*. In W.D. Shuford, R.E. Gill Jr., & C.M. Handel (Eds.), *Trends and traditions: Avifaunal change in western North America* (pp. 85 – 115). *Studies of Western Birds*, 3. Western Field Ornithologists, Camarillo, CA. DOI 10.21199/SWB3.4. https://westernfieldornithologists.org/docs/2020/Avifaunal_Change/Unitt-Hargrove/Unitt-Hargrove-Avifaunal_Change.pdf
- Connectivity**
- Boydston, E., Ordeñana, M., & Cooper, D.S. (2014). Landscape connectivity for medium and large mammals in the City of Los Angeles. *WatershedWise* 15(4).
- California Department of Fish and Wildlife. (2022). *Restoring California’s Wildlife Connectivity 2022. Biogeographic Data Branch*. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=204648&inline>. Accessed July 28, 2025.
- California Department of Fish and Wildlife. (2025, April 29). *Terrestrial Connectivity (Areas of Conservation Emphasis, ACE) [dataset]*. CDFW Open Data Portal.

DRAFT

https://wildlife.ca.gov/Data/Analysis/Connectivity?utm_source=chatgpt.com#589603664-terrestrial-connectivity-ace

California Department of Transportation. (2024). *Caltrans Wildlife Connectivity Program Report*. Headquarters Division of Environmental Analysis, Office of Fish and Wildlife Connectivity. <https://dot.ca.gov/-/media/dot-media/programs/environmental-analysis/documents/env/caltrans-wildlife-connectivity-report-ally.pdf>. Accessed July 28, 2025.

County of Los Angeles, Department of Regional Planning. (2018). *Guidance for Evaluating Impacts on Wildlife Movement in LA County*. Environmental Planning and Sustainability Section.

Mathewson, P., Spehar, S., & Cooper, D.S. (2008). An initial carnivore survey of Griffith Park, Los Angeles. *Bulletin of the Southern California Academy of Sciences*, 107, 57-67. DOI: 10.3160/0038-3872(2008)107[57:AICSOG]2.0.CO;2

Northwest Habitat Institute. (2024). *Conservation Framework and Assessment*. Prepared for the Southern California Association of Governments.

Penrod, K., Cabañero, C., Beier, P., Spencer, W., Luke, C., & Rubin, E. (2003). *South Coast Missing Linkages Project: A Linkage Design for the Tehachapi Connection*. Idyllwild, California: South Coast Wildlands. http://www.scwildlands.org/reports/SCML_Tehachapi_finalreport.pdf

Penrod, K., Beier, P., Garding, E., & Cabañero, C. (2012). *A Linkage Network for the California Deserts*. Prepared for Bureau of Land Management and The Wildlands Conservancy by Science and Collaboration for Connected Wildlands, Fair Oaks, CA, and Northern Arizona University, Flagstaff, AZ.

Spencer, W.D. (2005). *Maintaining Ecological Connectivity Across the “Missing Middle” of the Puente-Chino Hills Wildlife Corridor*. July 2005. Prepared by the Conservation Biology Institute for the Resources Legacy Fund Foundation, Sacramento, California. <https://consbio.org/reports/maintaining-ecological-connectivity-across-the-missing-middle-of-the-puente-chino-hills-wildlife-corridor/>

UCLA Institute of the Environment and Sustainability. (2023). *Quantifying and Validating Habitat Connectivity across Greater Los Angeles: Final Report for the City of Los Angeles & The Nature Conservancy*.

Regional and Local Studies and Reports

Anderson, I. (2009). Tejon Ranch—a conservation priority. *Fremontia* 37(1), 16-20.

Bilodeau, W. L., Bilodeau, S. W., Gath, E. M., Osborne, M., & Proctor, R. J. (2007). Geology of Los Angeles, California, United States of America. *Environmental and Engineering Geoscience* 13(2), 99 – 160. <https://doi.org/10.2113/gseegeosci.13.2.99>.

Bond, M. & Bradley, C. (2005). *Recommended Critical Biological Zones in Southern California’s Four National Forests*. Unpublished report. https://www.biologicaldiversity.org/programs/public_lands/forests/southern_california_forests/pdfs/hotspots.pdf

Bonebrake, T.C. & Cooper, D.S. (2014). A Hollywood drama of butterfly extirpation and persistence over a

DRAFT

- century of urbanization. *Journal of Insect Conservation*, 18(4), 683-692. DOI: 10.1007/s10841-014-9675-z
- Boyd, S. (1999). Vascular flora of the Liebre Mountains, western Transverse Ranges, California. Occasional Publ. No. 5. Rancho Santa Ana Botanic Garden, Claremont, Ca. Reprint of *Aliso*, 18(2), 93-139 + xii.
- Bureau of Land Management (BLM) California Desert District. (2004). *Final Environmental Impact Report and Statement for the West Mojave Plan, A Habitat Conservation Plan and California Desert Conservation Area Plan Amendment Vol 1*. https://eplanning.blm.gov/public_projects/lup/72544/97517/117675/wemo_plan_vol-1_2004.pdf
- California Native Plant Society & Center for Food Safety. (n.d.). *The world of seeds: Biodiversity and the role of seed saving*. Retrieved from materials provided by Los Angeles County.
- California Resources Agency, San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy, and Santa Monica Mountains Conservancy. (2001). *Common Ground from the Mountains to the Sea. Watershed and Open Space Plan San Gabriel and Los Angeles Rivers*.
- Campbell, J.E., Sharifi, M.R., & Rundel, P.W. (2017). Impact of ground water depletion on the mesquite community at Edwards Air Force Base, western Mojave Desert, California. *Aliso* 35 (2), 69 – 77. <https://doi.org/10.5642/aliso.20173502.03>.
- Cooper Ecological Monitoring, Inc. (2021). *Rapid Assessment of Wildland-Urban Interface Parcels for Conservation: Sierra Madre to La Verne in Los Angeles County*. Prepared for Watershed Conservation Authority, Azusa, CA. July 22, 2021.
- Cooper, D.S. (2004). *Important Bird Areas of California*. Audubon California, Pasadena. 286 pp.
- Cooper, D.S. (2011). Rare plants of Griffith Park, Los Angeles, California. *Fremontia* 38(4)/39(1),18-24.
- Cooper, D.S. (2011). Biological Assessment of Padua Park and regional reference sites. Prepared for City of Claremont, Nov. 4, 2011.
- Cooper, D.S. (2012). *Weed Management Plan for the Baldwin Hills, Los Angeles County*. Prepared for Council for Watershed Health, October 5, 2012.
- Cooper, D.S. (2015). Flora of Griffith Park, Los Angeles, California. *Crossosoma* 41(1&2): 1-86.
- Cooper, D.S. (2016). *Industrial-scale Solar Projects and Birds in the California Desert: Assessing Impacts and Developing Mitigation (Final)*. Prepared for Sonoran Joint Venture (USFWS). May 5, 2016.
- Cooper, D.S. (2024). *Chatsworth Nature Preserve Preliminary Biological Survey and Resource Management Plan Framework*. Prepared for Save Chatsworth. June 2024.
- Cooper, D.S. & C. Aiken (McCammon). (2018). *Puente Hills Wildlife Camera Study, 2018 Report (Final)*. Prepared for Puente Hills Habitat Preservation Authority, Whittier, CA. Aug. 6, 2018.
- Cooper, D.S. & Fiesler, E. (2012). *Madrona Marsh Preserve Biological Inventory (Final Report)*. Prepared for Friends of Madrona Marsh and City of Torrance, January 27, 2012.

DRAFT

- Cooper, D.S. & Hamilton, R.A. (2014). *A Conservation Analysis for the Santa Monica Mountains “Coastal Zone” in Los Angeles County*. Appendix A (Biota) in: Santa Monica Mountains Local Coastal Program, January 2014. <http://file.lacounty.gov/bos/supdocs/82928.pdf>
- Cooper, D.S. & Hamilton, R.A. (2015). *A Conservation Vision for Los Cerritos Wetlands, Los Angeles/Orange County, California (final draft)*. Prepared by D.S. Cooper & R.A. Hamilton for Audubon California. Oct. 27, 2015.
- Cypher, B.L., Job, C.V.H, Tennant, E.E., & Phillips, S.E. (2010). *Mammalian Species Surveys in the Acquisition Areas on the Tejon Ranch, California*. Unpublished Report. August 16, 2010.
- Dickey, R. (2009). The last floral blooms: Something to see, while they are still there. *Fremontia* 37(1), 3-15.
- Dimson, M., Harrigan, R., & Gillespie, T. W. (2023). *Biodiversity Atlas of Los Angeles* [Interactive website]. UCLA Sustainable LA Grand Challenge. <https://sustainablela.ucla.edu/research-portal/project/biodiversity-atlas-la>. Accessed July 28, 2025.
- Ethington, P.J., MacDonald, B., Stein, G., Deverell, W., & Longcore, T. (2020). *Historical Ecology of the Los Angeles River Watershed and Environs*. Report to the John Randolph Haynes and Dora Haynes Foundation, June 15, 2020. Spatial Sciences Institute University of Southern California Los Angeles.
- Hamilton Biological, Inc. (2019). *Biological Resources Report, City of Diamond Bar*. Biological technical report dated February 25, 2019, addressing the conservation and preservation of sensitive biological resources in the City of Diamond Bar and its Sphere of Influence. Prepared for a consortium of Diamond Bar residents for consideration by the City of Diamond Bar during an update of the City’s General Plan.
- ICF. (2021). *Antelope Valley Regional Conservation Investment Strategy (AVRCIS). Final*. November. (ICF 313.19.) San Diego, California. Prepared for Desert Mountains Conservation Authority, Quartz Hill, California.
- Los Angeles County Oak Woodlands Habitat Conservation Strategic Alliance. (2011). *Los Angeles County Oak Woodlands Conservation Management Plan*. Prepared for County of Los Angeles Department of Regional Planning.
- Los Angeles County Sanitation Districts. (2025). Santa Clara River Temperature Study. TAC Meeting (presentation). August 13, 2025. [https://ftp.sccwrp.org/pub/download/PROJECTS/Temperature Effects Study/Santa%20Clara/Meetings/2025-08-13%20SCR%20TAC%20Meeting.pdf](https://ftp.sccwrp.org/pub/download/PROJECTS/Temperature%20Effects%20Study/Santa%20Clara/Meetings/2025-08-13%20SCR%20TAC%20Meeting.pdf)
- Los Angeles Sanitation and Environment. (2025). *2025 LASD Biodiversity Index Monitoring Report*. City of Los Angeles.
- Mistretta, O. (2020). *Field guide to the flora of the San Gabriel Mountains*. Rancho Santa Ana Botanic Garden Occasional Publications, Number 18.
- National Park Service. (2011). *Draft San Gabriel Watershed and Mountains Special Resource Study & Environmental Assessment*.

DRAFT

- Power Engineers. (2011). Barren Ridge Renewable Transmission Project: Land Use Technical Report. Prepared for: LADWP. June 2011.
- Raven, P.H., Thompson, H.J. and Prigge, B.A. (1986). Flora of the Santa Monica Mountains. Southern California Botanists Special Publication No. 2, Fullerton, CA. 181 pp.
- Santa Monica Mountains Conservancy. (2021). *Eastern Santa Monica Mountains Natural Resource Protection Plan Adopted by the Santa Monica Mountains Conservancy State of California – December 13, 2021*. Unpublished report. <https://smmc.ca.gov/wp-content/uploads/2021/12/ESSM-NRPP.pdf>.
- Schneider-Ljubenkov, J. A. & Ross, T. S. (2001). An annotated checklist of the vascular plants of the Whittier Hills, Los Angeles County, California. *Crossosoma* 33, 55-69.
- Southern California Association of Governments. (2020). *Natural and Farm Lands Conservation Technical Report*, part of *Connect SoCal (2020 – 2045 RTP/SCS)*. Prepared by Northwest Habitat Institute.
- Soza, V. L., Gross, L, Boyd, S., & Fraga, N. (2013). Vascular flora of the Verdugo Mountains and San Rafael Hills, Los Angeles County, California. *Crossosoma* 39(1 & 2), 1-140.
- Stallcup, J.A, White, M.D., Stritholt, J.R., & Spencer, W.D. (2003). *Conservation aAssessment of Tejon Ranch*. Prepared by the Conservation Biology Institute for Preserving Wild California, Resources Legacy Fund Foundation (Grant #2003-0116).
- Stein et al. (2007). Historical ecology and landscape change of the San Gabriel River floodplain. SCCWRP Tech. Report #499. https://ftp.sccwrp.org/pub/download/DOCUMENTS/TechnicalReports/499_historical_ecology.pdf
- Stephenson, J.R., & Calcarone, G.M. (1999). *Southern California Mountains and Foothills Assessment: Habitat and Species Conservation Issues*. Gen. Tech. Rep. GTR-PSW-172. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture; 402 p.
- Swinney, D. (1994). *Glendora Foothills Plant Checklist, San Gabriel Mountains*. 65 pp. Self-published.
- Terrill, R.S., Dean, C.A., Garrett, J., Maxwell, D.J., Hill, L., Farnsworth, A., Dokter, A.M., & Tingley, M.W. (2021). A novel locality for the observation of thousands of passerine birds during spring migration in Los Angeles County, California. *Western Birds*, 52, 322 – 339. DOI 10.21199/WB52.4.4. <https://journal.westernfieldornithologists.org/index.php/westernbirds/article/view/704/666>
- The River Project. (2022). *Sepulveda Basin Restoration Feasibility Report*. Prepared by The River Project in collaboration with Balance Hydrologics, Wildling Design Studio, SALT-LAGHD. 204 pp.
- Tirrell, J., Fiddler, W., Strong, J., & Bothwell, G. (2013). A historical comparison of habitat and plant populations in the mid-range San Gabriel Mountains. *Fremontia* 41(2), 2-9.
- TRIP. (2018). Los Angeles Transportation by the Numbers. <https://tripnet.org/wp->

DRAFT

[content/uploads/2018/08/CA_Los_Angeles_Transportation_by_the_Numbers_TRIP_Report_Aug_2018.pdf](#)

[U.S. Forest Service. Angeles National Forest. 2018. San Gabriel Mountains National Monument Management Plan. <https://npshistory.com/publications/usfs/san-gabriel-mountains/mp-2018.pdf>](#)

[U.S. Forest Service. Pacific Southwest Region. 2005. Final Environmental Impact Statement \(2 vol.\). Land management plans: Southern California national forests vision: Angeles National Forest, Cleveland National Forest, Los Padres National Forest, San Bernardino National Forest. <https://www.fs.usda.gov/r05/angeles/planning>](#)

[U.S. Census Bureau. \(2024\). QuickFacts: Los Angeles County, California. Population Estimates. <https://www.census.gov/quickfacts/fact/table/losangelescountycalifornia/POP010220>. Accessed March 5, 2026.](#)

[Vandergast, A.G., Bohonak, A.J., Hathaway, S.A., Boys, J., & Fisher, R.N. \(2008\). Are hotspots of evolutionary potential adequately protected in southern California? *Biological Conservation* 141, 1648 – 1664. <https://doi.org/10.1016/j.biocon.2008.04.009>.](#)

Coastal Resource Area Descriptions

The following are detailed descriptions of each of the nine Coastal Resource Areas:

Alamitos Bay CRA

Location

General

The Alamitos Bay CRA is located upstream of the Alamitos Bay Marina, off the Los Cerritos Channel of the City of Long Beach. The CRA boundaries encompass the Alamitos Bay salt marsh, which is one of three remaining examples of salt marshes found in the County. The entire CRA area has been identified by the California Audubon as a Globally Important Bird Area (IBA)—part of the Los Cerritos Wetlands and adjacent oil fields section of the Orange Coast Wetlands unit. The entire CRA is located within the United States Geological Survey (USGS) 7.5' California Los Alamitos Quadrangle.

General Boundary and Resources Description

The CRA boundaries generally follow the Alamitos Bay salt marsh area, which is bordered by an oil field, main artery roads, the Cerritos Channel, and residential development. The CRA is situated to the west of Studebaker Road, north of Westminster Avenue, east of Pacific Coast Highway and south of Loynes Drive and the Cerritos Channel within the City of Long Beach.

The CRA is one of three remaining examples of salt marsh found in the County, and the last remnant of the extensive salt marshes once found in Los Alamitos Bay. The majority of this vegetation type has been lost to urbanization, flood control projects, harbors, and marinas. It is one of the most productive ecological communities and is an important breeding ground for terrestrial and marine organisms, including the majority of commercial fishes. This is due in part to the estuaries and salt marshes

DRAFT

interfacing between the terrestrial and marine worlds, and serving as important nutrient cycling centers for marine ecosystems. The Belding's savannah sparrow occurs in Alamitos Bay salt marsh. This species is restricted to salt marsh habitat, and has been placed on the state-endangered species list. The Orange Coast Wetlands IBA is believed to harbor one-eighth of the population of Belding's savannah sparrows, and there is a significant amount in Alamitos Bay. Least terns and other terns that breed in the area often use Alamitos Bay and the Cerritos arm as a foraging area. This type of habitat is an important wintering ground for migratory birds. It is estimated that in the fall and spring seasons, the Orange Coast Wetlands IBA hosts 15,000-20,000 individual migrating birds.

Vegetation

As discussed, the CRA encompasses the last remaining coastal salt marsh in Los Alamitos Bay's formerly extensive system of salt marshes and is one of only three such examples of this habitat remaining in County. Sensitive plant species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Coastal Salt Marsh: Consists of salt-tolerant plants that are mostly low-growing herbaceous perennials that are found on the borders of marine salt water bodies. The duration and extent of tidal inundation or influence causes a graduation in the prevalence of various species within this community. In the Alamitos Bay wetlands, this includes cordgrass as the dominant species and depending on the conditions, pickleweed, salt grass, alkali heath and spearscale can also be found.

Corresponding MCV communities:

- *Sarcocornia pacifica* (*Salicornia depressa*) ([*Salicornia pacifica*, *Salicornia depressa*] pickleweed mats) Herbaceous Alliance
- *Bolboschoenus maritimus* (salt marsh bulrush marshes) Herbaceous Alliance
- *Distichlis spicata* (salt grass flats) Herbaceous Alliance
- *Spartina foliosa* (California cordgrass marsh) Herbaceous Alliance

Intertidal Flats: Brackish coastal wetlands of low-lying basins of high evaporation and infrequent inputs of freshwater with low-growing salt tolerant plants.

Corresponding MCV Communities:

- *Ruppia (cirrhosa, maritima)* (ditch-grass or widgeon-grass mats) Aquatic Herbaceous Alliance

Wildlife

Coastal salt marshes and estuaries are productive habitats, which are used for foraging and breeding grounds, for both resident and migrating wildlife species. Estuaries and coastal salt marshes are the interface between the terrestrial and marine worlds, and are important nutrient recycling centers for marine ecosystems. In the past, this habitat was once extensive in the Los Alamitos Bay area.

Although little documentation regarding the types of animals present has been found, based on the apparent health of the ecosystem, it may be assumed that fishes that are commonly found in the vicinity can also be found in the CRA. These may include species, such as arrow goby, California halibut, cheekspot goby, diamond turbot, queenfish, shadow goby, shiner perch, topsmelt, longjaw

DRAFT

mudsucker, Pacific staghorn sculpin, and yellowfin goby.

Without more information, it is not possible to predict whether any reptiles or mammals can be found, but it is likely that amphibians, such as Baja California chorus frogs, are present.

The CRA habitat probably supports a variety of bird species found in the few coastal saltmarshes that remain in coastal Southern California. Belding's savannahsparrow has been observed in the CRA. Shallow water habitat exists and would be expected to attract wading birds and ducks. Foraging habitat that is attractive to raptors appears to be present on the outside perimeter of the marsh.

Wildlife Movement

The CRA provides a variety of saltwater, estuarine, mudflat and freshwater marsh habitats, and is an important stopover for many migratory birds traveling the Pacific Flyway migration route. Its suitability for many fishes and invertebrates allows populations that are capable of supporting further colonization and expansion of range. The area does not fall within any identified terrestrial movement routes for wildlife.

Sensitive Biological Resources

Sensitive biological resources are habitats or individual species that have special recognition by federal, state, or local conservation agencies and organizations as endangered, threatened, and/or rare. This is due to the species' declining or limited population sizes, which usually results from habitat loss. Watch lists of such resources are maintained by the California Department of Fish and Wildlife (CDFW), the United States Fish and Wildlife Service (USFWS), and special groups, such as the California Native Plant Society (CNPS). The following sections indicate the habitats as well as plant and animal species present, or potentially present within the CRA, which have been accorded special recognition.

Sensitive Plant Communities and Habitats

The CRA supports several habitat types considered sensitive by resource agencies. These are inventoried by California Department of Fish and Wildlife (CDFW) in the California Natural Diversity Database (CNDDB), or have been assigned a conservation status rank of 1 – 3 at the global or subnational scale. Conservation status ranks are assigned by CDFW, following methodology developed by NatureServe (<http://www.natureserve.org/conservation-tools/standards-methods/conservation-status-assessment>). Within this SEA, the CRA is configured to encompass the regionally significant community of a coastal salt marsh or coastal brackish marsh. This community or closely related designations are considered highest priority communities by the CDFW, indicating that they are declining in acreage throughout their range due to land use changes. The array and composition of these communities have been discussed in the Vegetation section. Changes to the classification system mentioned earlier in some cases divide plant communities into many possible vegetation alliances—not all of which may be considered sensitive. Previously listed communities with at least one sensitive alliance in the new format have been listed.

Sensitive Plant Species

The following special-status plant taxa have been reported or have the potential to occur within the CRA, based on known habitat requirements and geographic range information:

- Davidson's saltscale (*Atriplex serenana* var. *davidsonii*) RPR 1B.2

DRAFT

- Southern tarplant (*Centromadia parryi ssp. australis*) RPR 1B.1
- Salt marsh bird's-beak (*Chloropyron maritimum ssp. maritimum*) FE, SE, RPR 1B.2
- Coulter's goldfields (*Lasthenia glabrata ssp. coulteri*) RPR 1B.1
- Coast woolly-heads (*Nemacaulis denudata var. denudata*) FT, RPR 1B.1
- California Orcutt grass (*Orcuttia californica*) FE, SE, RPR 2.1
- Estuary seablite (*Suaeda esteroa*) RPR 1B.2
- San Bernardino aster (*Symphyotrichum defoliatum*) RPR 1B.2

Sensitive Animal Species

The following special-status animal species are reported or are likely to be present within the CRA based on habitat requirements and known range attributes:

- Belding's savannah sparrow (*Passerculus sandwichensis beldingi*) SE
- California least tern (*Sternula antillarum browni*) FE, SE, ABC, CDFW Fully Protected

CRA Regional Biological Value

The CRA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE ALAMITOS BAY CRA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	Los Alamitos Bay is one of only three remaining salt marshes found in the County, and a remnant of extensive salt marshes once found in its area. (The others are the Ballona Wetlands off of Santa Monica Bay and the Malibu Lagoon.) It is the type habitat of Belding's savannah sparrow, which is an endangered species that is still found in the much reduced habitat of the salt marshes of Southern California.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Salt marsh is reduced and fragmented from its former coverage in Southern California, so it is important to preserve any of these important estuarine areas.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that	Met	Los Alamitos Bay is one of only three remaining salt marshes found in the County, which is a remnant of extensive salt marshes once found in the area.

DRAFT

	Criterion	Status	Justification
	are either unique or are restricted in distribution.		
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The Belding's savannah sparrow lives its entire life in salt marshes and breeds, rests, and feeds in this area. Salt marshes are important nursery grounds for many marine animals, and diminished fisheries for many are attributed to the loss of salt marshes. Salt marshes are important habitat for migrating marine birds, which utilize the abundant forage produced by the marsh in the form of mud animals and insects.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	As one of only three salt marshes in the County, this area is scientifically interesting for the study of island bio-geography in that salt marshes are now islands, whereas they were once almost continuous in the seismically active coastline of Southern California. The three marshes differ from one another in dominant vegetation.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Los Alamitos Bay is one of only three remaining salt marshes found in the County, which is a remnant of extensive salt marshes once found in its area. The majority of this habitat type has been lost to urbanization, flood control projects, harbors, and marinas. Salt marshes are a very productive vegetative community and an important breeding and nursery area for marine, marsh, and terrestrial animals—an ecotone where multiple habitats meet and combine resident biota. Many commercially valuable fishes start life in salt marshes.

In conclusion, the area described in this report is a CRA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; and D) concentrated breeding, feeding, resting, or migrating grounds that are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.

Ballona Wetlands CRA

Location

General

The Ballona Wetlands CRA is located south of Marina del Rey, north of Playa Del Rey, and west and northwest of Playa Vista. One extending arm reaches north to the State Route-90 overcrossing and another reaches south to include the restored freshwater marsh adjacent to the Playa Del Rey and Playa Vista districts of the City of Los Angeles. The Ballona Wetlands are a remnant of what was the County's largest coastal lagoon. The Ballona watershed covers over 130 square miles, and the lagoon

DRAFT

area was so large (about 11-12 square miles) that it included freshwater peripheries. Incorporated in the lagoon complex were 10 kinds of habitat that ranged from coastal saltwater marsh to grassy prairie to oak and willow woodland adjacent to freshwater areas. The lagoon connected via Ballona Creek that sometimes was the Los Angeles River, to La Cienega, a large swampy area (about 13-14 square miles) that was north and east of the Baldwin Hills. The CRA lies at the base of the Ballona Creek watershed and includes part of the Ballona Creek flood control channel that drains the total of 130 square miles, from what is now a highly urbanized area. While the Ballona Wetlands ecosystem has been substantially degraded over the years due to human activity and urban development, it is still a rich ecological system that bridges the gap between aquatic marine and freshwater land environments. It provides crucial habitat for hundreds of plant and animal species.

The CRA is in part of the California Audubon-designated Ballona Wetlands State Important Bird Area (IBA). The IBA is more extensive than the CRA, and includes all of the Marina del Rey waterways, some of the Strand Beach to the south, Del Rey Lagoon and Ballona Lagoon, and extends up the Bluff Creek (source creek for the restored freshwater marsh) at the base of the south Ballona bluff as far as Centinela Avenue. This area is very important to avian life and has a great diversity of bird types ranging from those associated with salt water marsh to those associated with freshwater marsh to raptors that feed over grasslands.

The CRA is located on unincorporated land in the community of Marina del Rey north of the Ballona Creek Channel and in the City of Los Angeles south of the Ballona Creek Channel. It is within the United States Geological Survey (USGS) 7.5' California Venice Quadrangle.

General Boundary and Resources Description

The CRA is surrounded by urban and residential uses. The CRA is bordered by Fiji Way in Marina del Rey on the northeast, and has an arm that extends northeast to the State Route-90 overcrossing. This arm includes an undeveloped area of mixed native and non-native shrubs and grasses, and is bordered by Culver Boulevard on its south edge. The north boundary crosses the Ballona Creek Channel directly south of the end of Fiji Way, and goes west on the south side of Ballona Creek, which delineates the CRA's western half. The boundary turns southward, then eastward at the development in Playa Del Rey. The boundary goes north to Jefferson Boulevard along a gas line road, and then continues eastward on the north side of Jefferson Boulevard, as far as the area that has been restored as a freshwater marsh. The boundary turns southward and eastward to enclose the freshwater marsh. At the boundary of the freshwater marsh along Lincoln Boulevard, the boundary follows the west side of Lincoln Boulevard crossing Jefferson Boulevard, Ballona Wetlands, and connecting to the south side of the north arm where it crosses Culver Boulevard. Most of the area is a mixture of native and non-native shrubs and grasses, with a strong influence of brackish conditions. The northern side of the Ballona Creek Channel was a tidal mud flat in the original Ballona Lagoon that received a large amount of the dredge spoil when Marina del Rey was created. However, there is enough natural forage in the wetlands to sustain a breeding colony of great blue herons and a breeding colony of black-crowned night herons in the trees of nearby Marina del Rey.

Ballona Wetlands is one of three remaining remnants of salt marsh in the County (the other two are Malibu Lagoon in the Santa Monica Mountains SEA and the Cerritos Channel salt marsh in the Alamitos Bay CRA). This type of habitat is one of the most productive in the world, and is used as a breeding ground by many marine and terrestrial organisms. Belding's savannah sparrow (*Passerculus sandwichensis beldingi*), which is a state-recognized endangered species, resides in the pickleweed flats on the south side of the Ballona Wetlands. The California least tern (*Sternula antillarum browni*) breeds in the sandy areas around Ballona Lagoon and on Venice Beach, and is recognized as an

DRAFT

endangered species by the state and federal governments. The terns forage in the waterways that are included in the CRA.

The salt marsh, Ballona Creek Channel, Ballona Lagoon, and Del Rey Lagoon form an important complex of habitats that are heavily used by migratory birds. The area is recognized by ornithologists and bird watchers throughout the area for its rich birdlife during the spring and fall migrations, and during the winter season. This type of heavy use is common in salt marsh habitat, and has been made more intense by the loss of habitat in Marina del Rey, and throughout most of Southern California. This forces these birds to concentrate in the few remaining areas. Loss of this habitat type has led to reductions in the numbers of these birds present along the coast.

The salt marsh and lagoon at Ballona Creek are heavily used by academic institutions and conservation groups for educational field trips. This area serves as a type specimen of salt marsh habitat, and is the only easily accessible example in the County.

Vegetation

There are two categories of habitat in the CRA: wetland and upland altered or created by filling and grading. The wetland is characterized by coastal salt marsh and freshwater marsh. The upland is represented by a disturbed coastal sage scrub, non-native annual grassland, and coastal bluff and dune scrub. Sensitive plant species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Descriptions and general locations of the each plant community present within the CRA are given below.

Coastal Salt Marsh: Consists of salt tolerant plants that are mostly low-growing herbaceous perennials found on the borders of marine salt water bodies. The duration and extent of tidal inundation or influence causes a gradation of various species prevalence within this community. In the Ballona Wetlands this includes the areas where pickleweed (*Salicornia pacifica*) is a dominant species and depending on the conditions other plants such as California cordgrass (*Spartina foliosa*), salt grass (*Distichlis spicata*), alkali heath (*Frankenia salina*) and spearscale (*Atriplex prostrata*) can be found.

Corresponding MCV communities:

- *Sarcocornia* [*Salicornia*] *pacifica* (*Salicornia depressa*) (pickleweed mats) Herbaceous Alliance
- *Bolboschoenus maritimus* (salt marsh bulrush marshes) Herbaceous Alliance
- *Distichlis spicata* (salt grass flats) Herbaceous Alliance
- *Spartina foliosa* (California cordgrass marsh) Herbaceous Alliance

Intertidal Flat: Brackish coastal wetlands of low-lying basins of high evaporation and infrequent inputs of freshwater with low-growing salt tolerant plants.

Corresponding MCV Communities:

- *Ruppia* (*cirrrosa*, *maritima*) (ditch-grass or widgeon-grass mats) Aquatic Herbaceous Alliance

Freshwater Marsh: Develops in areas of still or slow-moving permanent freshwater. In the CRA, this

DRAFT

community may be dominated by perennial, emergent cattails (*Typha* spp.), which reach a height of four to five meters and often form a closed canopy. Bulrushes (*Schoenoplectus* spp.) are dominant below the cattail canopy. Freshwater marsh is relatively uncommon; it occurs in small patches in natural or created sinks with water sources.

Corresponding MCV communities:

- *Lepidium latifolium* (perennial pepper weed patches) Semi-Natural Herbaceous Stands
- *Schoenoplectus californicus* (California bulrush marsh) Herbaceous Alliance
- *Typha* (*angustifolia*, *domingensis*, *latifolia*) (cattail marshes) Herbaceous Alliance
- *Sarcocornia* [*Salicornia*] *pacifica* (*Salicornia depressa*) (pickleweed mats) Herbaceous Alliance
- *Juncus articus* (var. *balticus*, *mexicanus*) [*Juncus balticus* ssp. *ater*, *J. mexicanus*] (Baltic and Mexican rush marshes) Herbaceous Alliance
- *Juncus effusus* (soft rush marshes) Herbaceous Alliance
- *Lemna* (*minor*) and relatives (duckweed blooms) Provisional Herbaceous Alliance

Coastal Sage Scrub: Occurs in Southern California where moisture is available in the upper horizons during the winter-spring growing season. Plants that are adapted to these conditions are a mixture of herbaceous and weakly woody, shrubby and drought deciduous types. This includes species such as California sagebrush (*Artemisia tridentata*), coyote brush (*Baccharis pilularis*) and laurel sumac (*Malosma laurina*).

Corresponding MCV communities:

- *Artemisia californica* (California sagebrush scrub) Shrubland Alliance
- *Artemisia californica*-*Salvia mellifera* (California sagebrush-black sage scrub) Shrubland Alliance
- *Artemisia californica*-*Eriogonum fasciculatum* (California sagebrush-California buckwheat scrub) Shrubland Alliance
- *Encelia californica* (California brittle bush scrub) Shrubland Alliance
- *Salvia mellifera* (black sage scrub) Shrubland Alliance
- *Eriogonum cinereum* (ashy buckwheat scrub) Shrubland Alliance
- *Eriogonum fasciculatum* (California buckwheat scrub) Shrubland Alliance
- *Hazardia squarrosa* (sawtooth goldenbush scrub) Shrubland Alliance
- *Lotus scoparius* [*Acmispon glaber*] (deer weed scrub) Shrubland Alliance
- *Lupinus albifrons* (silver bush lupine scrub) Shrubland Alliance

DRAFT

Coastal Bluff and Dune Scrub: A remnant of the El Segundo Dune system that occupied coastal areas along much of the Santa Monica Bay. The system consists of fine, sandy soil that dries rapidly. Plants typical of the dune scrub include dune buckwheat (*Eriogonum parvifolium*), rattle pod (*Astragalus trichopodus* var. *lonchus*), bladderpod (*Peritoma arborea*), deer weed (*Acemisson glaber*), sawtooth goldenbush (*Hazardia squarrosa*), and California sunflower (*Helianthus californicus*)

Corresponding MCV communities:

- *Baccharis pilularis* (coyote brush scrub) Shrubland Alliance
- *Lupinus arboreus* (yellow bush lupine scrub) Shrubland Alliance and Semi-Natural Shrubland Stands
- *Lupinus chamissonis*-*Ericameria ericoides* (silver dune lupine–mock heather scrub) Shrubland Alliance

Non-native Grassland Communities: Consist of low, herbaceous vegetation that are dominated by invasive grasses that are primarily of Mediterranean origin, but can also harbor native forbs and bulbs, as well as naturalized annual forbs. Species found within this community include wild oat (*Avena fatua*), slender oat (*Avena barbata*), red brome (*Bromus madritensis* ssp. *rubens*), riggut brome (*Bromus diandrus*), and herbs such as black mustard (*Brassica nigra*) and wild radish (*Raphanus raphanistrum*). This community can be found throughout the CRA, but are less common in the moist soils found in and close to the salt and fresh water marshes.

Corresponding MCV communities:

- *Avena (barbata, fatua)* Semi-Natural Herbaceous Stands
- *Brassica (nigra)* and other mustards Semi-Natural Herbaceous Stands
- *Bromus (diandrus, hordeaceus) -Brachypodium distachyon* Semi-Natural Herbaceous Stands
- *Bromus rubens*-*Schismus (arabicus, barbatus)* ([*Bromus madritensis* ssp. *rubens*] red brome or Mediterranean grass grasslands) Semi-Natural Herbaceous Stands
- *Centaurea (solstitialis, melitensis)* (yellow star-thistle fields) Semi-Natural Herbaceous Stands
- *Lolium perenne [Festuca perennis]* (perennial rye grass fields) Semi-Natural Herbaceous Stands

Wildlife

Coastal salt marshes are productive habitats that are used for foraging and breeding grounds, for both permanent resident and migrating wildlife species. The CRA is an important site due to the rarity of this type of habitat, which was once more extensive in Southern California.

Studies in the last 30 years in the vicinity have identified a total of 44 species of fishes. Types found include arrow goby (*Clevelandia ios*), California halibut (*Paralichthys californicus*), cheekspot goby (*Ilypnus gilberti*), diamond turbot (*Hypsopsetta guttulata*), queenfish (*Seriphus politus*), shadow goby (*Acentrogobius nebulosus*), shiner perch (*Cymatogaster aggregata*), topsmelt (*Atherinops affinis*), longjaw mudsucker (*Gillichthys mirabilis*), Pacific staghorn sculpin (*Leptocottus armatus*), and yellowfin goby (*Acanthogobius flavimanus*).

DRAFT

Many native species of reptiles and amphibians have been found in the CRA. These include common kingsnake, San Diego gopher snake (*Pituophis catenifer annectens*), western side-blotched lizard (*Uta stansburiana elegans*), San Diego alligator lizard (*Elgaria multicarinata webbiai*), Great Basin fence lizard (*Sceloporus occidentalis longipes*), California slender salamander (*Batrachoseps attenuatus*) and California legless lizard (*Anniella pulchra*). Baja California chorus frog (*Pseudacris hypochondriaca*) and California toad (*Anaxyrus halophilus*) are very common in season. No invasive species of amphibians or reptiles have been reported in the CRA.

Native species of mammals found included western harvest mouse (*Reithrodontomys megalotis* (western harvest mouse), Botta's pocket gopher (*Thomomys bottae*), desert cottontail (*Sylvilagus audubonii*), striped skunk (*Mephitis mephitis*), raccoon (*Procyon lotor*), California ground squirrel (*Spermophilus beecheyi*), and the sensitive species south coast marsh vole (*Microtus californicus stephensi*). Invasive species found include house mouse (*Mus musculus*), black rat (*Rattus rattus*), Norway rat (*R. norvegicus*), domestic cat (*Felis domesticus*), Virginia opossum (*Didelphis virginiana*), and domestic dog (*Canis familiaris*). Low populations of small mammals are likely to be due to depredation by introduced red foxes, which are an immediate threat to the continued viability of the Ballona ecosystem.

The Ballona Wetlands provide habitat for a variety of bird species attracted to the few coastal saltmarshes that remain in coastal Southern California. Shallow water habitat for wading birds and ducks occurs in the northwestern portion of the CRA. No deep water is present for diving ducks and other birds that dive from the surface for fish. Foraging habitat for most raptors occurs in the western portion. Many bird species forage in the CRA, and the wetlands are used during migration and nesting. A resident population of Belding's savannah sparrow, which is a state-endangered species, was present in 2006 and the state and federally-endangered California least tern and fully protected American peregrine falcon (*Falco peregrinus anatum*) have been known to forage in this CRA. In 2010, least Bell's vireos (*least Bell's vireo*) successfully nested at the freshwater marsh, and many migrants have been observed there.

Sensitive wildlife species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Wildlife Movement

The CRA is an important stopover for many migratory birds traveling the Pacific Flyway migration route. For many birds this area, it is important because it provides a variety of salt-water, estuarine, mudflat and freshwater marsh habitats. It offers many fishes and invertebrates an opportunity to establish populations capable of supporting further colonization to expand their range. The area does not fall within any identified terrestrial movement routes for wildlife.

Sensitive Biological Resources

Sensitive biological resources are habitats or individual species that have special recognition by federal, state, or local conservation agencies and organizations as endangered, threatened, and/or rare. This is due to the species' declining or limited population sizes, which usually results from habitat loss. Watch lists of such resources are maintained by the California Department of Fish and Wildlife (CDFW), the United States Fish and Wildlife Service (USFWS), and special groups, such as the California Native Plant Society (CNPS). The following sections indicate the habitats as well as plant and animal species present, or potentially present within the CRA, which have been accorded special recognition.

Sensitive Plant Communities and Habitats

The CRA supports several habitat types considered sensitive by resource agencies. These are inventoried by California Department of Fish and Wildlife (CDFW) in the California Natural Diversity Database (CNDDDB), or have been assigned a conservation status rank of 1 – 3 at the global or subnational scale. Conservation status ranks are assigned by CDFW, following methodology developed by NatureServe (<http://www.natureserve.org/conservation-tools/standards-methods/conservation-status-assessment>). Within this SEA, the CRA is configured to encompass regionally significant communities, which include pickleweed mats, salt marsh bulrush marshes, California cordgrass marsh, ditch-grass or widgeon-grass mats, California brittle bush scrub, ashy buckwheat scrub, sawtooth goldenbush scrub, and silver dune lupine–mock heather scrub. These communities, or closely related designations, are considered high priority communities by the CDFW, which indicates that they are experiencing a decline throughout their range. The array and composition of these communities has been discussed in the Vegetation section.

Changes to the classification system mentioned earlier in some cases divides plant communities into many possible vegetation alliance communities, not all of which may be considered sensitive. For this CRA description, previously listed communities with a least one sensitive alliance in the new format have been listed.

Sensitive Plant Species

The statuses of rare plants are hierarchically categorized by the CNPS using a rank and decimal system. The initial category level of Rare Plant Rank is indicated by the ranks 1A (presumed extinct), 1B (rare or endangered in California and elsewhere), 2A (presumed extinct in California but extant elsewhere), 2B (rare or endangered in California but more common elsewhere), 3 (more information needed, a review list), and 4 (limited distribution). In cases where the CNPS has further identified the specific threat to the species, a decimal or Threat Code is added: .1 (seriously endangered in California), .2 (fairly endangered in California), or .3 (not very endangered in California).

The following special-status plant taxa have been reported or have the potential to occur within the CRA, based on known habitat requirements and geographic range information:

- *Aphanisma* (*Aphanisma blitoides*) RPR 1B.1
- Marsh sandwort (*Arenaria paludicola*) RPR FE, SE, 1B.1
- Ventura marsh milk-vetch (*Astragalus pycnostachyus* var. *lanosissimus*) FE, SE, RPR 1B.1
- South Coast saltscale (*Atriplex pacifica*) RPR 1B.2
- Parish's brittlescale (*Atriplex parishii*) RPR 1B.1
- Davidson's saltscale (*Atriplex serenana* var. *davidsonii*) RPR 1B.2
- Southern tarplant (*Centromadia parryi* ssp. *australis*) RPR 1B.1
- Orcutt's pincushion (*Chaenactis glabriuscula* var. *orcuttiana*) RPR 1B.1
- Coastal goosefoot (*Chenopodium littoreum*) RPR 1B.2

DRAFT

- Salt marsh bird's-beak (*Chloropyron maritimum* ssp. *maritimum*) FE, SE, RPR 1B.2
- Beach spectaclepod (*Dithyrea maritima*) ST, RPR 1B.1
- Los Angeles sunflower (*Helianthus nuttallii* ssp. *parishii*) RPR 1A
- Coulter's goldfields (*Lasthenia glabrata* ssp. *coulteri*) RPR 1B.1
- Mud nama (*Nama stenocarpum*) RPR 2B.2 Gambel's water cress (*Nasturtium gambelii*) FE, SE, RPR 1B.1
- Coast woolly-heads (*Nemacaulis denudata* var. *denudata*) RPR 1B.2
- Ballona cinquefoil (*Potentilla multijuga*) RPR 1A
- Estuary seablite (*Suaeda esteroa*) RPR 1B.2

Sensitive Animal Species

The following special-status animal species are reported or are likely to be present within the CRA based on habitat requirements and known range attributes:

- Mimic tryonia (*Tryonia imitator*) CDFW Special Animals List
- Wandering skipper (*Panoquina errans*) CDFW Special Animals List
- Silvery legless lizard (*Anniella pulchra pulchra*) FSS, SSC
- Coastal whiptail (*Aspidoscelis tigris stejnegeri*) CDFW Special Animals List
- San Bernardino ringneck snake (*Diadophis punctatus modestus*) FSS
- Western pond turtle (*Emys marmorata*) BLMS, FSS, SSC
- Two-striped garter snake (*Thamnophis hammondi*) BLMS, FSS, SSC
- Tricolored blackbird (*Agelaius tricolor*) BCC, BLMS, SSC, USBC, AWL, ABC
- Burrowing owl (*Athene cunicularia*) BCC, BLMS, SSC
- Western snowy plover (*Charadrius alexandrinus nivosus*) FT, BCC, SSC, ABC, AWL, USBC
- California black rail (*Laterallus jamaicensis coturniculus*) BCC, ST, CDFW Fully Protected, USBC, AWL, ABC
- Belding's savannah sparrow (*Passerculus sandwichensis beldingi*) SE
- California least tern (*Sternula antillarum browni*) FE, SE, CDFW Fully Protected, USBC, ABC
- Vireo bellii pusillus (*least Bell's vireo*) FE, SE, ABC

DRAFT

- South coast marsh vole (*Microtus californicus stephensi*) SSC
- Pacific pocket mouse (*Perognathus longimembris pacificus*) FE, SSC
- Southern California saltmarsh shrew (*Sorex ornatus salicornicus*) SSC

Regional Biological Value

The CRA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

Criteria Analysis of the Ballona Wetlands CRA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	Least Bell's vireos are beginning to use the restored freshwater marsh component for breeding. Given time, this could become one of the important breeding areas for them in the coastal area. Belding's savannah sparrow lives in the salt marsh savannah. Least terns breed nearby and forage year round in the shoal water areas of the CRA.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	There are three remaining salt marsh areas in the County: Cerritos Marsh in the Alamitos Bay CRA, Malibu Lagoon, and the CRA. In addition, there is a restored freshwater marsh on the periphery of Ballona Wetlands. The union of these two habitats is very rare in Southern California.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	There are three remaining salt marsh areas in the County: Malibu Lagoon, the marsh in Alamitos Bay CRA, and CRA. In addition there is a restored freshwater marsh on the periphery of Ballona Wetlands. Salt marsh coupled with freshwater marsh, is very rare in the County. There is no other such representation in the County.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	Endangered least Bell's vireos are using the freshwater marsh for breeding—they do not commonly occur in saltwater marshes, but were probably common on the peripheries in the freshwater portions. This restoration returns some of the former diversity and adds an important area for breeding. Many commercially valuable marine fishes start life in salt and estuarine marsh areas. The Ballona Wetlands are an important stopover point for migrations on the Pacific Flyway.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The union of the salt marsh and freshwater marsh habitats has no other representation in the County and is therefore of great scientific interest. As the only easily accessible saltwater marsh, the Ballona Wetlands have great educational value. This kind of ecotone is usually

DRAFT

			in very short supply, and will always be an extreme kind of physical habitat, a meeting point for the coastal strand, the brackish area of the marsh, and the freshwater area of the upland.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	NotMet	Without implementation of restoration activities, this CRA does not provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.

In conclusion, the area is a CRA because it contains A) habitat that hosts breeding for the endangered least Bell's vireo; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are unique and are restricted in distribution in the County and regionally; and D) concentrated breeding, feeding, resting, and migrating grounds, which are limited in availability in the County; and E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community

El Segundo Dunes CRA

Location

General

The El Segundo Dunes CRA is located between the west end of the Los Angeles International Airport (LAX) and the Pacific Ocean. The CRA is the largest remaining coastal dune habitat in Southern California. The federally-endangered El Segundo blue (*Euphilotes battoides allyni*) butterfly lives its entire life in these dunes.

The entire CRA is located in the United States Geological Survey (USGS) 7.5' California Quadrangle of Venice.

General Boundary and Resources Description

The CRA boundaries are as follows: Sandpiper Street to the north, Pershing Drive to the east, Imperial Highway to the south, and Vista Del Mar Road to the west. This is the largest of four proposed critical habitat areas for the El Segundo blue butterfly. The Vista Del Mar Park on Vista Del Mar Road is excluded from the CRA.

The ecological significance of the El Segundo Dunes is recognized by both federal and state governments, as well as by the City and County of Los Angeles and the California Coastal Commission (CCC). The vegetation found here does not occur anywhere else in the County, and is uncommon throughout Southern California. The vegetation is southern dune scrub, which is adapted to sandy, well-drained soils. The vegetation shows zonation, changing gradually as one moves from the foredunes facing west and the ocean over the dune crest and into coastal sage scrub. Many plants and invertebrates are specifically adapted and restricted to this environment and are not found elsewhere. One of these endemic organisms is the El Segundo blue butterfly, a federally-endangered butterfly species. The distribution of this butterfly is entirely restricted to the El Segundo Dunes and a few specks of dune habitat south to the edge of Malaga Cove (all once part of the El Segundo Dune system). Because of its rarity and highly limited range, the butterfly is officially recognized as an endangered species by the U.S. Fish and Wildlife Service. This small piece of dune habitat is extremely valuable as the final example of a community that was once more common along the County

and Southern California coastline.

Vegetation

Vegetation within the CRA is a remnant of a formerly more widespread distribution in the Los Angeles area of plant communities defined by their proximity to the immediate coast with its unique environmental conditions and sandy soils. The area has a long history of land disturbance and increasing isolation from related habitats. Today, the CRA has three plant communities with relatively few species. Importantly, the dune buckwheat (*Eriogonum parvifolium*), which is the only host plant for the larvae of the rare El Segundo blue butterfly, is found there, and the buckwheat population is isolated from other populations. Sensitive plant species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Descriptions and general locations of the each plant community present within the CRA are given below.

Coastal Strand Vegetation: A community that occurs in the loose sand just above the high tide line of the beach. This community is characterized by a low species diversity because few plants can tolerate the harsh conditions on this dry, sandy, saline soil with buffeting by high winds, salt spray and high summer temperatures.

Corresponding MCV communities:

- *Abronia latifolia* – *Ambrosia chamissonis* (dune mat) Herbaceous Alliance
- *Cakile* (*edentula*, *maritima*) (sea rocket stands) Semi-Natural Provisional Herbaceous Alliance

Coastal Bluff and Dune Scrub: A remnant of the El Segundo Dune system that occupied coastal areas along much of the Santa Monica Bay. In the CRA, the coastal bluff and dune scrub is found in a strip along the immediate coastal sections of the CRA. The system consists of fine, sandy soil that dries rapidly. Plants typical of the dune scrub include dune buckwheat (*Eriogonum parvifolium*), rattle pod (*Astragalus trichopodus* var. *lonchus*), bladderpod (*Peritoma arborea*), deer weed (*Acmispon glaber*), sawtooth goldenbush (*Hazardia squarrosa*), and California sunflower (*Helianthus californicus*).

Corresponding MCV communities:

- *Baccharis pilularis* (coyote brush scrub) Shrubland Alliance
- *Lupinus arboreus* (yellow bush lupine scrub) Shrubland Alliance and Semi-Natural Shrubland Stands
- *Lupinus chamissonis*-*Ericameria ericoides* (silver dune lupine–mock heather scrub) Shrubland Alliance

Coastal Sage Chaparral Scrub: Characterized by the summer drought deciduous vegetation found near the Southern California coast of low, mostly soft-woody shrubs with bare ground underneath and between shrubs. This community is dominated by California sagebrush, California buckwheat (*Eriogonum fasciculatum*), black sage (*Salvia mellifera*), purple sage (*Salvia leucophylla*), and California brittle bush (*Encelia californica*) and intergrades with the coastal bluff and dune scrub near the shore and gradually becomes more evident near the crest of the site

DRAFT

Corresponding MCV communities:

- *Artemisia californica* (California sagebrush scrub) Shrubland Alliance
- *Artemisia californica-Salvia mellifera* (California sagebrush-black sage scrub) Shrubland Alliance
- *Artemisia californica-Eriogonum fasciculatum* (California sagebrush-California buckwheat scrub) Shrubland Alliance
- *Encelia californica* (California brittle bush scrub) Shrubland Alliance
- *Salvia mellifera* (black sage scrub) Shrubland Alliance
- *Eriogonum fasciculatum* (California buckwheat scrub) Shrubland Alliance
- *Hazardia squarrosa* (sawtooth goldenbush scrub) Shrubland Alliance
- *Lotus scoparius* [*Acmispon glaber*] (deer weed scrub) Shrubland Alliance
- *Lupinus albifrons* (silver bush lupine scrub) Shrubland Alliance

Wildlife

An assessment of wildlife is made difficult due to the lack of data but the animal populations within the CRA probably reflect a somewhat lower diversity and abundance for the habitat types present for several reasons, including: the small area of the dune habitat, the homogeneity of the topography and habitat, influences of edge effect from developments to the north and south, the ocean to the west and the impact of being under the flight paths of aircraft departing from LAX to the east.

One key occupant of the area is the El Segundo blue butterfly, which is found here and also in a very few other locations to the south in remnants of the former dune complex. The El Segundo blue butterfly co-occurs with the only known plant species that supports its complete lifecycle, the dune buckwheat (*Eriogonum parvifolium*). The adult butterfly uses the flowers as a major source of nectar, and the larvae feed only on the flowers and seeds.

Amphibian populations are generally scarce in beachfront communities and no riparian habitat is available within the CRA. Many essential reptilian habitat characteristics, such as open habitats that allow free movement and high visibility and small mammal burrows for cover and escape from predators and extreme weather, are present within the CRA. The scrubland habitats in the CRA probably provide foraging and cover habitat for year-round resident and seasonal resident song birds. In addition, the CRA provides some raptor foraging habitat, so they can be expected visitors. Mammal populations probably reflect the generally disturbed environs influencing this CRA. Small mammals are expected to be uneven in their diversity, with more adaptive species and introduced European species occurring in higher numbers compared to others. Medium sized mammal populations are expected to exhibit the same characteristics. Resident large mammals are absent. Sensitive wildlife species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Wildlife Movement

The CRA is on the Pacific Flyway migration route used by many birds seasonally. The dune habitat

probably does not play an important role as a stopover because of its limited resources, due its small size; however, it will attract some birds because it is one of the few undeveloped places along the coast. The area does not fall within any identified terrestrial movement routes for wildlife.

Sensitive Biological Resources

Sensitive biological resources are habitats or individual species that have special recognition by federal, state, or local conservation agencies and organizations as endangered, threatened, and/or rare. This is due to the species' declining or limited population sizes, which usually results from habitat loss. Watch lists of such resources are maintained by the California Department of Fish and Wildlife (CDFW), the United States Fish and Wildlife Service (USFWS), and special groups, such as the California Native Plant Society (CNPS). The following sections indicate the habitats as well as plant and animal species present, or potentially present within the CRA, which have been accorded special recognition. When species are federally-listed as endangered or threatened, they often have federally-designated, geographically-specific "critical habitat areas." Critical habitat is a specific geographic area that contains features essential for the conservation of a threatened or endangered species and that may require special management and protection. Critical habitat may include an area that is not currently occupied by the species but that will be needed for its recovery. An area is designated as "critical habitat" after a proposed Federal regulation is published by USFWS in the Federal Register and public comments on the proposal are received and considered. However, for species listed previous to 1978, critical habitat areas rarely are designated officially. Sometimes such areas are called "essential habitat," which would be the case here for the EI Segundo blue butterfly.

Sensitive Plant Communities and Habitats

The CRA supports several habitat types considered sensitive by resource agencies. These are inventoried by California Department of Fish and Wildlife (CDFW) in the California Natural Diversity Database (CNDDB), or have been assigned a conservation status rank of 1 – 3 at the global or subnational scale. Conservation status ranks are assigned by CDFW, following methodology developed by NatureServe (<http://www.natureserve.org/conservation-tools/standards-methods/conservation-status-assessment>). Within this SEA, these include dune mat, silver dune lupine–mock heather scrub, California brittle bush scrub, and sawtooth goldenbush scrub. These communities, or closely related designations, are considered high priority communities by the CDFW, which indicates that they are experiencing a decline throughout their range. The array and composition of these communities has been discussed in the Vegetation section.

Sensitive Plant Species

The statuses of rare plants are hierarchically categorized by the CNPS using a rank and decimal system. The initial category level of Rare Plant Rank is indicated by the ranks 1A (presumed extinct), 1B (rare or endangered in California and elsewhere), 2A (presumed extinct in California but extant elsewhere), 2B (rare or endangered in California but more common elsewhere), 3 (more information needed, a review list), and 4 (limited distribution). In cases where the CNPS has further identified the specific threat to the species, a decimal or Threat Code is added: .1 (seriously endangered in California), .2 (fairly endangered in California), or .3 (not very endangered in California).

The following special-status plant taxa have been reported or have the potential to occur within the CRA, based on known habitat requirements and geographic range information:

- *Aphanisma* (*Aphanisma blitoides*) RPR 1B.1

DRAFT

- Coastal dunes milk-vetch (*Astragalus tener* var. *titi*) FE, SE, RPR 1B.1
- Orcutt's pincushion (*Chaenactis glabriuscula* var. *orcuttiana*) RPR 1B.1
- Beach spectaclepod (*Dithyrea maritima*) ST, RPR 1B.1
- Coulter's goldfields (*Lasthenia glabrata* ssp. *coulteri*) RPR 1B.1
- Brand's star phacelia (*Phacelia stellaris*) FC, RPR 1B.1

Sensitive Animal Species

The following special-status animal species are reported or are likely to be present within the CRA based on habitat requirements and known range attributes:

- Belkin's dune tabanid fly (*Brennania belkini*) CDFW Special Animals List
- Busck's gallmoth (*Carolella busckana*) CDFW Special Animals List
- Sandy beach tiger beetle (*Cicindela hirticollis gravida*) CDFW Special Animals List
- Senile tiger beetle (*Cicindela senilis frosti*) CDFW Special Animals List
- Globose dune beetle (*Coelus globosus*) CDFW Special Animals List
- Henne's eucosman moth (*Eucosma hennei*) CDFW Special Animals List
- El Segundo blue butterfly (*Euphilotes battoides allyni*) FE, Xerces: Critical
- Lange's El Segundo Dune weevil (*Onychobaris langei*) CDFW Special Animals List
- El Segundo flower-loving fly (*Rhaphiomidas terminatus terminatus*) CDFW Special Animals List
- Silvery legless lizard (*Anniella pulchra pulchra*) FSS, SSC
- Coastal whiptail (*Aspidoscelis tigris stejnegeri*) CDFW Special Animals List
- San Bernardino ringneck snake (*Diadophis punctatus modestus*) FSS
- Coast horned lizard (*Phrynosoma blainvillii*) BLMS, FSS, SSC
- Western snowy plover (*Charadrius alexandrinus nivosus*) FT, BCC, SSC, ABC, AWL, USBC
- California black rail (*Laterallus jamaicensis coturniculus*) BCC, ST, CDFW Fully Protected, USBC, AWL, ABC
- California least tern (*Sternula antillarum browni*) FE, SE, CDFW Fully Protected, USBC, ABC
- Pacific pocket mouse (*Perognathus longimembris pacificus*) FE, SSC

Regional Biological Value

The CRA meets several SEA designation criteria incorporating regional biological values. Each criterion and how it is met is described below.

Criteria Analysis of the El Segundo Dunes CRA

Criterion	Status	Justification
A) The habitat of core populations of endangered or threatened plant or animal species.	Met	The El Segundo Dunes is one of four known localities in the world where one can find the endangered El Segundo blue butterfly. All four areas are remnants of a once continuous three-mile system of seacoast dunes in the County. The blue's habitat is the coastal dune buckwheat, on which the butterfly lays its eggs and the larvae feed.
B) On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The southern dune scrub vegetation has become very uncommon in the Southern California region due to the conversion of the habitat to seacoast dwellings, hotels, and other development. The vegetation thrives on sandy, well-drained soils, and grades into coastal sage scrub on the inland side. Many plants and invertebrates are restricted to this environment and are not found elsewhere. This small piece of dune habitat is extremely valuable as the final example of a community that was once more common along the Southern California coast.
C) Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution	Met	The southern dune scrub vegetation found here on the three remnant localities of the El Segundo Dunes does not occur anywhere else in the County.
D) Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	A number of organisms call the sandy, well-drained soils of the El Segundo Dunes "home," and are restricted to this coastal environment. Like the El Segundo blue butterfly, their entire life cycles are here in the dunes.
E) Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	As a result of the uniqueness of this community and of the specializations of the organisms found here, the dunes have been closely scrutinized by biologists of many disciplines.
F) Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Not Met	Much of the original threemiles of dunes have been removed by development or disturbed. Scattered on the site are radio towers, roads, oil extraction equipment, and a reservoir. The vegetation is much impacted by human use, but can be restored. The designated area is the best remaining part of the original El Segundo Dunes.

DRAFT

In conclusion, the area is a CRA because it contains A) core habitat for the federally-endangered El Segundo blue butterfly; B-C) beach dunes, a rare community both regionally and within the County; D) a habitat that is rare and necessary for the life cycle of the El Segundo blue butterfly and other dune insects and fauna; and E) the geographic limit of El Segundo Dunes, a rare habitat much studied by biologists.

Malibu Coastline CRA

Location

General

The Malibu Coastline CRA is located in the shoreline and offshore coastal area of Malibu, which is adjacent to the Santa Monica Mountains. The CRA supports significant areas of aquatic plant and other subtidal communities, which provide habitat for a variety of fishes, birds, marine mammals, and other wildlife. Rocky outcrops intermixed with sandy spaces are found to a depth of 600 feet, and the nearshore area down to about 100 feet depth is considered the most productive and dynamic of all the marine communities outside the tropics. All of the many offshore rocks within 12 nautical miles of the coast are part of the California Coastal National Monument that is managed by the Bureau of Land Management in the U.S. Department of the Interior.

The site is located in the Triunfo Pass, Point Dume, Malibu and Topanga Quadrangles of the United States Geological Survey (USGS) 7.5 Minute Map Series (USGS, 1964).

General Boundary and Resources Description

The CRA boundary encompasses parts of the shoreline and about a 0.8 mile offshore from Malibu. The CRA stretches between the Ventura-Los Angeles County line at Leo Carillo State Beach, all the way east to Topanga State Beach. The CRA is generally adjacent to the Santa Monica Mountains SEA and Point Dume CRA. Parts of the shore are included along many of the state beaches. The sandy beaches along this CRA are the least disturbed beaches of the County, some of them completely natural. This is the remnant of the typical rock and sand shoreline that once occurred all along the coast of Southern California. From Mugu Lagoon to Latigo Point (the County portion starting at the Ventura-Los Angeles County line), is an Area of Special Biological Significance (ASBS), which is a marine area designated by the State Water Resources Control Board as having exceptionally good water quality and natural community features. Populated and disturbed areas along the shore are largely in the City of Malibu and excluded from the CRA.

This area is a relatively undisturbed coastal region where the upwelling of nutrient-rich waters and a variety of habitats support highly productive and extremely diverse marine communities. The area possesses some of the best kelp bed habitat south of Santa Barbara, and supports the only remaining natural kelp beds off the mainland coast of the County. These kelp bed areas may be one hundred times more productive than adjacent sand bottom communities, and they provide refuge, food, and nursery grounds for thousands of species.

Rocky outcrops alternate with sandy stretches along this coastline and subtidally, outcrops are found to a depth of 600 feet. The stability of the substrate and the variety of exposures provide microhabitats for a great number of organisms. Characteristically, rocky shorelines from the lower intertidal zone to about 100 foot depth can be the most biologically active areas in the world. The adjacent Point Dume CRA is one of the few places that rocky intertidal habitat occurs between Palos Verdes Peninsula and Point Mugu in Ventura County. The rocky tidepools off Point Dume, Big Rock Beach, and the

DRAFT

promontory where Topanga Canyon Creek enters the ocean are some of the very best remaining rocky intertidal habitat in the County, and these are within the CRA. The Marine Life Protection Act designated protected area in Malibu to be the Point Dume State Marine Conservation Area (SMCA) between El Matador State Beach west of Point Dume to the west side of Point Dume, which is approximately three miles of coastline, with the conservation area extending in due north-south lines from mean high tide to about three nautical miles offshore. In the SMCA, fishing is restricted to certain species. From the west side of Point Dume to Paradise Cove on the east side is designated as the Point Dume State Marine Reserve (SMR), which has no fishing or other take allowed, and extends in due north-south lines (adjacent to the SMCA on the west) from the intertidal to offshore, which is about three nautical miles.

This coastline possesses the only undisturbed sandy beaches that remain in the County. Although very dynamic in physical stability, there is a biological, subtidal, sand-bottom community that has a great diversity of resident organisms and invertebrates ranging from bacteria through jellyfish, mollusks, and echinoderms, such as seastars, to fishes that habitually choose the sand-bottom substrate. An important micro-community of decomposers is present. Sandy beaches with their diverse invertebrate communities of interstitial organisms provide feeding areas for many bird species. In addition, the soft substrate offers a repository for eggs and nursery grounds for many species, including the grunion (*Leuresthes tenuis*) that spawns and deposits its eggs in the highest intertidal areas of the sand. This shoreline remains in essentially a native state as a remnant of what once was typical of rock and sand shoreline in Southern California. Artificial modifications have been limited to small local areas. West of Point Dume, some minor pollution does occur, but the kelp is healthy. East of Point Dume there is minor to moderate pollution, and kelp does not grow below 35 feet.

The CRA has critical habitat for the federally-threatened western snowy plover (*Charadrius alexandrinus nivosus*) along Zuma Beach and the northwest sandy area along Point Dume. This will eventually extend along the entire west side of the headland. The western snowy plover will also have critical habitat on the seaside of the barrier berm of the Malibu Lagoon. The snowy plover has over 300 individuals using the beaches of the County for winter roosts, and there is potential for some to over-summer and breed in the County after a hiatus of many decades (Ryan Ecological Consulting, 2010).

Vegetation

The terrestrial component of this marine oriented CRA is limited to the narrow stretch of beach just above the high tide line. Sensitive plant species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Coastal Strand Vegetation: A community that occurs in the loose sand just above the high tide line of the beach. This community is characterized by a low species diversity because few plants can tolerate the harsh conditions on this dry, sandy, saline soil of high winds, salt spray and high summer temperatures.

Corresponding MCV communities:

- *Abronia latifolia* – *Ambrosia* spp. (dune mat) Herbaceous Alliance
- *Cakile* (*edentula*, *maritima*) (sea rocket stands) Semi-Natural Provisional Herbaceous Alliance

Marine Resources

The Malibu coastline is a relatively undisturbed coastal region where the upwelling of nutrient-rich waters and a variety of habitats support highly productive and extremely diverse marine communities. The area possesses some of the best kelp bed habitat south of Santa Barbara, and the only remaining natural kelp beds on the mainline coast of the County. Rocky outcrops alternate with sandy stretches along this coastline and are found to a depth of 600 feet. This coastline also possesses the only undisturbed sandy beaches remaining in the County.

Surfgrass and Eelgrass Habitats: Characterized by two flowering plants that resemble grass and form dense beds on different substrates and in different conditions. Both types form highly productive habitats for unique assemblages of marine species. NOAA Fisheries and the CDFW consider surfgrass and eelgrass as valuable habitats that should be protected. Surfgrass occurs from Mean Lower Low Water (MLLW) to about 25 feet depth on rocky substrate. Locations known to support substantial surfgrass beds include rocky points and rocky subtidal areas of canyons all along the northern border of the CRA. Eelgrass grows on soft substrate and typically occurs in bays. A form of eelgrass (thought to be *Zostera pacifica*) is found on open coast sand bottom. Eelgrass is an important habitat that supports a community of diverse species from sessile to mobile invertebrates and certain fishes, such as pipefish. A subtidal eelgrass bed is in the lee of Lechuza Point between about 25 and 45 feet depth, and subtidal eelgrass may still occur off Los Alisos Canyon.

Community dominant plant:

- *Phyllospadix scouleri*, or Scouler's surf grass and *P. torrey* or Torrey's surfgrass
- *Zosterapacifica* (eelgrass)

Intertidal Zone Habitats: Consist of a variety of coastal habitats periodically covered and uncovered by waves and tides. The rocky shores support a rich assortment of plants and animals including green, brown, and red algae. A wide variety of sessile invertebrates compete for space with the plants in this habitat. Mobile invertebrates, such as crabs and snails, can be found in great abundance. Fishes are limited to tidepools, although grunion do spawn and deposit their eggs in the high tide wash areas of sandy beaches. The sandy beaches have a diverse community of interstitial invertebrates. Shorebirds actively forage during low tide in all kinds of shoreline habitats. The federally-endangered black abalone (*Haliotus cracherodii* FE) was once abundant in this habitat all along the rocky coastlines of California. Whether it still persists in the Malibu coastline area is unknown, since its current populations are reduced and much more dispersed than before.

Nearshore Subtidal Habitat: Includes those marine habitats ranging from the lower level of the intertidal zone to 99 feet. This region supports a variety of assemblages of invertebrates and fishes, and in the CRA, this habitat is frequently dominated by giant kelp. Rocky areas have a diverse community of algae (in depths of sufficient light penetration), sessile and mobile invertebrates, and fishes. Subtidal areas are even more diverse than the intertidal areas, and this great variety can be appreciated by the examples of tidepools. There are a variety of subtidal sand-adapted organisms ranging from fishes to seastars, to many kinds of jellyfish, mollusks, and other invertebrates.

Kelp Forest Habitat: Giant kelp beds are located in many places along the CRA to a depth of approximately 99 feet in the ocean. The kelp beds are part of a productive habitat that provide food, attachment sites and shelter for invertebrates and fishes. Giant kelp, the dominant alga of this community, is the fastest lengthening organism known, and it thrives in nutrient-enriched waters of

DRAFT

upwelling. It has been “clocked” at two feet per day extension of its stipe and blades. The kelp beds are an important nursery habitat and recruitment area for juvenile fishes and invertebrates. The National Oceanographic and Atmospheric Administration (NOAA) Fisheries as well as the CDFW consider kelp beds as sensitive, and lush kelp beds such as those from the Ventura-Los Angeles County line to Malibu Point are designated as ESHA (Environmentally Sensitive Habitat Areas, Malibu Local Coastal Plan, 1986).

Community dominant plant:

- *Macrocystis pyrifera* (Giant kelp)

Wildlife

The terrestrial and aerial wildlife found in the CRA is dependent on the two basic regimes found there: marine and shoreline terrestrial. The shoreline beaches and rocky intertidal are home to or visited by a wide variety of shorebirds, migrating birds and marine life. The sandy beaches with their shifting sands present an unstable substrate on which organisms can establish themselves, and their resident wildlife is a set of small specialists that live in the sand interstices. An important microcommunity of decomposers exists, which feed on the materials washed up by the waves.

The Malibu Coastline cliffs, bluffs, offshore rocks and beaches offer many undisturbed habitats for roosting, feeding and nesting by numerous kinds of shore— and seabirds. Sandy beaches provide feeding areas for many species. In addition, the soft substrate offers a repository for eggs and nursery grounds for many species.

The marine habitat has the greatest diversity of wildlife, with representatives from nearly all the phyla (major groupings of animals such as jellyfish, mollusks, echinoderms, etc.) and all parts of the food web (trophic levels) for several communities. Many of the marine phyla do not have terrestrial representatives. Some of the vertebrates, such as gray whales, and the plankton use the area as a migratory corridor. The major vegetation communities, each with its own great diversity and all trophic levels, are the benthic algae of rocky substrates, the kelp beds based on giant kelp holding onto rocky subtidal substrate and extending into the water column, and the planktonic, based on photosynthesizers that are all microscopic.

Sensitive wildlife species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section of this document.

Wildlife Movement

The CRA is on the Pacific Flyway migration route used by many birds seasonally. This shoreline plays an important role as a stopover because of its extensive undisturbed marine resources. The offshore major part of the CRA is on the annual migration route of the California gray whale (*Eschrichtius robustus*), a CITES Appendix I animal (Convention on International Trade in Endangered Species, endangered status, no trade or harvest except by strictly controlled export and import permits issued in countries that are party to the convention and have legislation of adoption, which includes the U.S.). The gray whale migrates close to shore on its transit during the winter months from the Bering Sea to lagoons of Baja California. This is migration to the Baja lagoons for calving and breeding. The whales generally return north offshore. The area may well be a migration corridor for other marine animals, as this is part of the great Southern California Bight, which has an eddy circling counterclockwise to the north off the southward flow of the California Current. This flow brings marine organisms (fishes, invertebrates, and plants) that may have major population concentrations in the south to this part of

their ocean habitat. Most of these organisms have a planktonic stage that is absolutely dependent on the current system to maintain their populations and distribution. Pollution of the ocean waters by development on land will have significant effects that may not be appreciated immediately because of the cryptic nature of subtidal animals and plants.

Sensitive Biological Resources

Sensitive biological resources are habitats or individual species that have special recognition by federal, state, or local conservation agencies and organizations as endangered, threatened, and/or rare. This is due to the species' declining or limited population sizes, which usually results from habitat loss. Watch lists of such resources are maintained by the California Department of Fish and Wildlife (CDFW), the United States Fish and Wildlife Service (USFWS), and special groups, such as the California Native Plant Society (CNPS). The following sections indicate the habitats as well as plant and animal species present, or potentially present within the CRA, which have been accorded special recognition. When species are federally-listed as endangered or threatened, they often have federally-designated, geographically-specific "critical habitat areas." Critical habitat is a specific geographic area that contains features essential for the conservation of a threatened or endangered species and that may require special management and protection. Critical habitat may include an area that is not currently occupied by the species but that will be needed for its recovery. An area is designated as "critical habitat" after a proposed Federal regulation is published by USFWS in the Federal Register and public comments on the proposal are received and considered. There are two critical habitat areas for the western snowy plover in this CRA.

Sensitive Plant Communities and Habitats

The CRA supports no terrestrial habitat types considered sensitive by resource agencies, but the California Department of Fish and Wildlife (CDFW) does not list intertidal or subtidal plants in its California Natural Diversity Data Base (CNDDDB, 2011). NOAA Fisheries (National Oceanographic and Atmospheric Administration) as well as the CDFW do consider kelp beds as sensitive, and lush kelp beds such as those from the Ventura-Los Angeles County line to Malibu Point are designated as ESHAs by the County's Malibu Local Coastal Plan, 1986.

Kelp beds disappeared off of Palos Verdes Peninsula during the height of pollution by the White Point outfall in the mid 20th century. The Palos Verdes Peninsula's kelp beds are subsequently returning after reduction of pollution, some prodigious planting efforts, and incursion of the motile zoospores from areas like the Malibu coastline. The kelp beds are sensitive to effects like pollution and excessive sedimentation from development. They are one of the most productive communities of the world, and should be treated with care. Any development plan that might impact them, even with indirect effects, deserves scrutiny.

Sensitive Plant Species

NOAA Fisheries as well as the CDFW consider kelp beds as sensitive, and lush kelp beds, such as those from the Ventura-Los Angeles County line to Malibu Point, are designated as ESHA by the Malibu Local Coastal Plan, 1986. The principle structural component of Malibu offshore kelp beds is the giant kelp (*Macrocystis pyrifera*). NOAA Fisheries and the CDFW consider surfgrass a valuable habitat that should be protected. The principle structural component of surfgrass beds is Scouler's surf grass (*Phyllospadix scouleri*) and Torrey's surfgrass (*P. torreyi*).

Sensitive Animal Species

DRAFT

The western snowy plover (*Charadrius alexandrinus nivosus* [FT, CSC]), which feeds and resides in the wrack line areas, has designated critical habitat on Zuma Beach from Trancas Canyon to the northwestern side of Point Dume. This is proposed to expand to Point Dume. Critical habitat is also proposed to include Malibu Beach from Malibu Point to an area east of the pier, which is the seaward side of Malibu Lagoon. The Malibu Lagoon is part of the Santa Monica Mountains SEA, and the CRA is contiguous with the Santa Monica Mountains SEA at the barrier berm of the Lagoon.

The southern steelhead (*Oncorhynchus mykiss irideus* [FE, CSC]) lives in the oceanic and coastal waters for most of its life and uses the coastal streams for breeding and the first year of its young fish's lives. After one to two years in fresh water, the young fish change to smolts and make their run to the ocean, where they spend the majority of their lives. The lower Arroyo Sequit and its West Fork is designated critical habitat for the southern steelhead. This area has naturally occurring spawning beds and young fish habitat. The coastal ocean waters are within this CRA, and the linkage paths from the ocean to the coastal streams are within this CRA. The southern steelhead is known to currently use the Arroyo Sequit, Malibu Creek, and Topanga Canyon. Historically the steelhead was known from Solstice and Zuma canyons, and was probable in all the major drainages, which once had perennial water and extended to the shore in the rainy season.

The California gray whale (*Eschrichtius robustus* CITES Appendix I) uses this CRA during its calving-breeding migration cycle. The entire order of cetaceae (whales and beaked dolphins) is considered CITES Appendix I by the Australian rating system, the most stringent adoption.

The black abalone (*Haliotis cracherodii*) is a federally-endangered species and critically endangered species of the International Union for Conservation of Nature-Red List (IUCN). At one time, the black abalone was plentiful in the rocky intertidal and nearshore rocky subtidal areas in the CRA, down to about 20 feet depth. Whether it still exists in the CRA is unknown, because its current occurrences are widely scattered and much reduced by overfishing and wasting disease.

- Black abalone (*Haliotis cracherodii*) FE
- Southern steelhead (*Oncorhynchus mykiss irideus*) FE, CSC
- Tidewater goby (*Eucyclogobius newberryi*) FE, SSC
- Western snowy plover (*Charadrius alexandrinus nivosus*) FT, CSC
- California gray whale (*Eschrichtius robustus*) CITES Appendix I

Regional Biological Value

The CRA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE MALIBU COASTLINE CRA

	Criterion	Status	Justification
	The habitat of core populations		California gray whale, tidewater goby, southern steelhead, western

DRAFT

	Criterion	Status	Justification
A)	of endangered or threatened plant or animal species.	Met	snowy plover, and black abalone, all spend critical periods of their life cycles in this CRA. Other CITES-listed marine mammals also use this CRA.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	This area is a relatively undisturbed coastal region where the upwelling of nutrient-rich waters provides for highly productive and extremely diverse marine communities. The area possesses some of the best kelp bed habitat south of Santa Barbara and is recognized as ESHA by NOAA and CDFW. The Malibu coastline section from the Ventura-Los Angeles County line boundary, southeast to Latigo Point is recognized as one of the principal natural areas in the (coastal) State Water Quality Protection Area Program. It is area ASBS-24 (Area of Special Biological Significance). The CRA has critical habitat for the southern steelhead, tidewater goby, and the western snowy plover.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	This relatively undisturbed coastal region possesses some of the best kelp bed habitat south of Santa Barbara and supports the only remaining natural kelp beds off the mainland coast of the County. The kelp beds are recognized as ESHA by NOAA, CDFW, and the Malibu Local Coastal Program. It has critical habitat for the southern steelhead, tidewater goby, and the western snowy plover
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	This area protects the entry point of two of three of the (endangered) southern steelhead spawning streams in the County and provides connective area for the endangered tidewater goby, which breeds in the brackish-water areas of the streams of the Santa Monica Mountains. Kelpbeds are the macroforest of the ocean, and the habitat and breeding and/or spawning ground for many marine animals. The sandy beaches provide feeding areas for many bird species, and the soft substrate offers a repository for eggs and nursery grounds for many species. The migration of marine species occurs in this area for the California gray whale and for innumerable marine species whose plankton ride the ocean currents.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	As the main kelp bed of the County, this area is of great interest to marine scientists. It is also an edge kelp forest, separated by the long stretch of beaches of the Santa Monica Bay from the kelp beds of the Palos Verdes Peninsular area.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	This shoreline remains in essentially a native state as a remnant of what once was typical of rock and adjacent sand shoreline in Southern California. The coastline possesses the only complete, undisturbed sandy beaches remaining in the County. An important microcommunity of decomposers is present. Artificial modifications have been limited to small local areas. West of Point Dume, there is minor pollution; east of Point Dume, there is minor to moderate pollution, and kelp does not grow below 35 feet.

DRAFT

In conclusion, the area is a CRA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County, or regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Palos Verdes Peninsula and Coastline CRA

For a description of the resources in the Palos Verdes Peninsula and Coastline CRA, please refer to the description for the Palos Verdes Peninsula and Coastline SEA above.

Point Dume CRA

Location

General

The Point Dume Coastal Resource Area (CRA) is located on the Malibu coastline and forms the northern end of Santa Monica Bay. Zuma Beach lies to the immediate northwest. Paradise Cove is just east. The CRA is important as a headland on the Pacific Flyway, for its nearby marine and its terrestrial habitats, and for critical habitat of the western snowy plover.

The CRA is located in the Point Dume United States Geological Survey (USGS) 7.5' California Quadrangle.

General Boundary and Resources Description

This headland extends into Santa Monica Bay more than a mile beyond the rest of the Malibu coast, and it is located within the Pacific Flyway. The CRA consists of the rocky coastal headland edge and six fingers extending into the interior canyons on the headland. Point Dume is a significant headland used during migration times on the Pacific Flyway. Point Dume is the southernmost habitat for coastal cliff flora such as the giant coreopsis (*Leptosyne gigantea*) and sea-lettuce (*Dudleya caespitosa*). The CRA is contiguous with the Malibu Coastline CRA, which has notable tidepools with rich marine habitat on the eastern side of Point Dume. Southern California is known for its extensive sandy beaches, so Point Dume, which is one of very few places with rocky intertidal habitat between Palos Verdes Peninsula and the middle coastline of Ventura County, is a rare habitat in the County. The limited public access, the coastal strand vegetation, and the canyon slopes covered by Venturan coastal sage scrub, protect the resource values of this headland, both terrestrial and marine. The terrestrial and marine habitats combined make Point Dume an exceptional and unique CRA. The Marine Life Protection Act designated protected area in Malibu to be the Point Dume State Marine Conservation Area (SMCA) between El Matador State Beach west of Point Dume to the west side of Point Dume (east end of Zuma Beach), approximately a three-mile coastline. In the Conservation area, take is limited to certain designated fish and squid species for a distance of three nautical miles proceeding due north-south out from the shore. Adjacent, from the west side of Point Dume to the east side of Point Dume at Paradise Cove is designated as the Point Dume State Marine Reserve (SMR), with no take extending to three nautical miles offshore. This includes very fine tidepools on the east side.

DRAFT

Point Dume is one of two remaining areas in the County where a diverse and healthy mixture of terrestrial and marine habitats can be found in close association. Point Dume incorporates the terrestrial habitats: the interior canyons, the unprotected rocky shore with numerous small caves, and the sandy beach pockets; and the Malibu Coastline CRA has the marine habitats that are adjacent: outlying reefs, rocks, kelp beds, and tide pools. The tide pools off of Point Dume are some of the very best remaining in the County. Due to strong upwelling along the coast bringing in nutrient-rich waters, the nearby marine habitat is characterized by highly diverse and productive marine communities. This relative healthiness of both the terrestrial and marine habitats is largely due to limited public access, which has protected the fragile marine and shoreline ecosystems.

Coastal strand vegetation is found on sandy beaches below bluffs that rise 100 to 200 feet above the coast. Giant coreopsis (*Leptosyne gigantea*) and sea-lettuce (*Dudleya caespitosa*) are found in these communities at the southern limit of their range. Several small drainages cut through the bluffs and extend over a mile inland. The slopes are covered by Venturan coastal sage scrub. The value of these communities is increased by the unique geographic position of Point Dume. The area is an important resting and jumping-off point for migratory birds. Without the remaining terrestrial habitats, this refuge would be lost.

The CRA will have critical habitat for the federally-threatened western snowy plover (*Charadrius alexandrinus nivosus*). Currently this critical habitat goes eastward from Trancas Beach, along Zuma Beach, and extends to the northwest sandy area along Point Dume. The plovers use the west sandy habitat of the Point Dume headland, and the critical habitat will eventually extend along the entire west side of the headland into the CRA. (The western snowy plover will also have critical habitat on the seaside of the barrier berm of the Malibu Lagoon.) The snowy plover has over 300 individuals using the beaches of the County for winter roosts, and there is potential for some to over-summer and breed in the County after a hiatus of many decades (Ryan Ecological Consulting, 2010).

Vegetation

Vegetation within the CRA is comprised of several community types. Coastal strand vegetation occurs along the immediate shore and on the cliffs and bluffs, overlooking the ocean coastal dune, and bluff scrub also can be found. In the ravines and upper slopes examples of Venturan coastal sage scrub intergrade with maritime succulent scrub. This is a type of coastal sage scrub, defined by the unusual giant coreopsis, which is found in some of the less disturbed areas of the CRA. Sensitive plant species and plant communities occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Coastal Strand Vegetation: A community that occurs in the loose sand just above the high-tide line of the coast. This community is characterized by low species diversity because few plants can tolerate the harsh conditions on this dry, sandy, saline soil of high winds, salt spray and high summer temperatures.

Corresponding MCV communities:

- *Abronia latifolia* – *Ambrosia* spp. (dune mat) Herbaceous Alliance
- *Cakile* (*edentula*, *maritima*) (sea rocket stands) Semi-Natural Provisional Herbaceous Alliance

Coastal Bluff and Dune Scrub: Found on the cliffs and bluffs overlooking the ocean. This community formerly was found in many places along the coast, but is now restricted to a few locations. The system

DRAFT

consists of fine, sandy soil that dries rapidly. Plants typical of the dune scrub include dune buckwheat, rattle-pod, bladderpod, deer weed, sawtooth goldenbush, and California sunflower.

Corresponding MCV communities:

- *Baccharis pilularis* (coyote brush scrub) Shrubland Alliance
- *Lupinus arboreus* (yellow bush lupine scrub) Shrubland Alliance and Semi-Natural Shrubland Stands
- *Lupinus chamissonis-Ericameria ericoides* (silver dune lupine–mock heather scrub) Shrubland Alliance

Venturan Coastal Sage Scrub: Characterized by the summer drought-deciduous vegetation found near the Southern California coast south of Ventura. It has low, mostly soft-woody shrubs with bare ground underneath and between shrubs. This community is dominated by California sagebrush, California buckwheat, black sage, purple sage, and California encelia.

Corresponding MCV communities:

- *Artemisia californica* (California sagebrush scrub) Shrubland Alliance
- *Artemisia californica-Salvia mellifera* (California sagebrush-black sage scrub) Shrubland Alliance
- *Artemisia californica-Eriogonum fasciculatum* (California sagebrush-California buckwheat scrub) Shrubland Alliance
- *Encelia californica* (California brittle bush scrub) Shrubland Alliance
- *Salvia apiana* (white sage scrub) Shrubland Alliance
- *Salvia leucophylla* (purple sage scrub) Shrubland Alliance
- *Salvia mellifera* (black sage scrub) Shrubland Alliance
- *Eriogonum fasciculatum* (California buckwheat scrub) Shrubland Alliance
- *Hazardia squarrosa* (sawtooth goldenbush scrub) Shrubland Alliance
- *Lotus scoparius* ([*Acmispon glaber*] deerweed scrub) Shrubland Alliance
- *Lupinus albifrons* (silver-bush lupine scrub) Shrubland Alliance
- *Malacothamnus fasciculatus* (bush mallow scrub) Shrubland Alliance

Maritime Succulent Scrub: A form of coastal sage scrub with many members of that community, as well as one or more endemic succulent evergreen species. Examples of this can be found throughout the CRA.

Corresponding MCV communities:

DRAFT

- *Leptosyne gigantea* (giant coreopsis scrub) Shrubland Alliance
- *Opuntia littoralis* (coast prickly-pear scrub) Shrubland Alliance
- *Lycium californicum* (desert thorn scrub) Provisional Shrubland Alliance

Marine Resources

The Malibu coastline is a relatively undisturbed coastal region where upwelling of nutrient-rich waters and a variety of habitats support highly productive and extremely diverse marine communities. The terrestrial habitat (here the Point Dume headland) and the marine habitats are integrally tied to one another. Disruption in one will impact the other. The marine habitats feed the seabirds that use the headland, and marine organisms, such as the grunion (fish, *Leuresthes tenuis*) use the sandy shore for spawning. The headland protects the nearshore habitats and controls and filters runoff so that the offshore habitats remain clean and have the terrestrial input of nutrients for which they are adapted. The nearby areas in the Malibu Coastline CRA possesses some of the best kelp bed habitat south of Santa Barbara, and the only remaining natural kelp beds on the mainland coast of the County. Rocky outcrops alternate with sandy stretches along this coastline and are found to a depth of 600 feet. The Point Dume headland typifies the alternating sand and rock on several scales. The Malibu coastline possesses the only complete, undisturbed sandy beaches remaining in the County. The west side of the headland has critical habitat for the snowy plover.

Intertidal Zone Habitats: Consist of a variety of coastal habitats that are periodically covered and uncovered by waves and tides. The rocky shores support a rich assortment of plants and animals including green, brown, and red algae. A wide variety of sessile invertebrates compete for space with the plants in this habitat. Mobile invertebrates, such as crabs and snails, can be found in great abundance. Fishes are chiefly limited to tidepools during high tide. (Grunion spawn on the high-tide areas of the sandy beach.) There is a community of small and microscopic organisms adapted to the interstices of the sand grains of the beach. Shorebirds actively forage in both rocky and sand areas during low tide. The rocky tidepools adjacent to the CRA on its east side are some of the finest remaining in the County. They are in the Malibu Coastline CRA and the Point Dume State Marine Reserve—no take of any kind is permitted.

Nearshore Subtidal Habitat: The Malibu Coastline CRA has many interactions with the biota of the CRA. Marine habitats include the rocky substrate of the lower level of the intertidal zone on the east side of Point Dume, to subtidal rocky areas that support a great diversity of invertebrates and fishes. Near the CRA this habitat is frequently dominated by giant kelp. The sandy offshore habitat on the west side of Point Dume has a wonderful variety of sand-adapted organisms, ranging from fishes to seastars, to many kinds of jellyfish, mollusks, and other invertebrates.

Kelp Forest Habitat: Giant kelp beds are located down to a depth of approximately 99 feet in the ocean surrounding Point Dume in the Malibu Coastline CRA. The kelp beds are part of one of the temperate zone's most productive habitats, by providing food, attachment sites and shelter for invertebrates and fishes. It is a supremely important nursery and recruitment habitat for juvenile fishes and invertebrates, and an important feeding area for seabirds that use Point Dume for roosting.

Wildlife

The wildlife found in the CRA is dependent upon the two basic regimes found there: marine and terrestrial. The shoreline and bluffs overlooking the coastal sections are home to or visited by a wide variety of shorebirds, migrating birds and marine life. The interior grasslands and ravines have a very

DRAFT

different assemblage than other nearby coastal areas. The CRA ravines are separate and isolated from one another. They exhibit a reduced number of animal species than otherwise might be expected and probably suffer from the effects of fragmentation of a larger, more contiguous ecosystem.

Point Dume is a peninsula that projects its rocky cliff sides out into the Pacific Ocean relative to low bluffs and beaches to the east and west that are heavily impacted by human activity. Point Dume offers a variety of undisturbed habitats for marine and shorebirds. These cliffs and offshore rocks offer ideal roosting and feeding sites for numerous shorebirds, gulls, and other seabirds.

All wildlife species previously recorded, as well as those expected to occur, within the CRA are indicated in the *Comprehensive Floral & Faunal Compendium of the Los Angeles County SEA Update Study 2000 Background Report*. Sensitive wildlife species occurring or potentially occurring within the CRA are discussed in the Sensitive Biological Resources section.

Wildlife Movement

The CRA is on the Pacific Flyway migration route used by many birds seasonally. It plays an important role as a stopover because of its extensive undisturbed marine resources. Point Dume does not fall within any identified terrestrial movement routes for wildlife because it is isolated by the ocean on one side, and miles of developed land on the remaining sides.

Sensitive Biological Resources

Sensitive biological resources are habitats or individual species that have special recognition by federal, state, or local conservation agencies and organizations as endangered, threatened, and/or rare. This is due to the species' declining or limited population sizes, which usually results from habitat loss. Watch lists of such resources are maintained by the California Department of Fish and Wildlife (CDFW), the United States Fish and Wildlife Service (USFWS), and special groups, such as the California Native Plant Society (CNPS). The following sections indicate the habitats as well as plant and animal species present, or potentially present within the CRA, which have been accorded special recognition. When species are federally-listed as endangered or threatened, they often have federally-designated, geographically-specific "critical habitat areas." Critical habitat is a specific geographic area that contains features essential for the conservation of a threatened or endangered species and that may require special management and protection. Critical habitat may include an area that is not currently occupied by the species but that will be needed for its recovery. An area is designated as "critical habitat" after a proposed Federal regulation is published by USFWS in the Federal Register and public comments on the proposal are received and considered. The CRA has proposed critical habitat for the federally-threatened western snowy plover.

Sensitive Plant Communities and Habitats

The CRA supports several habitat types considered sensitive by resource agencies. These are inventoried by California Department of Fish and Wildlife (CDFW) in the California Natural Diversity Database (CNDDDB), or have been assigned a conservation status rank of 1 – 3 at the global or subnational scale. Conservation status ranks are assigned by CDFW, following methodology developed by NatureServe (<http://www.natureserve.org/conservation-tools/standards-methods/conservation-status-assessment>). The CNDDDB includes state and federally-listed endangered, threatened, and rare vascular plants, as well as several sensitive vertebrate species. These communities include coastal sage chaparral scrub and maritime succulent scrub, which occur throughout the CRA. These communities, or closely related designations, are considered high priority communities by the CDFW, which indicates that they are experiencing a decline throughout their

DRAFT

range. The array and composition of these communities has been discussed in the Vegetation section.

Sensitive Plant Species

The following special-status plant taxa have been reported or have the potential to occur within the CRA, based on known habitat requirements and geographic range information:

- Malibu baccharis (*Baccharis malibuensis*) RPR 1B.2
- Coulter’s saltbush (*Atriplex coulteri*) RPR 1B.2
- Slender mariposa lily (*Calochortus clavatus* var. *gracilis*) RPR 1B.1
- Plummer’s mariposa lily (*Calochortus plummerae*) RPR 4.2
- Parry’s spineflower (*Chorizanthe parryi* var. *parryi*) RPR 1B.1
- Blochman’s dudleya (*Dudleya blochmaniae* ssp. *blochmaniae*) RPR 1B.1
- Agoura Hills dudleya (*Dudleya cymosa* ssp. *agourensis*) FT, CSC, RPR 1B.2
- Marcescent dudleya (*Dudleya cymosa* ssp. *marcescens*) FT, SR, RPR 1B.2
- Lyon’s pentachaeta (*Pentachaeta lyonii*) FE, SE, RPR 1B.1
- Sonoran maiden fern (*Thelypteris puberula* var. *sonorensis*) RPR 2.2

Sensitive Animal Species

The following special-status animal species are reported or are likely to be present within the CRA based on habitat requirements and known range attributes:

- Coast horned lizard (*Phrynosoma blainvillii*) CSC
- Two-striped garter snake (*Thamnophis hammondi*) CSC
- Western snowy plover (*Charadrius alexandrinus nivosus*) FT, CSC

Regional Biological Value

The CRA meets all SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE POINT DUME CRA

	Criterion	Status	Justification
	The habitat of core populations of endangered or threatened		This CRA includes habitat that is used by the western snowy plover (<i>Charadrius alexandrinus nivosus</i>). The west side of the headland

DRAFT

	Criterion	Status	Justification
A)	plant or animal species.	Met	includes federally-designated critical habitat that will expand into the CRA.
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The CRA has coastal bluff vegetation on 100-200 foot cliffs, and on the sandy beaches below, is good coastal strand vegetation and habitat. The southern limit of the range for two plants characteristic of the bluffs to the north, giant coreopsis and sea-lettuce is on Point Dume. On the east side of Point Dume, the Malibu Coastline CRA has an excellent rocky intertidal area, which is a habitat that is no longer common in Southern California.
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	Point Dume is one of the two remaining areas in the County where a diverse and healthy mixture of terrestrial Venturan coastal sage scrub and coastal strand vegetation are in close association with marine habitats. Its marine habitats consist of an unprotected rocky shore, with sandy beach pockets, and numerous small caves. The contiguous Malibu Coastline CRA has outlying reefs, rocks, and kelp beds. Due to strong upwelling along the coast, the waters are nutrient-rich and foster highly diverse and productive marine communities, which are protected somewhat by limited public access on the Point Dume headland.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding, feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.	Met	The headland of Point Dume extends into Santa Monica Bay more than a mile beyond the rest of the Malibu coast, and it is located within the Pacific Flyway. As a result, the area is an important jumping off point for migratory birds. Without the remaining terrestrial habitats, this refuge would be lost.
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The close association of native terrestrial vegetation and diverse marine habitats enriched by the nutrient upwelling make the Point Dume area scientifically important for the County. The southern-most point for giant coreopsis and sea lettuce are extreme points for vegetation that are more common to the north.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Met	Relative to other coastal areas, the terrestrial vegetation communities at Point Dume are in good condition, in spite of some degradation from increased runoff and human usage. The marine flora and fauna are in excellent condition, and the association is important to conservation of the County's biodiversity.

In conclusion, the area is a CRA because it contains A) core habitat of the western snowy plover, a federally-threatened species, B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are restricted in distribution in the County and also regionally restricted; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; E) biotic resources that are of scientific interest because they are either an extreme in

DRAFT

physical/geographical limitations, or represent unusual variation in a population or community; and F) areas that provide for the preservation of relatively undisturbed examples of original natural biotic communities in the County.

Santa Catalina Island CRA

For a description of the resources in the Santa Catalina Islands CRA, please refer to the description of Santa Catalina Island in Section V below.

Santa Monica Mountains CRA

For a description of the resources in the Santa Monica Mountains CRA, please refer to the description for the Santa Monica Mountains SEA above.

Terminal Island (Pier 400) CRA

Location

General

The Terminal Island (Pier 400) CRA is located in the southernmost part of the County within the Port of Los Angeles. Terminal Island (Pier 400) supports one of the few remaining areas for breeding utilized by the endangered least terns. The CRA is an artificial dredge spoil island located generally in San Pedro Bay within the Port of Los Angeles. Pier 400 is in the outer harbor area, and the least tern nesting area is on the southern end. It is protected from future development per inter-agency agreement among the Port of Los Angeles, U.S. Fish and Wildlife Service, California Department of Fish and Wildlife, and the U.S. Army Corps of Engineers. The area was specifically designed for least tern nesting and is maintained with buffer from the surrounding Port development, close range to foraging areas, and clearing of the flat sandy area to have little vegetation. During the nesting season (April-August) the site is designated as a no-trespassing area. When least terns nest outside the designated boundaries, a buffer is established for the nest(s) until the chicks are fledged. At other times, the Port may use the site for temporary purposes as long as it is restored prior to the following nesting season. The CRA corresponds directly to the California Audubon-designated Globally Important Bird Area (IBA), Pier 400 Tern Colony IBA.

The site is located in the Long Beach and San Pedro Quadrangles of the United States Geological Survey (USGS) 7.5 Minute Map Series (USGS, 1964).

General Boundary and Resources Description

The CRA is not on a natural island. It is an approximately 15-acre fenced area on the south end of Pier 400, part of Terminal Island, in the Los Angeles Harbor. The area is maintained, prepared, and monitored annually by the Port of Los Angeles for the least tern. The nesting site was created, along with the entire Pier 400 peninsula, of material dredged from the Los Angeles Harbor from 1995-2003. The substrate consists of sand with small rocks and many shell pieces. Non-native flora colonize the area every year. Every year before the breeding period, the Port of Los Angeles has the plants surveyed and then cleared to prepare the terrain for what least terns prefer—a flat sandy area that is virtually free of vegetation that the terns can scrape to provide a shallow depression for their eggs. The northern part of Pier 400 is now a shipping container terminal. There is an undeveloped area, covered with low, volunteer vegetation, west of the CRA that may someday be used by the Port. The CRA site has been used by the least tern for nesting since it was created in 1997. The CRA and the adjacent field on the west are separated and protected from Port activity by a chainlink fence with

DRAFT

locked gate. In addition, the CRA has a peripheral chick fence to keep the chicks from wandering off the CRA area.

Vegetation

The vegetation in this CRA was removed in 2003 and 2004 to allow additional nesting area for the least tern. The common species seen include sea rocket (*Cakile maritima*), tree tobacco (*Nicotiana glauca*), Bermuda grass (*Cynodon dactylon*), puncture vine (*Tribulus terrestris*), and sow thistle (*Sonchus oleraceus*). This removal was judged a success, and clearing has been continued in recent years. Sensitive plant communities have not been found on the site.

Wildlife

Wildlife diversity and abundance within the CRA is limited by its remote location, only accessible by terrestrial animals after traversing long distances through industrial facilities and the narrow, paved causeway of access. Feral cats, rats and mice, and birds associated with development in the region, such as gulls, crows, rock doves, house finches, house sparrows and mockingbirds have been recorded in the area. Native herons that could prey upon the least tern chicks have been observed in the area. The site was also used for breeding in 2011 by two other species that prefer bare, scraped areas: black skimmers (*Rynchops niger*) and Caspian terns (*Hydroprogne caspia*). Many types of shore birds are found nearby.

Wildlife Movement

The site is a low quality habitat for terrestrial animal foraging, and it does not lie within any identified terrestrial movement routes for wildlife. However, it is well located as a linkage site for migratory waterfowl and some birds, specifically the least tern. Surrounding bay waters on three sides provide suitable area for the least tern foraging. The principle prey items they seek are small, schooling fish that frequent the surface area of bay and estuarine waters.

Sensitive Biological Resources

Sensitive biological resources are habitats or individual species that have special recognition by federal, state, or local conservation agencies and organizations as endangered, threatened, and/or rare. This is due to the species' declining or limited population sizes, which usually results from habitat loss. Watch lists of such resources are maintained by the California Department of Fish and Wildlife (CDFW), the United States Fish and Wildlife Service (USFWS), and special groups, such as the California Native Plant Society (CNPS). The following sections indicate the habitats as well as plant and animal species present, or potentially present within the CRA, which have been accorded special recognition.

The site of this CRA is entirely artificial, but it provides the least terns the ideal kind of substrate they prefer: a flat sandy area, devoid of vegetation, that they can scrape for a nesting depression. Artificially created sites (inadvertently created by development) are frequently chosen for breeding by least terns. The site has been used by the least tern for nesting since it was created in 1997 and has been continually managed for this use.

Sensitive Plant Communities and Habitats

The CRA supports no habitat types considered sensitive by resource agencies, namely the CDFW California Natural Diversity Data Base (CNDDB) [2011].

DRAFT

Sensitive Plant Species

No sensitive plants are expected to occur within this CRA.

Sensitive Animal Species

The California least tern (*Sternula antillarum browni*) is both state and federally-endangered. It nests in the CRA and forages in shallow water near the nest site. The western snowy plover (*Charadrius alexandrus nivosus*) is listed as a federally-threatened species and California species of concern, and migrants have been seen but have not nested. Black skimmers ([F-BCC, CA-SSC]) used the site for nesting in the 2011 season.

Regional Biological Value

The CRA meets several SEA designation criteria and supports many regional biological values. Each criterion and how it is met is described below.

CRITERIA ANALYSIS OF THE TERMINAL ISLAND (PIER 400) CRA

	Criterion	Status	Justification
A)	The habitat of core populations of endangered or threatened plant or animal species.	Met	This area was set aside because endangered least terns use it for breeding in the summer. Their breeding depends on availability of forage fish, and they may not use the area consistently. In addition there may be some breeding by the black skimmer (bird of concern for the State and USFWS).
B)	On a regional basis, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	This area was set aside because endangered least terns use it for breeding in the summer. Their breeding depends on availability of forage fish, and they may not use the area consistently. In addition there may be some breeding by the black skimmer (bird of concern for the State and USFWS).
C)	Within the County, biotic communities, vegetative associations, and habitat of plant or animal species that are either unique or are restricted in distribution.	Met	The least tern typically breeds in flat sandy areas that are created by disturbance, but are free of disturbance and vegetation during the breeding period. The area needs to be close to a source of the terns' prey, which are small, estuarine or surface-schooling marine fishes. The choice breeding areas require no disturbance because the terns make an unsheltered scoop for a nest, and eggs and nestlings would be crushed by constant traffic of vehicles or pedestrians. This kind of habitat was once common along the beaches of Southern California, but has nearly disappeared as estuaries have been filled and channelized, and sandy beaches have become a favorite Southern California recreation area.
D)	Habitat that at some point in the life cycle of a species or group of species, serves as concentrated breeding,	Met	Los Angeles has many fine beaches that are potential nesting sites, but nearly all have become recreation areas. The terns have continued to use the areas with protection from disturbance, which are a small number compared to the previously available undisturbed sites.

DRAFT

	Criterion	Status	Justification
	feeding, resting, or migrating grounds and is limited in availability either regionally or in the County.		
E)	Biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.	Met	The Terminal Island site is named as a CRA for the summer bird breeding that occurs.
F)	Areas that would provide for the preservation of relatively undisturbed examples of the original natural biotic communities in the County.	Not Met	Due to the artificial construct of the area, the CRA does not serve as a relatively undisturbed example of the original natural biotic communities in the County.

In conclusion, the area is a CRA because it contains A) the habitat of core populations of endangered and threatened plant and animal species; B-C) biotic communities, vegetative associations, and habitat of plant and animal species that are either unique or are restricted in distribution in the County and regionally; D) concentrated breeding, feeding, resting, or migrating grounds, which are limited in availability in the County; and E) biotic resources that are of scientific interest because they are either an extreme in physical/geographical limitations, or represent unusual variation in a population or community.

CRA Description Sources

Alamitos Bay CRA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 2006. *The Alamitos Bay SEA Description*, Los Angeles County, California.

Fish and Wildlife Services. 2000. *Federal Register*. "Endangered and Threatened Wildlife and Plants;

DRAFT

Proposed Determination of Critical Habitat for the Coastal California Gnatcatcher; Proposed Rule". Fish and Wildlife Services; 32 pp.

Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.

LSA Associates, Inc. October 2009. *Draft Environmental Impact Report: Alamitos Bay Marina Rehabilitation Project*, Prepared for the City of Long Beach.

Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.

U.S. Fish and Wildlife Service. 1997. *Vernal Pools of Southern California Draft Recovery Plan*. U.S. Fish and Wildlife Service, Portland, Oregon. 113+pp.

Zedler, P.H. 1987. *The Ecology of Southern California Vernal Pools: A Community Profile*. U.S. Fish and Wildlife Service Biological. Report 85(7.11). 136 pp.

Ballona Wetlands CRA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals* (898 taxa). 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 2006. *The Ballona Wetlands SEA Description*, Los Angeles County, California.

Fish and Wildlife Services. 2000. *Federal Register*. "Endangered and Threatened Wildlife and Plants; Proposed Determination of Critical Habitat for the Coastal California Gnatcatcher; Proposed Rule". Fish and Wildlife Services; 32 pp.

Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.

Philip Williams & Associates, Ltd. 2006. *Ballona Wetland Existing Conditions DRAFT Report, Section 5. Biological Resources*. Prepared for California State Coastal Conservancy.

Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.

U.S. Fish and Wildlife Service. 1997. *Vernal Pools of Southern California Draft Recovery Plan*. U.S. Fish and Wildlife Service, Portland, Oregon. 113+pp.

DRAFT

Zedler, P.H. 1987. *The Ecology of Southern California Vernal Pools: A Community Profile*. U.S. Fish and Wildlife Service Biological. Report 85(7.11). 136 pp.

El Segundo Dunes CRA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 2006. *The El Segundo Dunes SEA Description*, Los Angeles County, California.

England & Nelson Environmental Consultants. 1976. *Land Capability/Suitability Study, Los Angeles County General Plan Revision Program, Significant Ecological Areas Report*, Riverside, California.

Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.

Sierra Delta Corporation. 1988. *Biota Report for Zond Systems, Inc., Los Angeles Wind Electric Generating Station, Gorman, California, Conditional Use Permit 86453*, Las Vegas, Nevada.

Sapphos Environmental, Inc. January 2005. *Los Angeles/El Segundo Dunes Habitat Restoration Plan*. Prepared for Los Angeles World Airports and U.S. Department of Transportation, Federal Aviation Administration.

Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation*, Second Edition Sacramento: California Native Plant Society Press.

Malibu Coastline CRA Sources

Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti (editors, 2012) *The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

DRAFT

- City of Malibu. 2009. Malibu General Plan, *Section II, Elements of the General Plan, Chapter 3.0 Conservation Element, 3.2 Existing Natural Resources*.
- City of Malibu. 2007. *Draft Proposed Environmentally Sensitive Habitat Areas (ESHA) Findings for the City of Malibu LCP*.
- County of Los Angeles, Department of Regional Planning. 1986. *Malibu Local Coastal Program Land Use Plan*. Los Angeles County, California.
- County of Los Angeles, Department of Regional Planning. 2006. *The Malibu Coastline SEA Description*, Los Angeles County, California.
- Fish and Wildlife Services. 2000. *Federal Register*. "Endangered and Threatened Wildlife and Plants; Proposed Determination of Critical Habitat for the Coastal California Gnatcatcher; Proposed Rule". Fish and Wildlife Services; 32 pp.
- Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.
- Ryan Ecological Consulting. 2010. The Western Snowy Plover in Los Angeles County, California. Report to California Department of Fish and Wildlife. 136 pp.
- <http://losangelesaudubon.org/images/stories/pdf/westernsnowyploverin%20losangelescounty.pdf>
- Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.
- U.S. Fish and Wildlife Service. 1997. *Vernal Pools of Southern California Draft Recovery Plan*. U.S. Fish and Wildlife Service, Portland, Oregon. 113+pp.
- Zedler, P.H. 1987. *The Ecology of Southern California Vernal Pools: A Community Profile*. U.S. Fish and Wildlife Service Biological. Report 85(7.11). 136 pp.
- Point Dume CRA Sources
- Baldwin, Bruce G., Douglas H. Goldman, David J. Keil, Robert Patterson, and Thomas J. Rosatti(editors, 2012)*The Jepson Manual: Vascular Plants of California, Thoroughly Revised and Expanded*. Second Edition. Jepson Herbarium, Berkeley, CA. 1600 pp., illus.
- California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.
- California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.
- California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.
- City of Malibu. 2009. Malibu General Plan, *Section II, Elements of the General Plan, Chapter 3.0 Conservation Element, 3.2 Existing Natural Resources*.
- Holland RF. *Preliminary descriptions of the terrestrial natural communities of California*. 1986 and

DRAFT

1992 Update. California Department of Fish and Wildlife unpublished report.

Ryan Ecological Consulting. 2010. The Western Snowy Plover in Los Angeles County, California. Report to California Department of Fish and Wildlife. 136 pp.

<http://losangelesaudubon.org/images/stories/pdf/westernsnowyploverin%20losangelescounty.pdf>

Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.

U.S. Department of Commerce. National Oceanic and Atmospheric Administration. National Marine Sanctuary Program. 2008. *Channel Islands National Marine Sanctuary Management Plan/Final Environmental Impact Statement*. Silver Spring, MD

Terminal Island (Pier 400) CRA Sources

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *Special Animals (898 taxa)*. 60 pp.

California Department of Fish and Wildlife, Natural Diversity Database. April 2011. *Special Vascular Plants, Bryophytes, and Lichens List*. Quarterly publication. 71 pp.

California Department of Fish and Wildlife, Natural Diversity Database. January 2011. *State and Federally Listed Endangered and Threatened Animals of California*. 13 pp.

County of Los Angeles, Department of Regional Planning. 2006. *The Pier 400, Terminal Island SEA Description*, Los Angeles County, California.

Holland RF. Preliminary descriptions of the terrestrial natural communities of California. 1986 and 1992 Update. California Department of Fish and Wildlife unpublished report.

Science Applications International Corporation. 2008. Draft Biological Assessment, Pacific L.A. Marine Terminal Crude Oil Terminal Project. Prepared for the U.S. Army Corps of Engineers, Los Angeles District and Port of Los Angeles, Harbor Department.

Sawyer, Keeler-Wolf and Evens. 2009. *A Manual of California Vegetation, Second Edition* Sacramento: California Native Plant Society Press.

V. Coastal Zone Resources

Coastal Zone Resources include natural resources in the the County's coastal areas, which are neither SEAs nor CRAs, and areas that contain SEAs or CRAs but are larger than one single SEA or CRA.

Ballona Wetlands

For a description of the resources in the Ballona Wetlands, please refer to the description for the Marina del Rey below, and the description for the Ballona Wetlands CRA in Section IV.

Marina del Rey⁷

Marina del Rey is a man-made feature that occupies part of the historical estuary near the mouth of Ballona Creek. Prior to the arrival of Europeans, the Ballona area was an ancient estuary of the Los Angeles River, with the mouth of the river entering the Pacific Ocean north of the Westchester Bluffs, forming a broad coastal plain with seasonal and permanent wetlands extending north and east toward higher ground in present-day Santa Monica.⁸ A long range of natural sand dunes cut off the ocean from the majority of the low-lying ground, which featured a network of tidal channels and extensive salt marsh (i.e., the historical "Venice Marshes" or "Ballona Wetlands"). Until a flood in the early 1800s, the Los Angeles River emptied at Santa Monica Bay, along the current course of Ballona Creek.⁹ The historical landscape along the coast west of present-day Lincoln Boulevard (i.e., an area encompassing all of Marina del Rey) likely consisted of wide tidal channels and mudflats, salt marshes, coastal dunes, pockets of freshwater and/or brackish marsh, as well as riparian scrub. Also present was a coastal prairie community described by researchers as far back as the 1930s (e.g., "the meadow" referred to by von Bloeker 1943¹⁰). These are generally the habitat types typical of coastal estuaries throughout Southern California and northwestern Baja California, Mexico.¹¹ Comparable coastal estuaries on broad plains in Southern California include Carpinteria Marsh, Mugu Lagoon, Alamitos Bay, Bolsa Chica, Upper Newport Bay, and the Tijuana River Estuary.

By the mid-1900s, much of Ballona Creek had been excavated and routed through a channel, at first

⁷Adapted from Hamilton, RA and Cooper, DS, 2010. Conservation and Management Plan for Marina del Rey, Los Angeles County, California. Prepared for the County Los Angeles Department of Beaches and Harbors and the Department of Regional Planning.

⁸ Cooper, D. S. 2008. The use of historical data in the restoration of the avifauna of the Ballona Wetlands, Los Angeles County, California. *Natural Areas Journal* 28:83–90.

⁹ Gumprecht B. 1999. *The Los Angeles River: Its Life, Death, and Possible Rebirth*. Johns Hopkins University Press, Baltimore, MD.

¹⁰von Bloeker, J. C. 1943. The fauna and flora of the El Segundo sand dunes: Birds of El Segundo and Playa del Rey. *Bulletin of the Southern California Academy of Sciences* 42:1–30 (Part 1) and 90–103 (Part 2).

¹¹ Grewell, B. J., Callaway, J. C., and Ferren, W. R., Jr. 2007. Estuarine wetlands. Pp. 124–154 in *Terrestrial Vegetation of California* (M. G. Barbour, T. Keeler-Wolf, and A. A. Schoenherr, eds.). University of California Press, Berkeley. Pickart, A. J., and Barbour, M. G. 2007. Beach and dune. Pp. 155–179 in *Terrestrial Vegetation of California* (M. G. Barbour, T. Keeler-Wolf, and A. A. Schoenherr, eds.). University of California Press, Berkeley.

DRAFT

earthen (1920s), then concrete-lined (late 1930s), principally to control floods in the Ballona Valley that regularly destroyed cropland and generally hindered development. The most serious and final impact to lower Ballona Creek and the majority of its natural wetlands came in the early 1960s, with the completion of Marina del Rey, which eliminated nearly all the functional wetlands north of the Ballona Creek channel and left only a small remnant to the south, along Culver Boulevard. However, just as the creation of Marina del Rey development entailed the elimination of certain natural habitats, it created novel ones, with the addition of hundreds of evergreen, semi-tropical, trees, as well as irrigated lawns and man-made structures.

Today the channel waters of Marina del Rey and its breakwater are designated by California Audubon as a part of the Ballona Wetlands State Important Bird Area (IBA), recognizing the important natural resources that still occur in this district of the County. The IBA includes the Ballona Lagoon in nearby Venice, the Del Rey Lagoon across Ballona Creek in Playa Del Rey, the least tern nesting area on Venice City Beach, the Strand Beach south of the Ballona Channel opening, the Ballona Channel, the Ballona Wetlands both north and south of the Ballona Channel, and Bluff Creek that feeds the Ballona Freshwater Marsh. The IBA designation recognizes this as a unit, all the parts contributing to the welfare of the whole. As an example, the federally-endangered least terns that breed on Venice City Beach need to forage widely, in all parts of the IBA waters and beyond in order to successfully raise their young to fledglings.

Vegetation

Vegetation of the Marina del Rey has not been formally described; however, the Conservation & Management Plan for Marina del Rey, Los Angeles County, California (Hamilton and Cooper, 2010.) discusses six areas within Marina del Rey with regard to their value as avian resource areas. Because no vegetation map has been prepared for the Marina, these areas are listed individually below, along with any corresponding natural or Semi-Natural vegetation types that are to be expected within each of the six areas.

Oxford Stormwater Retention Basin: Provides a flood-control function for the development and is lined with native and non-native species. The shoreline of the basin has been landscaped extensively with non-native trees and shrubs, especially small-flowered myoporum (*Myoporum laetum*). The myoporum landscaping currently in poor health, presumably due to an infestation of the myoporum thrip (*Klambothrips myopori*).

Corresponding MCV communities:

- *Schinus (molle, terebinthifolius)-Myoporum laetum* (Pepper tree or Myoporum groves) Semi-Natural Woodland Stands
- *Eucalyptus (globulus, camaldulensis)* (Eucalyptus groves) Semi-Natural Woodland Stands

Ballona Lagoon: The northern extent of the former coastal lagoon at the mouth of the Ballona Wetlands, and now the southern extension of the “Grand Canal” in Venice (adjacent to and just west of Marina del Rey), this site has been known as “Ballona Lagoon” since 1996 when extensive habitat restoration was completed in an effort to bring back a native coastal scrub community. The lagoon is tidal, and a band of mudflat is usually exposed around the entire lagoon, but only the upper/northern end drains completely except during the most extreme low tides. Saltmarsh vegetation forms a ring around the upper mudflat, below the coastal scrub.

Corresponding MCV communities

DRAFT

- *Schinus (molle, terebinthifolius)-Myoporum laetum* (pepper tree or myoporum groves) Semi-Natural Woodland Stands
- *Carpobrotus edulis* or other ice plants (ice plant mats) Semi-NaturalHerbaceous Stands
- *Sarcocornia pacifica (Salicornia depressa)* (pickleweed mats) Herbaceous Alliance
- Orcutt's pincushion (*Chaenactis glabriuscula* var. *orcuttiana*) new Alliance

Del Rey Lagoon: This area is entirely landscaped with lawn and ornamental non-native *Ficus* and *Acacia* trees.

Corresponding MCV communities: None at this time.

Ballona Wetlands: This, the main tidal marsh area remaining at Ballona, is located between the Ballona Creek channel and Culver Boulevard. It features extensive pickleweed (*Salicornia* spp.) marsh habitat, muddy tidal channels, and a large saltpan that is irregularly moistened by rain, dense fog, and high tides.

Corresponding MCV communities

- *Sarcocornia pacifica (Salicornia depressa)* (pickleweed mats) Herbaceous Alliance

Ballona Freshwater Marsh: This marsh, constructed in 2003 at the corner of Lincoln and Jefferson Boulevards, just south of Marina del Rey, supports large expanses of tules, cattails, and other freshwater marsh vegetation.

Corresponding MCV communities

- *Schoenoplectus acutus* (hardstem bulrush marsh) Herbaceous Alliance
- *Schoenoplectus californicus* (California bulrush marsh) Herbaceous Alliance
- *Typha (angustifolia, domingensis, latifolia)* (cattail marshes) Herbaceous Alliance

Centinela Confluence: This refers to the tidally-influenced confluence of Ballona Creek at the Centinela Channel, just south of the State Route-90 bridge. A patch of tall, lush grasses serves as a consistent roosting and foraging location for herons, egrets, brown pelicans, gulls, terns, and shorebirds.

Corresponding MCV communities: None at this time.

Wildlife

As reviewed by Cooper¹², many bird species associated with freshwater marsh, coastal lagoon, and

¹² Cooper, D. S. 2008. The use of historical data in the restoration of the avifauna of the Ballona Wetlands, Los Angeles County, California. *Natural Areas Journal* 28:83–90.

riparian habitats were lost from the Ballona area during the early period of development (pre-1930s); many saltmarsh species, including waterfowl and shorebirds that occurred in large flocks, suffered heavy losses during the middle period (1940s to 1960s); and since the 1960s, many open-country species, particularly those of agricultural fields and extensive grasslands, have either been extirpated or experienced serious declines.

Cooper (2006¹³) documented the ongoing colonization of the local area by bird species that require tall trees for breeding and/or foraging, and by species frequently associated with human habitation. This colonization phenomenon has intensified as the Marina's non-native landscaping has matured, providing much more structural complexity than was present formerly, but at the expense of numerous species that depend on natural, wild habitats for their persistence in the landscape or for refueling during long migrations. Currently, the area is used extensively by nesting and foraging great blue heron (*Ardea herodias*), great egret (*A. alba*), snowy egret (*Egretta thula*), black-crowned night heron (*Nycticorax nycticorax*), and double-crested cormorant (*Phalacrocorax auritus*).

A well-documented least tern colony (federally and state-listed Endangered) is present nearby at Venice Beach and is known to have suffered predation by crows and black-crowned night herons, both of which are dependent on non-native landscaping and artificial structures within and surrounding Marina del Rey for nesting substrate.

Sensitive Biological Resources

The statuses of rare plants are hierarchically categorized by the CNPS using a rank and decimal system. The initial category level of Rare Plant Rank (RPR) is indicated by the ranks 1A (presumed extinct), 1B (rare or endangered in California and elsewhere), 2A (presumed extinct in California but extant elsewhere), 2B (rare or endangered in California but more common elsewhere), 3 (more information needed, a review list), and 4 (limited distribution). In cases where the CNPS has further identified the specific threat to the species, a decimal or Threat Code is added: .1 (seriously endangered in California), .2 (fairly endangered in California), or .3 (not very endangered in California).

The following special-status plant taxa have been reported or have the potential to occur within the SEA, based on known habitat requirements and geographic range information:

- Southern tarplant (*Centromadia parryi* ssp. *australis*)RPR 1B.1
- Orcutt's pincushion (*Chaenactis glabriuscula* var. *orcuttiana*)RPR 1B.1
- Coastal goosefoot (*Chenopodium littoreum*)RPR 1B.2

¹³ Cooper, D. S. 2006. Annotated checklist of extirpated, reestablished, and newly-colonized avian taxa of the Ballona Valley, Los Angeles County, California. Bulletin of the Southern California Academy of Sciences 105:91–112.

DRAFT

- Coulter's goldfields (*Lasthenia glabrata* ssp. *coulteri*) RPR 1B.1
- Ballona cinquefoil (*Potentilla multijuga*)RPR 1A
- Estuary seablite (*Suaeda esteroa*)RPR 1B.2

The following special-status animal species are reported or are likely to be present within the SEA based on habitat requirements and known range attributes:

- Mimic tryonia (*Tryonia imitator*)CDFW Special Animals List
- Wandering skipper (*Panoquina errans*)CDFW Special Animals List
- Silvery legless lizard (*Anniella pulchra pulchra*)FSS, SSC
- Two-striped garter snake (*Thamnophis hammondi*)BLMS, FSS, SSC
- Tricolored blackbird (*Agelaius tricolor*)BCC, BLMS, SSC, USBC, AWL, ABC
- Western snowy plover (*Charadrius alexandrinus nivosus*)FT, BCC, SSC, ABC, AWL, USBC
- California black rail (*Laterallus jamaicensis coturniculus*)BCC, ST, CDFW Fully Protected, USBC, AWL, ABC
- Belding's savannah sparrow (*Passerculus sandwichensis beldingi*)SE
- California least tern (*Sternula antillarum browni*)FE, SE, CDFW Fully Protected, USBC, ABC
- South coast marsh vole (*Microtus californicus stephensi*)SSC
- Southern California saltmarsh shrew (*Sorex ornatus salicornicus*)SSC

San Clemente Island

San Clemente Island lies approximately 63 miles south of the City of Long Beach and 78 miles west of the City of San Diego. San Clemente Island is approximately 24 miles long and 5 miles across at its widest point. It has a land area of approximately 57 square miles. Since 1934, San Clemente Island has been owned and operated by the U.S. Navy. More than a dozen range and operational areas are clustered within a 60 mile radius of San Clemente Island. The Commander-in-Chief, Naval Forces, Pacific (CINCPACFLT) is the major claimant for San Clemente Island, and the Naval Air Station, North Island (NASNI) is responsible for its administration.

The climate of the Island is classified as semi-arid, which is similar to the Southern California coastal areas, except for more fog and overcast weather and cooler year-round temperatures. Aided by the Santa Ana winds, temperatures exceeding 90 degrees occasionally occur between August and October. Rainfall is slight and occurs mostly between the months of November and April (San Clemente Island Master Plan).

The Island is a very important biological resource for the County, California, and the U.S. Any island is a microcosm of evolutionary processes, and studies of island biota lead to many insights on biological community processes—how food webs are structured and how to maintain them in the

DRAFT

context of impacts. The Island has 14 endemic plants (known only from the Island; 6 of which are federally-endangered) and 29 other plants known only from the Island and other Channel Islands (the northern ones and Santa Catalina Island). Some of the Island's endemics have their closest relatives on islands off of Baja California, which is over 500 miles away, and not the nearest neighboring islands. The plants had a major setback with about 400 years of goat and pig grazing, but the Navy has finally removed the last of these (a goat in 1992), and plants have responded vigorously to predation relief in most cases.

There is multiple federal recognition of the Island's resources. The Island has endemic birds, notably the federally-endangered shrike (*Lanius ludovicianus mearnsi*) and the federally-threatened San Clemente sage sparrow (*Amphispiza belli clementeae*). The shrike crashed to a population of 14 birds in 1998, but with a Navy recovery program of incubating eggs, the shrikes have rebounded to about 200 individuals. The chief predators were or are the introduced black rat and feral cats, but the goats did major indirect damage by undermining and browsing the trees that the shrike needs to breed. The Island has an endemic federally-threatened Island night lizard (*Xantusia riversiana*), and a subspecies of the island fox (*Urocyon littoralis clementae*) that is state-threatened and monitored by the Navy. Although it does not include offshore reefs, rocks, and pinnacles under the jurisdiction of the armed forces, the California Coastal National Monument, under the Bureau of Land Management in the U.S. Department of the Interior, manages the rocks within one nautical mile of the coast of the Island under this program.

State recognition has been equally strong. The Island has many state-recognized sensitive species. The Island is designated by the California State Water Resources Control Board as an ASBS (Area of Special Biological Significance), which is a marine area with exceptionally good water quality and natural community features. Populated and disturbed areas along the shore are zones of exception, and there is one small designated area at the sewage outfall out of 100 natural gullies and ephemeral streams and 23 known Navy discharges. The Island is reputedly a favorite of snorkelers and divers for the marine life, and its waters are on the main migration path for the CITES Appendix I California gray whale (*Eschrichtius robustus*).

The Island and its nearshore out to about 0.8 mile offshore is designated by California Audubon as the San Clemente Island Globally Important Bird Area (IBA). Over 240 bird species regularly use the Island, and about 30 species inhabit the Island year-round.

The Island has unique features that are mysterious in some respects—phenomena waiting for discovery. The Island is nearly surrounded by sea cliffs, which in their steepness, provide the protection from predators (including introduced rats) that seabirds need for breeding. It is probable that some breeding by marine birds is unknown, since they often forage and return to feed their offspring at night, which is a difficult phenomenon to observe for predators and humans alike. Colonies are yet to be discovered. A known marine bird that has this breeding pattern is the federal candidate for listing, Xantus' murrelet (*Synthliboramphus hypoleucus*). Because it is so distant from shore, this Island has many oceanic influences that are unique for the Channel Islands—much more than the other nearby islands. For example, an oceanic bird, the red-tailed tropic bird (*Phaethon rubricauda*), once had some breeding on Pyramid Head before this area became part of the shore bombardment range. The Ashy storm-petrel (*Oceanodroma homochroa*), which is a state species of concern and one of the rarest storm-petrels, is known to breed in the Seal Cove area, and could also be breeding elsewhere on the Island.

Vegetation

Despite the Island's barren look, a large variety of plant life is present. The Island harbors 14 endemic plants and 29 insular endemic plants restricted to the Channel Island system (San Clemente Island Master Plan).

Domestic animals introduced to the Island, including cats, pigs, and goats, have greatly impacted the native plant communities. Beginning in 1875, the introduction of Spanish goats drastically altered the ecological structure of the Island due to constant foraging and grazing. Native plants and animals were significantly reduced, which resulted in an effort to remove the goats from the Island. By 1992, the U.S. Fish and Wildlife Service removed all goats and pigs from the Island (USFWS, California Channel Islands Species Recovery Plan).

Plant communities on the Island were classified using standard methodology and terminology. Most of the communities discussed correspond directly with those listed in Holland's Preliminary Descriptions of the Terrestrial Natural Communities of California (1986 and 1992 update). Other communities are named based on dominant species within them and/or commonly used terminology. Descriptions and general locations of each plant community on the Island, including maritime desert scrub, maritime sage scrub, island woodland, grassland, coastal salt marsh, and disturbed areas are given below.

In recent years, ecologists have refined previous vegetation classification approaches to define communities primarily by their constituent plant assemblages, and have now widely adopted the classification system described by Sawyer, Keeler-Wolf and Evens in *A Manual of California Vegetation, Second Edition 2009 (MCV)*. This has become the accepted standard recognized by the California Department of Fish and Wildlife, the California Native Plant Society and the U.S. Fish and Wildlife Service.

The important difference between current and earlier methods is that earlier classifications were based on a variety of factors, such as physiographic features, as in the case of vernal pools; or by specific plants, as in the case of coast live oak woodland; or by the use of commonly accepted terms, as in chaparral. In the MCV, plant communities are defined with more precision as botanical alliances where one, or occasionally two, plant species are dominant or co-dominant with a host of other possible associated plants. The MCV lists no one plant community called "chaparral" because in habitats of this type, any one of a variety of shrubs can be dominant and influence the character of the vegetation. For example, in a location where chamise (*Adenostoma fasciculatum*) is predominant, the alliance is classified as an *Adenostoma fasciculatum* Shrubland Alliance, while close by, greenbark ceanothus (*Ceanothus spinosus*) may emerge as the most common shrub and this is termed a *Ceanothus spinosus* Shrubland Alliance.

The transition to the new MCV format is ongoing. Not all alliances have been fully described and new ones are still being recognized. As such, a significant amount of field work and site surveying for accuracy would be needed to verify all existing alliances on the Island, as well as extensive research to define every alliance in the new MCV plant communities' format. The more familiar nomenclatures will continue to be used for the Island when appropriate.

An effort has been made to conform to this new format. Descriptions and general locations of each plant community as described in the California Channel Islands Species Recovery Plan (USFWS, 1984) appear below. The plant communities correspond to the classifications used in that document. Each was considered baseline information and evaluated for the potential presence of alliances as

DRAFT

described in the MCV. Alliances with matching profiles of the given criteria are listed. In many cases, only with further investigation can the presence of some alliances be confirmed. In addition, it should be noted that not all alliances are listed within this description as many alliances have yet to be defined and new alliances are still being discovered. Descriptions and general locations of the each plant community present on the Island are given below.

Maritime Desert Scrub - (Lycium Phase, Typical Phase, and Cholla Phase): Includes a mixture of low growing, dry-season deciduous scrubs and cactus located along the southern and western shores. The predominant deciduous shrubs include California desert-thorn (*Lycium californicum*) and island senecio (*Senecio lyonii*), while California brittle-bush (*Encelia californica*) and California sagebrush (*Artemisia californica*) are the predominant maritime sage scrub species. Coastal prickly-pear (*Opuntia littoralis*) also exists in abundance. The south-facing slopes at the southern end of the island are dominated by coastal cholla (*Cylindropuntia prolifera*), but box thorn and cunyado (*Bergerocactus emoryi*) also exist.

Corresponding MCV communities:

- *Lycium californicum* (California desert-thorn) Shrubland Provisional Alliance
- *Encelia californica* (California brittle bush scrub) Shrubland Alliance
- *Artemisia californica* (California sagebrush scrub) Shrubland Alliance
- *Opuntia littoralis* (coast prickly pear scrub) Shrubland Alliance

Maritime Sage Scrub: Located on steep eastern slopes and in some canyons on the west side of the Island. Lemonadeberry (*Rhus integrifolia*) can be found on both sides of the Island on relatively level ground. Many types of chaparral shrubs are present, including chamise, island ceanothus (*Ceanothus megacarpus* var. *insularis*), Catalina crossosoma (*Crossosoma californicum*), Channel Island tree poppy (*Dendromecon harfordii*), toyon (*Heteromeles arbutifolia*), laurel sumac (*Malosma laurina*), Catalina cherry (*Prunus ilicifolia* ssp. *lyonii*), island redberry (*Rhamnus pirifolia*), blue elderberry (*Sambucus nigra* var. *caerulea*), and poison oak (*Toxicodendron diversilobum*).

Corresponding MCV communities:

- *Rhus integrifolia* (lemonade berry scrub) Shrubland Alliance
- *Ceanothus megacarpus* (big pod ceanothus chaparral) Shrubland Alliance
- *Malosma laurina* (laurel sumac scrub) Shrubland Alliance

Island Woodland: Located in the western and eastern canyons, is the only tree vegetation type present on the Island. It includes Catalina cherry, Santa Cruz Island ironwood (*Lyonothamnus floribundus* ssp. *aspleniifolius*), island oak (*Quercus tomentella*), blue elderberry, and toyon.

Corresponding MCV communities:

- *Prunus ilicifolia* (holly leaf cherry chaparral) Shrubland Alliance
- *Lyonothamnus floribundus* (Catalina ironwood groves) Woodland Special Stands

DRAFT

- *Quercus tomentella* (Island oak groves) Woodland Special Stands

Grasslands: Cover the central uplands of the Island. Although dominated by exotic annual herbs, scattered native wildflowers also exist. Native perennial needlegrass, larkspur, and brodiaea are also present.

Corresponding MCV communities:

- *Bromus (diandrus, hordeaceus)-Brachypodium distachyon* (annual brome grasslands) Semi-Natural Herbaceous Stands
- *Bromus rubens-Schismus (arabicus, barbatus)* (red brome or Mediterranean grass grasslands) Semi-Natural Herbaceous Stands
- *Lasthenia californica-Plantago erecta-Vulpia microstachys* (California goldfields-dwarf plantain-six-weeks fescue flower fields) Herbaceous Alliance
- *Nassella [Stipa] pulchra* (purple needlegrass grassland) Herbaceous Alliance

Coastal Strand and Dunes: Coastal strand and dune vegetation is restricted to small areas along the south and northwest coasts of the Island. Predominantly associated with sandy substrates, several species typical of this vegetation type also occur in scattered rocky areas behind beaches. Introduced crystalline iceplant (*Mesembryanthemum crystallinum*) and grasses are abundant.

Corresponding MCV communities:

- *Carpobrotus edulis* or other ice plants (ice plant mats) Semi-Natural Herbaceous Stands
- *Abronia latifolia - Ambrosia chamissonis* (dune mat) Herbaceous Alliance

Coastal Salt Marsh: Exist on the south side of the Island. These areas are proximate to the impact zone of the shore bombardment range, and therefore, the biological information is somewhat limited. However, pickleweed (*Salicornia pacifica*), alkali heath (*Frankenia salina*), saltgrass (*Distichlis spicata*), and California sea-blite (*Suaeda californica*) are known to exist.

Corresponding MCV communities:

- *Sarcocornia [Salicornia] pacifica(Salicornia depressa)(pickleweed mats)Herbaceous Alliance*
- *Frankenia salina(alkali heath marsh)Herbaceous Alliance*
- *Distichlis spicata(salt grass flats)Herbaceous Alliance*

Disturbed Areas: Mainly consist of plants that are native to the Old World. Once limited to ruderal areas, these areas have expanded and exotic species are now common throughout the Island.

Corresponding MCV communities:

- *Avena (barbata, fatua)(wild oats grasslands)Semi-NaturalHerbaceous Stands*
- *Brassica nigra* and other mustards (upland mustards)Semi-NaturalHerbaceous Stands

DRAFT

- *Bromus (diandrus, hordeaceus)-Brachypodium distachyon* (annual brome grasslands)Semi-NaturalHerbaceous Stands
- *Bromus rubens-Schismus (arabicus, barbatus)*(red brome or Mediterranean grass grasslands)Semi-NaturalHerbaceous Stands

(USFWS, California Channel Islands Species Recovery Plan)

Wildlife

An abundance of marine mammals thrive in waters surrounding the Island. The most prominent are the California sea lion (*Zalophus californianus*) and the harbor seal (*Phoca vitulina*); however, porpoises and whales are frequently sighted. The Island is in the path of a migration route, which along with great depths and caves, creates a rich and varied marine environment (San Clemente Island Master Plan).

The Island supports a unique assortment of terrestrial fauna. There are two lizard species on the Island: western side-blotched lizard (*Uta stansburiana elegans*) and the San Clemente Island night lizard (*Xantusia riversiana reticulata*). As many as 240 bird species inhabit the Island at various periods of the year. Around 30 species breed on the Island, while the other species are mainly migrants. Some of the bird species include island horned lark (*Eremophila alpestris insularis*), western meadowlark (*Sturnella neglecta*), white-crowned sparrow (*Zonotrichia leucophrys*), American kestrel (*Falco sparverius*), Gambel's quail (*Callipepla gambelii*), and the exotic chukar (*Alectoris chukar*). Waterbirds are attracted to the rocky shore of the Island, which provides opportunities for foraging and resting. Some of the waterbirds on the Island include western gull (*Larus occidentalis*), Brandt's cormorant (*Phalacrocorax penicillatus*), California brown pelican (*Pelecanus occidentalis californicus*), royal tern (*Thalasseus maximus*), black oystercatcher (*Haematopus bachmani*), and Xantus' murrelet (*Synthliboramphus hypoleucus*)(USFWS, California Channel Islands Species Recovery Plan).

The Island is also home to the following bat species: California myotis (*Myotis californicus*), fringed myotis (*M. thysanodes*), Townsend's big-eared bat (*Corynorhinus townsendii townsendii*), and Mexican free-tailed bat (*Tadarida brasiliensis*). The San Clemente deer mouse (*Peromyscus maniculatus clementis*) is also native to the Island. One of the most notable mammals on the Island is the San Clemente Island fox (*Urocyon littoralis clementae*), which inhabits most of the Island (USFWS, California Channel Islands Species Recovery Plan).

Sensitive Biological Resources

The statuses of rare plants are hierarchically categorized by the CNPS using a rank and decimal system. The initial category level of Rare Plant Rank (RPR) is indicated by the ranks 1A (presumed extinct), 1B (rare or endangered in Californiaand elsewhere), 2A (presumed extinct in California but extant elsewhere), 2B (rare or endangered in California but more common elsewhere), 3 (more information needed, a review list), and 4 (limited distribution). In cases where the CNPS has further identified the specific threat to the species, a decimal or Threat Code is added: .1 (seriously endangered in California), .2 (fairly endangered in California), or .3 (not very endangered in California).

The following special-status plant taxa have been reported or have the potential to occur on San

DRAFT

Clemente Island, based on known habitat requirements and geographic range information:¹⁴

- Woven-spored lichen (*Texosporium sancti-jacobi*) CDFW Special Plants List, RPR 1B.2
- Red sand-verbena (*Abronia maritima*) RPR 4.2
- San Clemente Island bird's-foot trefoil (*Acemisson argophyllus* var. *adsurgens*) SE, RPR 1B.1
- San Clemente Island lotus (*Acemisson dendroideus* var. *traskiae*) FE, SE, RPR 1B.1
- Aphanisma (*Aphanisma blitoides*) RPR 1B.1
- Island sagebrush (*Artemisia nesiotica*) RPR 4.3
- San Miguel Island milk-vetch (*Astragalus miguelensis*) RPR 4.3
- San Clemente Island milk-vetch (*Astragalus nevinii*) RPR 1B.2
- Coulter's saltbush (*Atriplex coulteri*) RPR 1B.2
- South Coast saltscale (*Atriplex pacifica*) RPR 1B.2
- Golden-spined cereus (*Bergerocactus emoryi*) RPR 2.2
- Island morning-glory (*Calystegia macrostegia* ssp. *amplissima*) RPR 4.3
- San Clemente Island evening-primrose (*Camissonia guadalupensis* ssp. *clementina*) RPR 1B.2
- San Clemente Island paintbrush (*Castilleja grisea*) FE, SE, RPR 1B.2
- Island ceanothus (*Ceanothus megacarpus* var. *insularis*) RPR 4.3
- Seaside cistanthe (*Cistanthe maritima*) RPR 4.2
- Nevin's woolly sunflower (*Constancea nevinii*) RPR 1B.3
- Small-flowered morning-glory (*Convolvulus simulans*) RPR 4.2

¹⁴ Status abbreviations: ABC: American Bird Conservancy Green List; BLMS: Bureau of Land Management Sensitive Species; CDF: California Department of Forestry and Fire Protection Sensitive Species; CDFW: California Department of Fish and Wildlife; FD: Federally delisted; FE: Federally listed as Endangered; FSS: USDA Forest Service Sensitive Species; FT: Federally listed as Threatened; RPR: Rare Plant Rank; SE: State-listed as Endangered; SD: State delisted; SSC: CDFW Species of Special Concern; WBWG: Western Bat Working Group: High, Medium or Low priority

DRAFT

- Catalina crossosoma (*Crossosoma californicum*) RPR 1B.2
- Trask's cryptantha (*Cryptantha traskiae*) RPR 1B.1
- Island tarplant (*Deinandra clementina*) RPR 4.3
- San Clemente Island larkspur (*Delphinium variegatum* ssp. *kinkiense*) FE, SE, RPR 1B.1
- Thorne's royal larkspur (*Delphinium variegatum* ssp. *thornei*) RPR 1B.1
- South island bush-poppy (*Dendromecon harfordii* var. *rhamnoides*) RPR 1B.1
- Bright green dudleya (*Dudleya virens* ssp. *virens*) RPR 1B.2
- San Clemente Island buckwheat (*Eriogonum giganteum* var. *formosum*) RPR 1B.2
- Island buckwheat (*Eriogonum grande* var. *grande*) RPR 4.2
- Island poppy (*Eschscholzia ramosa*) RPR 4.3
- Cliff spurge (*Euphorbia misera*) RPR 2.2
- San Clemente Island bedstraw (*Galium catalinense* ssp. *acrispum*) SE, RPR 1B.2
- Showy island snapdragon (*Gambelia speciosa*) RPR 1B.2
- Nevin's gilia (*Gilia nevinii*) RPR 4.3
- San Clemente Island hazardia (*Hazardia cana*) RPR 1B.2
- Decumbent goldenbush (*Isocoma menziesii* var. *decumbens*) RPR 1B.2
- Island jepsonia (*Jepsonia malvifolia*) RPR 4.2
- Southern island mallow (*Lavatera assurgentiflora* ssp. *glabra*) RPR 1B.1
- Robinson's pepper-grass (*Lepidium virginicum* var. *robinsonii*) RPR 1B.2
- Pygmy leptosiphon (*Leptosiphon pygmaeus* ssp. *pygmaeus*) RPR 1B.2
- San Clemente Island woodland star (*Lithophragma maximum*) FE, SE, RPR 1B.1
- San Nicolas Island lomatium (*Lomatium insulare*) RPR 1B.2
- Guadalupe Island lupine (*Lupinus guadalupensis*) RPR 1B.2
- Santa Catalina Island desert-thorn (*Lycium brevipes* var. *hassei*) RPR 1B.1
- California box-thorn (*Lycium californicum*) RPR 4.2
- Santa Cruz Island ironwood (*Lyonothamnus floribundus* ssp. *aspleniifolius*) RPR 1B.2

DRAFT

- San Clemente Island bush-mallow (*Malacothamnus clementinus*) FE, SE, RPR 1B.1
- Leafy malacothrix (*Malacothrix foliosa* ssp. *foliosa*) RPR 4.2
- Dunedelion (*Malacothrix incana*) RPR 4.3
- Small-flowered microseris (*Microseris douglasii* ssp. *platycarpha*) RPR 4.2
- Island bush monkeyflower (*Mimulus flemingii*) RPR 4.3
- Blair's munzothamnus (*Munzothamnus blairii*) RPR 1B.2
- Mud nama (*Nama stenocarpum*) RPR 2B.2
- Many-flowered phacelia (*Phacelia floribunda*) RPR 1B.2
- Island oak (*Quercus tomentella*) RPR 4.2
- Island redberry (*Rhamnus pirifolia*) RPR 4.2
- Santa Catalina figwort (*Scrophularia villosa*) RPR 1B.2
- Santa Cruz Island rock cress (*Sibara filifolia*) FE, RPR 1B.1
- Woolly seablite (*Suaeda taxifolia*) RPR 4.2
- Southern island clover (*Trifolium palmeri*) RPR 4.2
- San Clemente Island brodiaea (*Brodiaea kinkiensis*) RPR 1B.2
- California dissanthelium (*Dissanthelium californicum*) RPR 1B.2
- Vernal barley (*Hordeum intercedens*) RPR 3.2
- Appressed muhly (*Muhlenbergia appressa*) RPR 2.2
- San Clemente Island triteleia (*Triteleia clementina*) RPR 1B.2

The following special-status animal species are reported or are likely to be present on San Clemente Island based on habitat requirements and known range attributes:

- San Clemente Islandsnail (*Micrarionta gabbi*) CDFW Special Animals List
- San Clemente Island blunt-top snail (*Sterkia clementina*) CDFW Special Animals List
- Horseshoe snail (*Xerarionta intercosa*) CDFW Special Animals List
- Senile tiger beetle (*Cicindela senilis frosti*) CDFW Special Animals List
- Island night lizard (*Xantusia riversiana*) FT

DRAFT

- San Clemente sage sparrow (*Amphispiza belli clementeae*) FT, BCC, SSC, ABC
- Western snowy plover (*Charadrius alexandrinus nivosus*) FT, BCC, SSC, ABC, AWL, USBC
- San Clemente loggerhead shrike (*Lanius ludovicianus mearnsi*) FE, SSC
- Ashy storm-petrel (*Oceanodroma homochroa*) BCC, SSC, ABC
- Xantus' murrelet (*Synthliboramphus hypoleucus*) FC, BCC, ST, ABC
- San Clemente Island fox (*Urocyon littoralis clementae*) ST

Santa Catalina Island

Santa Catalina Island, part of the Channel Islands chain, is approximately 21 miles long and 8 miles wide. The Island consists of two parts connected by a low-lying isthmus at Two Harbors. The larger (southeastern) portion can be generally characterized by rolling hills with a gradual descent into the sea. The smaller (northwestern) portion is extremely steep and rugged with steep shoreline palisades. Level terrain on the Island is limited to the floors of a few large coastal canyons and areas, such as Avalon, Pebbly Beach, White's Landing, Middle Ranch, Two Harbors, and Emerald Bay. Mount Orizaba, which is located in the central portion of the Island, represents the highest peak at 2,069 feet above mean sea level (MSL).

The climate of the Island is similar to the mainland with wet mild winters and long dry periods. The majority of the Island is relatively undisturbed, consisting of grasslands, coastal sage scrub, woodlands, and chaparral. Disturbed areas include minor camping areas, paved roads, dirt roads, radio tower pads, reservoirs and a landfill.

Because of habitation, proximity to Los Angeles, the work of the Catalina Island Conservancy, and the Wrigley Marine Science Center, the Island has been studied more than any of the other California offshore islands, with national, state, and County recognition in many ways. Nationally, all of the many offshore rocks and islets within 12 nautical miles of the coast are part of the California Coastal National Monument, which is managed by the Bureau of Land Management in the U.S. Department of the Interior. Of the 64 State-recognized sensitive plants, 3 are federally endangered and 1 is federally threatened. Two vertebrates have federal status: the federal candidate marine bird Xantus' murrelet (*Synthliboramphus hypoleucus*) and the federally-endangered Santa Catalina Island fox (*Urocyon littoralis catalinae*). The fox has recently recovered from an epidemic of distemper by a combination of the luck, in that the northwestern population was so isolated and hardly suffered, and by managed captive breeding.

The State Water Resources Control Board has named a number of areas as ASBS (Area of Special Biological Significance) for the water quality and the coincident diversity of the marine resources. These are the Northwest Santa Catalina Island ASBS #25 (20.9 miles of coastline from Isthmus Cove to Catalina Head), the Southeast Santa Catalina Island ASBS #28 (2.9 miles centered on Seal Rocks), the Western Santa Catalina ASBS #26 (4 miles of coastline from Little Harbor south to Ben Weston Beach), and the Farnsworth Bank ASBS #27 (37 acres on a submerged pinnacle, about 60 feet deep at the top, and covered with many kinds of encrusting organisms, such as the purple hydrocoral *Allopora californica*).

The State has also designated a number of State Marine Conservation Areas (SMCA) and three *de facto* State Marine Reserves (SMR) around the Island. In these conservation areas, the take method

DRAFT

and what may be fished is limited very specifically and differs for each area. (1) Centered on Farnsworth Bank is the Farnsworth Offshore SMCA that extends from 1 to 3 nautical miles offshore and (2) the Farnsworth Onshore SMCA that extends from the coastline to meet the Farnsworth Offshore SMCA about 1 nautical mile. (3) All of CatHarbor on the south side of the Isthmus is in Cat Harbor SMCA with take restricted to recreational fishing for marine aquatic plants and finfish. (4) Arrow Point to Lion Head Point SMCA extends from mean high tide out to 1000 feet offshore. (5) Bird Rock SMCA is centered on Bird Rock off the northwest coast and extends from the offshore boundary of Blue Cavern SMCA to 3 nautical miles offshore, with boundaries drawn due N-S off the end points of Blue Cavern SMCA. (6) Blue Cavern SMCA is really a reserve, with no take allowed even of sport fish. (7) Long Point SMR has no take allowed, which is usual for reserves. (8) Casino Point SMCA on the north end of AvalonBay is really a reserve, with no take allowed. (9) Lover's Cove SMCA on the south side of Avalon Bay has common provisions, but includes the unusual one of the allowance of fish feeding (for the viewers on the glass-bottomed boats).

The California Department of Fish and Wildlife maintains a list of sensitive species (all rare and uncommon), which includes those designated federally. There are 64 plants, 5 snails, and 5 vertebrates listed for the Island.

California Audubon has designated the Island and its surrounding waters to about 0.8 miles offshore as part of the Northern Channel Islands Globally Important Bird Area (IBA).

The County requires special biological review of development on the Island if the project involves a parcel that has a designated Significant Ecological Area (SEA). These areas were all specified on the basis of significant island vegetation, and in a few cases, included marine algae. There are 37 designated SEAs on the Island: Arrow Point; Avalon Canyon; Toyon Canyon; Ben Weston Beach-Mills Landing-Sentinel Rock; Bird Rock; Black Point, Black Jack Mountain and Echo Lake; Blue Cavern Point-Fishermans Cove; Buffalo Springs Reservoirs; Bulrush Canyon; Cactus Peak; Cape Canyon; Cherry Valley; Cottonwood Canyon; Descanso Canyon; Fourth of July Cove; Gallagher Canyon; Geiger Coves; Haypress Area-Hamilton Canyon; Indian Rock; Isthmus; Isthmus Canyon; Johnsons Landing; Little Harbor-Shark Harbor-Indian Head Point; Middle Ranch Canyon; Mount Orizaba; Parsons Landing; Pebbly Beach Canyon; Renton Mine Road; Silver Peak; Skull Canyon; Sweetwater Canyon; White Cove; and Wild Boar Gully.

Vegetation

Vegetation on the Island is composed of a large variety of plant community types. The rugged topography, steep and rocky shoreline, and generally undisturbed condition of the Island has produced a unique diversity of vegetative communities. Historically, the Island was mostly brushland dominated by chamise (*Adenostoma fasciculatum*) and island ceanothus (*Ceanothus megacarpus* var. *insularis*) on the northern slopes, and California sagebrush (*Artemisia californica*) and Santa Catalina Island buckwheat (*Eriogonum giganteum* var. *giganteum*) on the south-facing slopes. Following the introduction of feral herbivores (goats, pigs, deer, and bison), this brushland was replaced in most areas by island scrub oak (*Quercus pacifica*), laurel sumac (*Malosma laurina*), toyon (*Heteromeles arbutifolia*), lemonadeberry (*Rhus integrifolia*), black sage (*Salvia mellifera*), and white sage (*S. apiana*), which dominate today. The lack of a significant fire history and minimal differences in vegetation along elevation gradients (due to an abundance of moisture) has resulted in slope orientation as a major determinant for species presence/absence.

Plant communities on the Island were classified using standard methodology and terminology. Most of the communities discussed correspond directly with those listed in Holland's Preliminary

DRAFT

Descriptions of the Terrestrial Natural Communities of California (1986 and 1992 update). Other communities are named based on dominant species within them and/or commonly used terminology. Descriptions and general locations of each plant community on the Island, including maritime succulent scrub, southern coastal bluff scrub, island chaparral, island oak woodland, ironwood woodland, island cherry woodland, non-native grassland, native grassland, and disturbed are given below.

In recent years, ecologists have refined previous vegetation classification approaches to define communities primarily by their constituent plant assemblages, and have now widely adopted the classification system described by Sawyer, Keeler-Wolf and Evens in *A Manual of California Vegetation*, Second Edition 2009 (MCV). This has become the accepted standard recognized by the California Department of Fish and Wildlife, the California Native Plant Society and the United States Fish and Wildlife Service.

The important difference between current and earlier methods is that earlier classifications were based on a variety of factors, such as physiographic features, as in the case of vernal pools; or by specific plants, as in the case of coast live oak woodland; or by the use of commonly accepted terms, as in chaparral. In the MCV, plant communities are defined with more precision as botanical alliances where one, or occasionally two, plant species are dominant or co-dominant with a host of other possible associated plants. The MCV lists no one plant community called chaparral because in habitats of this type any one of a variety of shrubs can be dominant and influence the character of the vegetation. For example, in a location where chamise (*Adenostoma fasciculatum*) is predominant it is classified as an *Adenostoma fasciculatum* Shrubland Alliance, while close by greenbark ceanothus (*Ceanothus spinosus*) may emerge as the most common shrub and this is termed a *Ceanothus spinosus* Shrubland Alliance.

The transition to the new MCV format is ongoing. Not all alliances have been fully described and new ones are still being recognized. As such, a significant amount of field work and site surveying for accuracy would be needed to verify all existing alliances on the Island, as well as extensive research to define every alliance in the new MCV plant communities' format. The more familiar nomenclatures will continue to be used for the Island when appropriate.

An effort has been made to conform to this new format. Descriptions and general locations of each plant community as described in the California Channel Islands Species Recovery Plan (USFWS, 1984) appear below. The plant communities correspond to classifications used in that document. Each was considered baseline information and evaluated for the potential presence of alliances as described in the MCV. Alliances with matching profiles of given criteria are listed. In many cases only with further investigation can the presence of some alliances be confirmed. In addition, it should be noted that not all alliances are listed within this description, as many alliances have yet to be defined and new alliances are still being discovered. Descriptions and general locations of the each plant community present on the Island are given below.

Maritime Succulent Scrub: A low, open scrub of soft-leaved shrubs and herbs with a rich admixture of stem and leaf succulents occurring on steep coastal slopes. This community is dominated by California sagebrush and coastal prickly-pear (*Opuntia littoralis*), which is located mainly on the exposed, dry south-facing slopes typically with well-drained soils. Other species associated with this community include Catalina crossosoma (*Crossosoma californica*), California brittle-bush (*Encelia californica*), Santa Catalina Island buckwheat, bedstraw (*Galium* spp.), island broom (*Acmispon dendroideus* var. *dendroideus*), laurel sumac, lemonadeberry, and black sage.

DRAFT

Corresponding MCV communities:

- *Artemisia californica* (California sagebrush scrub) Shrubland Alliance
- *Opuntia littoralis* (coast prickly pear scrub) Shrubland Alliance
- *Encelia californica* (California brittle bush scrub) Shrubland Alliance
- *Deinandra clementina-Eriogonum giganteum* (island buckwheat-island tarplant scrub) Provisional Shrubland Alliance
- *Malosma laurina* (laurel sumac scrub) Shrubland Alliance
- *Rhus integrifolia* (lemonade berry scrub) Shrubland Alliance
- *Salvia mellifera* (black sage scrub) Shrubland Alliance

Southern Coastal Bluff Scrub: A low scrub community adapted to exposed areas with nearly constant winds and high salt content. It consists of the largest reservoir of sensitive species and island endemics due to its location within inaccessible areas. This community is dominated by giant coreopsis (*Leptosyne gigantea*), Catalina crossosoma, *Dudleya* spp., Santa Catalina Island buckwheat, and island tarplant (*Deinandra clementina*) (David Carroll and Associates (DCA), 1994). Southern coastal bluff scrub, which occurs on the precipitous cliff, faces typically near the mouths of canyons and adjacent to some of the Island's disturbed areas on the steep seaward (east-facing) slopes and bluffs.

Corresponding MCV communities:

- *Leptosyne gigantea* (giant coreopsis scrub) Shrubland Alliance
- *Deinandra clementina-Eriogonum giganteum* (island buckwheat-island tarplant scrub) Provisional Shrubland Alliance

Island Chaparral: Consists of tall broad-leaved shrubs that form a dense cover on steep slopes. Dominant species found within this community include island and MacDonal's scrub oaks (*Quercus pacifica* and *Q. macdonaldii*), feltleaf ceanothus (*Ceanothus arboreus*), chamise, island red berry, and Santa Catalina Island manzanita (*Arctostaphylos catalinae*). Island chaparral occupies canyon bottoms, most of the higher elevations, and steep, north-facing slopes.

Corresponding MCV communities:

- *Quercus pacifica* (island scrub oak chaparral) Shrubland Alliance
- *Ceanothus megacarpus* (big pod ceanothus chaparral) Shrubland Alliance
- *Adenostoma fasciculatum* (chamise chaparral) Shrubland Alliance

Island Oak Woodland: Dominated by canyon oak (*Quercus chrysolepis*) with a poorly developed shrub layer, which includes wild blackberry (*Rubus ursinus*), poison oak, heart-leaved penstemon (*Keckiella cordifolia*), and southern chaparral honeysuckle (*Lonicera subspicata* var. *denudata*). Some island oak woodlands along riparian habitat include scattered arroyo willows (*Salix lasiolepis*). This community occurs in relatively moist, protected canyon bottoms with rich alluvial soils.

DRAFT

Corresponding MCV communities:

- *Quercus chrysolepis* (canyon live oak forest) Forest Alliance

Island Ironwood Forest: An upland community characterized by a dominance of Catalina ironwood (*Lyonothamnus floribundus* ssp. *floribundus*). The Island endemic is a broad-leaved tree and occurs in groves of 50-100 trees located along the north- and east-facing slopes (DCA 1994). Other species occasionally associated with the ironwood forest include island scrub oak and Santa Catalina Island manzanita. The understory is sparse, supporting a number of herbaceous annuals and ferns. This community is typically found in protected canyons with rich alluvial soils in the northern portion of the island.

Corresponding MCV communities:

- *Lyonothamnus floribundus* (Catalina ironwood groves) Woodland Special Stands

Island Cherry Woodland: An open, dense woodland dominated by Catalina cherry (*Prunus ilicifolia* ssp. *lyonii*) with an understory consisting of Santa Catalina figwort (*Scrophularia villosa*), cudweed (*Gnaphalium* spp.), common chickweed (*Cerastium glomeratum*), wild cucumber (*Marah macrocarpa*), Santa Catalina Island bushmallow (*Malacothamnus fasciculatus* var. *catalinensis*), island morning-glory (*Calystegia macrostegia* ssp. *amplissima*), and many of weedy forb and grass species. This community occurs mostly along riparian habitats and in valley/canyon bottoms in the northern portion of the island.

Corresponding MCV communities:

- *Prunus ilicifolia* (Holly leaf cherry chaparral) Shrubland Alliance

Grassland: Consist of low, herbaceous vegetation that are dominated by grasses, but generally also harbor native forbs and bulbs as well as naturalized annual forbs. Non-native grassland consists of dominant invasive annual grasses that are primarily of Mediterranean origin. Dominant species found within this community include slender oats (*Avena barbata*), wild oats (*A. fatua*), ripgut brome (*Bromus diandrus*), red brome (*B. madritensis* ssp. *rubens*), and wild mustards (*Brassica*, *Hirschfeldia*, and *Sisymbrium* spp.). Non-native grasslands are located in small patches throughout the island; along many of the ridges and gentle slopes with shallow clay or clay loam substrates; and in more significant acreage, on rolling hills in the southeastern portion of the island. Native grassland consists of at least 10 percent cover of native grass species with the remaining coverage similar to non-native grasslands. Small patches of native grassland can be found on the island mostly mixed with non-native grasslands.

Corresponding MCV communities:

- *Avena (barbata, fatua)* (Wild oats grasslands) Semi-Natural Herbaceous Stands
- *Bromus (diandrus, hordeaceus) -Brachypodium distachyon* (Annual brome grasslands) Semi-Natural Herbaceous Stands
- *Bromus rubens-Schismus (arabicus, barbatus)* (Red brome or Mediterranean grass grasslands) Semi-Natural Herbaceous Stands
- *Brassica nigra* and other mustards (Upland mustards) Semi-Natural Herbaceous Stands

DRAFT

Disturbed Areas: Areas that either completely lack vegetation or are dominated by ruderal species within developed areas. Vegetation typically include horehound (*Marrubium vulgare*) and tree tobacco (*Nicotiana glauca*). Several disturbed areas occur throughout the Island and take the form of residential developments, paved roads, fire breaks, dirt access roads, trails, and other similarly disturbed areas.

Corresponding MCV communities: None at this time.

Wildlife

Wildlife on the Island is diverse and abundant due to the large acreage of natural open space and the diversity of habitat types. While a few wildlife species are entirely dependent on a single vegetative community, the vegetation communities within the area and adjoining areas constitute a functional ecosystem for a variety of wildlife species.

The analysis of invertebrates is severely limited due to the lack of data. The Island, however, supports healthy populations of a diverse assortment of countless invertebrate species. Amphibian populations are generally abundant and diverse due to the high moisture content provided under the shade of woodlands and the abundance of drainages. Many essential reptilian habitat characteristics are present. These include open habitats that allow free movement and high visibility and small mammal burrows for cover and escape from predators and extreme weather. These characteristics as well as the variety of habitat types present are likely to support a wide variety of reptilian species.

The scrubland, woodland, riparian, and grassland habitats provide foraging and cover habitat for year-round residents, seasonal residents, and migrating song birds. In addition, several year-round water sources and abundant raptor foraging, perching, and nesting habitat are located throughout the area. The combination of these resources as well as the confluence of many community types provides for a high diversity of bird species. Unlike other taxonomic groups, mammal populations are diverse and reflective of the unique island habitat types.

Wildlife Movement

Wildlife movement occurs throughout the Island. Concentrated movement corridors or bottlenecks are uncommon on the Island due to the abundance of uninterrupted open space and the lack of disturbed areas. In general, movement takes place in large drainages, along ridgelines, and along dirt roads. However, the small isthmus at Two Harbors represents a significant reduction in the ability for animals to move freely between the two parts of the Island. Movement across the isthmus has been further restricted by human encroachment of the Two Harbors community and Island visitors. Although a lack of movement across the isthmus may isolate some animal populations and reduce the genetic diversity on either side, this division has provided a unique opportunity for restoration by isolating and removing feral animals from the Island.

Sensitive Biological Resources

The statuses of rare plants are hierarchically categorized by the CNPS using a rank and decimal system. The initial category level of Rare Plant Rank is indicated by ranks 1A (presumed extinct), 1B (rare or endangered in California and elsewhere), 2 (rare or endangered in California but more common elsewhere), 3 (more information needed, a review list), and 4 (limited distribution). In cases where the CNPS has further identified the specific threat to the species, a decimal or Threat Code is added: .1 (seriously endangered in California), .2 (fairly endangered in California) or .3 (not very endangered in California).

DRAFT

The following special-status plant taxa have been reported or have the potential to occur on Santa Catalina Island, based on known habitat requirements and geographic range information:

- Baja rock lichen (*Graphis saxorum*) CDFW Special Plants List
- Red sand-verbena (*Abronia maritima*) RPR 4.2
- Island broom (*Acemispom dendroideus* var. *dendroideus*) RPR 4.2
- Aphanisma (*Aphanisma blitoides*) RPR 1B.1
- Santa Catalina Island manzanita (*Arctostaphylos catalinae*) RPR 1B.2
- Coulter's saltbush (*Atriplex coulteri*) RPR 1B.2
- South Coast saltscale (*Atriplex pacifica*) RPR 1B.2
- Davidson's saltscale (*Atriplex serenana* var. *davidsonii*) RPR 1B.2
- Golden-spined cereus (*Bergerocactus emoryi*) RPR 2.2
- Round-leaved filaree (*California macrophylla*) RPR 1B.1
- Island ceanothus (*Ceanothus megacarpus* var. *insularis*) RPR 4.3
- Southern tarplant (*Centromadia parryi* ssp. *australis*) RPR 1B.1
- Island mountain-mahogany (*Cercocarpus betuloides* var. *blancheae*) RPR 4.3
- Catalina Island mountain-mahogany (*Cercocarpus traskiae*) FE, SE, RPR 1B.1
- Seaside cistanthe (*Cistanthe maritima*) RPR 4.2
- Nevin's woolly sunflower (*Constancea nevinii*) RPR 1B.3
- Small-flowered morning-glory (*Convolvulus simulans*) RPR 4.2
- Catalina crossosoma (*Crossosoma californicum*) RPR 1B.2
- Island tarplant (*Deinandra clementina*) RPR 4.3
- South island bush-poppy (*Dendromecon harfordii* var. *ramnoides*) RPR 1B.1
- Western dichondra (*Dichondra occidentalis*) RPR 4.2
- Beach spectaclepod (*Dithyrea maritima*) ST, 1B.1
- Greene's dudleya (*Dudleya greenei*) RPR 4.2
- Catalina Island dudleya (*Dudleya virens* ssp. *hassei*) RPR 1B.2

DRAFT

- Island green dudleya (*Dudleya virens* ssp. *insularis*) RPR 1B.2
- Bright green dudleya (*Dudleya virens* ssp. *virens*) RPR 1B.2
- Santa Catalina Island buckwheat (*Eriogonum giganteum* var. *giganteum*) RPR 4.3
- Island buckwheat (*Eriogonum grande* var. *grande*) RPR 4.2
- Island poppy (*Eschscholzia ramosa*) RPR 4.3
- Cliff spurge (*Euphorbia misera*) RPR 2.2
- Santa Catalina Island bedstraw (*Galium catalinense* ssp. *catalinense*) RPR 1B.2
- Nuttall's island bedstraw (*Galium nuttallii* ssp. *insulare*) RPR 4.3
- Showy island snapdragon (*Gambelia speciosa*) RPR 1B.2
- Nevin's gilia (*Gilia nevinii*) RPR 4.3
- Palmer's grapplinghook (*Harpagonella palmeri*) RPR 4.2
- Island rush-rose (*Helianthemum greenei*) FT, RPR 1B.2
- Decumbent goldenbush (*Isocoma menziesii* var. *decumbens*) RPR 1B.2
- Island jepsonia (*Jepsonia malvifolia*) RPR 4.2
- Southern island mallow (*Lavatera assurgentiflora* ssp. *glabra*) RPR 1B.1
- Fragrant pitcher sage (*Lepechinia fragrans*) RPR 4.2
- Santa Barbara honeysuckle (*Lonicera subspicata* var. *subspicata*) RPR 1B.2
- Santa Catalina Island desert-thorn (*Lycium brevipes* var. *hassei*) RPR 1B.1
- California box-thorn (*Lycium californicum*) RPR 4.2
- Santa Catalina Island ironwood (*Lyonothamnus floribundus* ssp. *floribundus*) RPR 1B.2
- Small-flowered microseris (*Microseris douglasii* ssp. *platycarpha*) RPR 4.2
- Santa Catalina Island monkeyflower (*Mimulus traskiae*) RPR 1A
- Coast woolly-heads (*Nemacaulis denudata* var. *denudata*) RPR 1B.2
- Short-lobed broomrape (*Orobanche parishii* ssp. *brachyloba*) RPR 4.2
- Lyon's pentachaeta (*Pentachaeta lyonii*) FE, SE, RPR 1B.1
- Engelmann oak (*Quercus engelmannii*) RPR 4.2

DRAFT

- Island scrub oak (*Quercus pacifica*) RPR 4.2
- Island oak (*Quercus tomentella*) RPR 4.2
- Island redberry (*Rhamnus pirifolia*) RPR 4.2
- Santa Catalina Island currant (*Ribes viburnifolium*) RPR 1B.2
- Santa Catalina figwort (*Scrophularia villosa*) RPR 1B.2
- Chaparral ragwort (*Senecio aphanactis*) RPR 2B.2
- Santa Cruz Island rock cress (*Sibara filifolia*) FE, RPR 1B.1
- Wallace's nightshade (*Solanum wallacei*) RPR 1B.1
- Woolly seablite (*Suaeda taxifolia*) RPR 4.2
- Southern island clover (*Trifolium palmeri*) RPR 4.2
- Catalina mariposa lily (*Calochortus catalinae*) RPR 4.2
- California dissanthelium (*Dissanthelium californicum*) RPR 1B.2
- Vernal barley (*Hordeum intercedens*) RPR 3.2
- Chaparral rein orchid (*Piperia cooperi*) RPR 4.2

The following special-status animal species are reported or are likely to be present on Santa Catalina Island based on habitat requirements and known range attributes:

- Santa Catalina lancetooth (*Haplotrema catalinense*) CDFW Special Animals List
- Shepard's snail (*Pristiloma shepardae*) CDFW Special Animals List
- Catalina mountainsnail (*Radiocentrum avalonense*) CDFW Special Animals List
- San Clemente Island blunt-top snail (*Sterkia clementina*) CDFW Special Animals List
- Sandy beach tiger beetle (*Cicindela hirticollis gravida*) CDFW Special Animals List
- Santa Catalina garter snake (*Thamnophis hammondi* ssp.) CDFW Special Animals List
- Bald eagle (*Haliaeetus leucocephalus*) SE, CDF, CDFW Fully Protected
- Xantus' murrelet (*Synthliboramphus hypoleucus*) FC, BCC, ST, ABC
- Santa Catalina Island shrew (*Sorex ornatus willetti*) SSC
- Santa Catalina Island fox (*Urocyon littoralis catalinae*) FE, ST

Santa Monica Mountains

For a description of the resources in the Santa Monica Mountains, please refer to the description for the Santa Monica Mountains Significant Ecological Area (SEA) in Section IV.

VI. Watersheds

Antelope Valley Watershed

The southern half of the Lahontan hydrologic region is located in the Antelope Valley. Unlike the coastal watersheds in Los Angeles County, it is a closed basin on the edge of the Mojave Desert, having no outlet to the ocean or major river system. Numerous streams drain the north-facing San Gabriel Mountains, carrying rainfall and snow melt from the Angeles National Forest into the Antelope Valley. Significant stream systems in the Antelope Valley are Amaroosa Creek, Big Rock Creek, and Little Rock Creek.

During most years, the rainfall in the Antelope Valley is scant, averaging less than eight inches per year. Every few years, major storms cause flooding, sending sheets of water flow across the eastern portion of the Antelope Valley to the dry lakebeds of Rosamond and Rodgers lakes in Kern County. Uninhibited by development, the sheet flow filters into the groundwater basin or evaporates on the lakebeds, leaving the surface smooth and flat. This natural runoff process is important for two reasons: 1) it benefits the local communities with groundwater recharge; and 2) it seasonally resurfaces the dry lake beds, which are used for aircraft landings at Edwards Air Force Base.

The Lahontan Regional Water Quality Control Board monitors the Antelope Valley watershed through its Basin Plan for the region. The Basin Plan calls for land use controls to help reduce pollutants in stormwater runoff. In particular, the Basin Plan advocates for limiting impervious surfaces, restoring natural vegetation and protecting the headwaters of stream channels and riparian areas.

Los Angeles River Watershed

The Los Angeles River watershed covers approximately 870 square miles, a small part of which extends into Ventura County. It includes the San Fernando Valley and is the largest watershed in the Los Angeles Basin. The river extends 51 stream miles, from the confluence of Bell Creek and Arroyo Calabasas, to the Pacific Ocean. The first 32 miles of the river flow through the cities of Los Angeles, Burbank, and Glendale, and then, subsequently, through Vernon, Commerce, Maywood, Bell, Bell Gardens, Lynwood, Compton, South Gate, Paramount, Cudahy, and Long Beach. Numerous tributaries feed the Los Angeles River, as it flows through the San Fernando Valley and the coastal plain to the Long Beach Harbor. These tributaries include Tujunga Wash, Verdugo Wash, Arroyo Seco, Rio Hondo, and Compton Creek. Several important biotic communities exist in the northern tributaries that feed the river, including freshwater marsh areas in Tujunga Canyon and the Hansen Flood Control Basin. The natural habitat in these tributaries provides a semi-protected corridor for wildlife between the Angeles National Forest, Santa Monica Mountains National Area, and the Los Angeles River.

By 1960, the Los Angeles River was lined with concrete along most of its length by the U.S. Army Corps of Engineers in order to prevent the loss of lives and property from flood damage. As a result, the Los Angeles River's sole purpose for years was efficient water conveyance—carrying stormwater from the land to the ocean as quickly as possible. Efforts continue under the auspices of the Los Angeles County Flood Control District to capture as much stormwater as possible and redirect it to regional groundwater recharge areas to replenish groundwater basins, saving thousands of acre-feet

of water every year.

The volume of pollutants that enters the Los Angeles River is extremely high due to accumulated urban stormwater runoff from the hundreds of square miles of impervious land uses that flank the Los Angeles River. To address these problems, the County, the Flood Control District, local jurisdictions, a variety of stakeholders, and the Los Angeles Regional Water Quality Control Board are implementing programs to reduce the number and concentration of pollutants that enter the Los Angeles River.

Over the past two decades, interest in the Los Angeles River's recreational and ecological functions has reemerged, culminating in a riverwide planning effort in the 1990s, which resulted in the adoption of the *Los Angeles River Master Plan* by the Board of Supervisors in 1996. The Plan was created through a cooperative effort by the County and many river stakeholder groups for the enhancement of aesthetic, recreational, flood protection and environmental functions of the Los Angeles River. The Plan seeks to do so by expanding bikeway, walking and equestrian trails to and along the Los Angeles River, enhancing existing trails and habitat with landscaping, and promoting economic development opportunities. Since the adoption of the Plan, an advisory committee has overseen many new projects, including bike trails, pocket parks, equestrian trail enhancements, art and signage. So much public interest in the river has been generated that many more improvements are anticipated in the future. The County's Bicycle Master Plan also prioritizes the Los Angeles River bike path.

The County is also working with various organizations and agencies that are involved in watershed-related planning activities, such as the San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy, the Council for Watershed Health, and the Flood Control District. The attention being paid to the watershed has resulted in a better understanding of its functions and generated an unprecedented network of residents, private organizations and government entities dedicated to watershed management. The County has also partnered with the City of Los Angeles on implementation of its 2007 Los Angeles River Revitalization Master Plan. Subsequently, the County Board of Supervisors and Los Angeles City Council adopted the Los Angeles River Memorandum of Understanding, which established the Los Angeles River Cooperation Committee to prioritize cooperative implementation of Los Angeles River projects. In addition, the County is a partner in the U.S. Army Corps of Engineers' Los Angeles River Ecosystem Restoration Feasibility Study (started in 2006 for which the City of Los Angeles is serving as primary local sponsor). The County is also a partner with the U.S. Bureau of Reclamation on the Los Angeles Basin Study to prioritize stormwater capture and infiltration that will result in watershed-wide conservation.

Compton Creek Sub-Watershed

Compton Creek drains an area of approximately 42 square miles, including portions of the cities of Carson, Compton, Long Beach, Los Angeles, Lynwood, and South Gate and portions of the unincorporated communities of Florence-Firestone, Willowbrook, and Rancho Dominguez. Compton Creek drains into the Los Angeles River, which then empties into San Pedro Bay at the eastern edge of Long Beach Harbor.

With more than 700,000 residents, the Compton Creek watershed is highly urbanized and most of its drainage courses are lined with concrete. Approximately 97 percent of the land area is occupied by homes, businesses, roads, and similar uses. As a result of excess levels of trash and coliform bacteria, surface and groundwater quality in Compton Creek has been degraded, natural hydrologic functions modified, and plant and wildlife diversity reduced.

In recent years, various groups, cities, and agencies have worked to transform Compton Creek into a

valued community asset, improve and expand open space, optimize water resources, preserve and restore habitat, and create a network of trails and bike paths. Some of these efforts have been informally coordinated, in recognition of the potential to extend benefits beyond the borders of individual cities, create opportunities to leverage benefits, and maximize funding resources.

Los Angeles Harbor Watershed

The major river system within the area that drains to Los Angeles Harbor is the Dominguez Channel. The Dominguez Channel watershed, part of the larger Los Angeles Harbor watershed, is located within the southern portion of Los Angeles County and encompasses approximately 110 square miles of intensely urban area. Approximately 81 percent of the watershed is developed. Residential development covers nearly 40 percent of the watershed, and another 41 percent is made up by industrial, commercial and transportation uses. With a population of nearly 1 million, considerable demands are made on infrastructure and services within the watershed. Local water supply is limited and the majority of water use is provided by imported sources. Parkland and open space are in short supply and generally are deficient. Another significant stream system within the Los Angeles Harbor watershed is the Wilmington Drain.

There are significant stormwater pollution issues in this watershed. For example, old waste disposal practices have left DDT and PCBs deposited in the channel bottom, which are carried to the harbor in land and aquatic sediment swept up and re-suspended by stormwater.

Nine unincorporated areas are located within this urban-industrial watershed and each of these areas affects and is affected by the health and function of Dominguez Channel and its tributaries.

San Gabriel River Watershed

The San Gabriel River watershed encompasses part of the Angeles National Forest, the San Gabriel Valley, and large urban areas in southeast portion of Los Angeles County. It is bounded by the Los Angeles River on much of its western flank, and extends to San Bernardino and Orange counties. Totalling more than 640 square miles, the watershed has extensive areas of un-channeled tributaries, which support riparian and woodland habitats. Its northern reaches in the Angeles National Forest are dramatically different from the developed 167 square miles in the Los Angeles Basin. The U.S. Congress has preserved two wilderness areas within this watershed: the San Gabriel Wilderness Area, 36,215 acres along the west fork of the San Gabriel River, and Sheep Mountain Wilderness Area, 31,680 acres along the east fork of the San Gabriel River.

The main watercourse in this watershed is the San Gabriel River. The San Gabriel River extends 59 stream miles from the Angeles National Forest to the Pacific Ocean, draining 350 square miles of land. It also recharges groundwater tables in several basins. The major tributaries that feed the San Gabriel River include Coyote Creek, Walnut Creek, Puente Creek and San Jose Creek. The upper section of the San Gabriel River and its tributaries are still considered relatively pristine. However, intensive recreational use and erosion due to wildfires in this area may threaten water quality and wildlife that depend on the river. The middle section of the river has been extensively modified throughout the San Gabriel Valley to diminish flood damage and encourage groundwater recharge. The lower section, similar to the Los Angeles River, is lined with concrete from Firestone Boulevard to the bay. In contrast to the upper and middle sections of the river, dry weatherflow in the lower section stems primarily from urban runoff and treated effluent from municipal wastewater treatment facilities.

A clear link exists between the health of this watershed and the quality of life for millions of Los Angeles

County residents. The upper reaches of the San Gabriel River support wildlife, deliver drinking water and provide a myriad of recreational opportunities. To protect and enhance the multiple benefits of this resource a riverwide planning effort entitled *San Gabriel River Master Plan* was adopted in 2006. This effort, spearheaded by the County, brings together a dynamic group of stakeholders, including the 13 cities along the San Gabriel River, residents, environmental groups and many business and community leaders.

The County is working with stakeholders, such as the San Gabriel and Lower Los Angeles Rivers and Mountains Conservancy, the Santa Monica Mountains Conservancy, and the Flood Control District. Together, stakeholders developed a watershed and open space plan in 2001 entitled *Common Ground: From the Mountains to the Sea* that provides general guidelines for improvement of the San Gabriel and Lower Los Angeles Rivers watersheds through community development, public awareness, preservation of open space and creation of recreational opportunities—particularly along the rivers.

Santa Clara River Watershed

The Santa Clara River watershed is an extensive hydrologic system that encompasses the western portion of the Angeles National Forest in Los Angeles County and the eastern portion of Los Padres National Forest in Ventura County. The Santa Clara River—an essential component of this watershed—recharges local groundwater, provides riparian habitat and supplies water to downstream agricultural lands in Ventura County. It is the largest relatively unaltered river system in Southern California, and the single most important natural wildlife corridor in Los Angeles County. The Santa Clara River and its tributaries provide drainage for approximately 654 square miles of the upper watershed within Los Angeles County. The Santa Clara River's major tributaries include Soledad Canyon, Castaic Creek, San Francisquito Canyon Creek, Bouquet Canyon Creek, Sand Canyon Creek, Mint Canyon Creek and Santa Clara River South Fork. Several endangered species are found in this watershed, including the arroyo toad and the unarmored three-spine stickleback. Another important stretch of the Santa Clara River supports a variety of riparian-obligate songbirds and birds of prey between Castaic Junction and Blue Cut near the Ventura County line, where the groundwater basin thins and narrows, forcing groundwater toward the surface.

A link exists between the health of this watershed, particularly its tributaries, and development in the area. Urban expansion in the 1990s and early 2000s impacted the watershed on several levels, including a reduction in local water supplies and disappearing open space. Furthermore, the land use activities in this area have created many square miles of impervious surfaces, which have created more urban runoff and reduced the amount of water that would naturally percolate into groundwater basins. By employing watershed management techniques, the County aims to curb this trend.

Santa Monica Bay Coastal Watersheds

The County, other agencies, cities, and stakeholders coordinate the management of the coastal watersheds of Santa Monica Bay as two distinct management areas, due to the vast differences in land use and population density: North Santa Monica Bay watersheds (North Bay) and the South Santa Monica Bay watersheds (South Bay). The North Bay consists of the Malibu Creek and Topanga Creek watersheds as well as sixteen other rural coastal watersheds. North Bay watersheds are primarily natural open space with low-density residential developments. The South Bay includes Ballona Creek watershed, Marina del Rey, and 10 urban coastal sub-watersheds. Although vastly different from one another, these watersheds have a direct impact on the quality and quantity of water delivered to the Santa Monica Bay. The Malibu Creek watershed traverses a rural mountainous area,

while the Ballona Creek watershed is comprised of intensely urban development.

The coastal watersheds of the Santa Monica Bay extend from the Ventura County-Los Angeles County line to outer Cabrillo Beach in San Pedro. This length includes 44 beaches along 55 miles of coastline, attracting over 55 million beach visitors each year. While the Malibu Creek and Ballona Creek watersheds also drain to Santa Monica Bay, they are typically managed as separate areas due to their significant size.

Ballona Creek Sub-Watershed

The Ballona Creek watershed is located in the western portion of Los Angeles County and is approximately 130 square miles in size. It is highly urbanized and home to more than 1.6 million residents. It includes the cities of Beverly Hills and West Hollywood; portions of the cities of Culver City, Inglewood, Los Angeles, Santa Monica; Caltrans facilities; the Ballona Wetlands, and the unincorporated communities of Marina del Rey, Baldwin Hills, Ladera Heights and a portion of Playa Del Rey. Centinela Creek, Sepulveda Canyon Channel and Benedict Canyon Channel are significant tributaries of Ballona Creek.

Over the years, the urbanization of the Ballona Creek watershed routed many small tributaries through storm drains. These storm drains collect runoff from city streets and carry it to major tributaries and eventually to Ballona Creek, which flows into the Santa Monica Bay. Major contributors to the impaired water quality in Ballona Creek are urban runoff and illegal dumping. These pollutants significantly contribute to pollution in the Santa Monica Bay, degrading ecosystems and recreational opportunities.

The Ballona Creek Trash TMDL became effective on August 11, 2005. To comply with this regulation, the County adopted an aggressive strategy to reduce the amount of trash entering Ballona Creek from unincorporated areas.

Malibu Creek Sub-Watershed

Malibu Creek watershed is the largest rural watershed in North Santa Monica Bay. It is approximately 109 square miles and consists of over 75 percent natural open space. The Malibu Creek watershed encompasses a major portion of the Santa Monica Mountains and is one of many sub-watersheds that drain the mountain range. Over the past twenty years, the number of residents living in the Malibu Creek watershed has doubled. This growth and development has increased runoff, sedimentation and demand for imported water, and caused various tributaries that feed Malibu Creek to be channelized. As a result, the natural flow of water within the watershed has changed, degrading oak and riparian woodlands, steelhead trout populations, and the Malibu Lagoon.

The primary watercourse draining this watershed is Malibu Creek, which flows into Malibu Lagoon. The health and function of Malibu Creek and its tributaries is an important issue as these waterways drain 109 square miles of the watershed into Malibu Lagoon and the Santa Monica Bay—a National Estuary.¹⁵ Two important plant communities comprise the lagoon: the coastal salt marsh and coastal

¹⁵ The Santa Monica Bay is designated a National estuary under the U.S. EPA's National Estuaries Program.

strand, and over 200 species of birds use the lagoon as a refuge.

A clear link exists between the health of Malibu Creek watershed, particularly Malibu Creek, and development in the mountains. Land use activities account for about half of all pollutants that enter the Malibu watershed drainage. Pollution sources include roadway runoff, septic system overflow, new construction, and vegetation clearance.

The Santa Monica Mountains North Area Plan, adopted by the Board of Supervisors in 2000, and the 1986 Local Coastal Plan address the adverse affect of development on the Santa Monica Mountains. These plans, which cover portions of the mountains north of the coastal zone, significantly restrict the potential number of dwelling units that may be built in the mountains. The guiding principle of the plan is to let the land dictate the site and type of development that should be allowed.

Marina del Rey Sub-Watershed

The Marina del Rey watershed is approximately twosquare miles in size and its drainage area is mostly within the City of Los Angeles. The unincorporated marina and harbor were mainly constructed from remnants of the Ballona Creek Wetlands and Estuary.

VI. Agricultural Resources

Agricultural Resource Areas Methodology

Figure 9.5in the Conservation and Natural Resources Element shows the County's Agricultural Resource Areas (ARAs), where the County promotes the preservation of agricultural activities. The ARA boundaries were derived from farmland identified by the State Department of Conservation, including Prime Farmland, Farmland of Statewide Importance, Farmland of Local Importance, and Unique Farmland. In addition, the ARAs include lands that received permits from the Los Angeles County Agricultural Commissioner/Weights and Measures.

To reflect changes in land uses and address environmental concerns, the following were excluded from the ARAs:

- Significant Ecological Areas (SEA);
- Approved specific plan areas;
- Approved large-scale renewable energy facilities;
- Lands outside of the Santa Clarita and Antelope Valleys, where farming is concentrated; and
- Lands that are designated Public and Semi-Public (P).

Data from the U.S. Census of Agriculture

Table E.1: Change in Number and Acreage of Farms in Los Angeles County, 1987-2007

DRAFT

	1987	1992	1997	2002	2007	2017	2022
Farms (number)	2,035	1,446	1,226	1,543	1,734	1,035	766
Change from previous year	-	-589	-220	317	191	-259	-269
Percent change from previous year	-	-28.94%	-15.21%	25.86%	12.38%	-20.02%	-25.99%
Land in farms (acres)	280,156	183,569	130,838	111,458	108,463	57,809	69,224
Change from previous year	-	-96,587	-52,731	-19,380	-2,995	-33,880	+11,415
Percent change from previous year	-	-34.48%	-28.73%	-14.81%	-2.69%	-36.95%	+19.75%

Source: U.S. Census of Agriculture, 1987-~~2007~~ 2022.

DRAFT